(f) Measurements of the magnetic susceptibility of the compound in N_2 and in CO carried out by Theorell & Agner do not in any way contradict the view that the iron of this compound is in the ferrous state.

6. The close relationship between the properties of azide-catalase and the free enzyme suggests that the catalytic decomposition of H_2O_2 by catalase involves a cyclical reduction and oxidation of its iron, as has been established in the case of azide-catalase.

REFERENCES

- Agner, K. (1938). *Biochem. J.* **32**, 1702.
 Agner, K. (1941). *Acta physiol. Scand.* **2**, Suppl. VIII.
 Agner, K. (1942). *Ark. Kemi Min. Geol.* **16 A**, no. 6.
 Coryell, C. D. (1939). *J. phys. Chem.* **43**, 841.
 Dounce, A. L. & Frampton, O. D. (1939). *Science*, **89**, 300.
 Dounce, A. L. & Howland, J. W. (1943). *Science*, **97**, 21.
 Euler, H. v. (1934). *Chemie der Enzyme*. München: Bergmann.
 Euler, H. v. & Josephson, K. (1927). *Liebigs Ann.* **452**, 158.
 Haber, F. & Willstätter, R. (1931). *Ber. dtsh. chem. Ges.* **64**, 2844.
 Keilin, D. (1930). *Proc. roy. Soc. B*, **106**, 418.
 Keilin, D. (1936). *Proc. roy. Soc. B*, **121**, 165.
 Keilin, D. & Hartree, E. F. (1935). *Proc. roy. Soc. B*, **117**, 1.
 Keilin, D. & Hartree, E. F. (1936). *Proc. roy. Soc. B*, **121**, 173.
 Keilin, D. & Hartree, E. F. (1938). *Proc. roy. Soc. B*, **124**, 397.
 Keilin, D. & Hartree, E. F. (1939). *Proc. roy. Soc. B*, **127**, 167.
 Keilin, D. & Hartree, E. F. (1943). *Nature, Lond.*, **152**, 626.
 Kuhn, R. & Wassermann, A. (1933). *Liebigs Ann.* **503**, 203.
 Lemberg, R. & Legge, J. W. (1943). *Biochem. J.* **37**, 117.
 Lemberg, R., Legge, J. W. & Lockwood, W. H. (1941). *Biochem. J.* **35**, 339.
 Lemberg, R., Norrie, M. & Legge, J. W. (1939). *Nature, Lond.*, **144**, 551.
 Pauling, L. (1942). *The Nature of the Chemical Bond*. New York: Cornell University Press.
 Pauling, L. & Coryell, C. D. (1936a). *Proc. nat. Acad. Sci., Wash.*, **22**, 159.
 Pauling, L. & Coryell, C. D. (1936b). *Proc. nat. Acad. Sci., Wash.*, **22**, 210.
 Sumner, J. B. (1941). *Advances in Enzymology*, **1**, 163.
 Sumner, J. B. & Dounce, A. L. (1937). *J. biol. Chem.* **121**, 417.
 Sumner, J. B., Dounce, A. L. & Frampton, O. D. (1940). *J. biol. Chem.* **136**, 343.
 Theorell, H. & Agner, K. (1942). *Ark. Kemi Min. Geol.* **16 A**, no. 7.
 Weiss, J. (1937). *J. phys. Chem.* **41**, 1107.
 Zeile, K., Fawaz, G. & Ellis, V. (1940). *Hoppe-Seyl. Z.* **263**, 181.
 Zeile, K. & Hellström, H. (1930). *Hoppe-Seyl. Z.* **192**, 171.

The Chemical Conversion of Di-iodotyrosine into Thyroxine

By C. R. HARINGTON AND ROSALIND V. PITT RIVERS, *National Institute for Medical Research, Hampstead, N.W. 3*

(Received 3 February 1945)

The fact observed by Ludwig & von Mutzenbecher (1939), that iodination of certain proteins under controlled conditions gave rise to products from which thyroxine could be isolated, could be explained in one of two ways: either the proteins in question contained thyronine as a hitherto unrecognized constituent which was directly converted by iodination into thyroxine, or the tyrosine of the protein was iodinated to 3:5-di-iodotyrosine, part of which subsequently formed thyroxine. The latter mode of formation of thyroxine had indeed been suggested by one of us many years ago (Harington & Barger, 1927) and evidence, both chemical and biological, which had accumulated over a number of years (cf. Harington, 1944) favoured the supposition; its likelihood was further increased by the later observation of von Mutzenbecher (1939) that di-iodotyrosine on incubation in alkaline solution spontaneously gave rise to traces of thyroxine.

The over-all reaction representing the conversion of di-iodotyrosine into thyroxine can be crudely formulated as involving the loss, between two molecules of di-iodotyrosine, of one molecule of alanine; in such a formulation no oxidative process need necessarily be assumed. In actual fact, however, it is scarcely possible to conceive such an intermolecular dissociation, and it seems almost certain that the first stage of the reaction must be an oxidative process whereby the molecule of di-iodotyrosine is converted into one or more less stable and more reactive forms.

The supposition that oxidation must play a part is strengthened by a further consideration. By altering the conditions of iodination of a protein two entirely different results can be obtained. If the experiment is conducted as rapidly as possible in cold concentrated solution in presence of excess of ammonia the predominant and sometimes the

exclusive result is substitution of the tyrosine groups to form 3:5-di-iodotyrosine (cf. Neuberger, 1934; Harington & Neuberger, 1936). If, on the other hand, it is desired to form products containing thyroxine similar to those prepared by Ludwig & von Mutzenbecher the iodination must be conducted in warm dilute solution at a slightly alkaline reaction (pH about 8), and the thyroxine content of the products is increased by prolonged incubation of the reaction mixtures under aerobic conditions. In the present paper direct evidence will be given that oxidation is involved in the conversion of di-iodotyrosine into thyroxine, since this conversion fails when the incubation of di-iodotyrosine in alkaline solution is conducted in the complete absence of oxygen.

The suggestion that the formation of thyroxine during iodination of proteins under these conditions and during incubation of solutions of di-iodotyrosine was essentially an oxidative process was developed by Johnson & Tewkesbury (1942) who postulated hypiodite as the effective oxidizing agent in both cases; these authors indeed claimed increased yields of thyroxine as the result of addition of hypiodite to an alkaline solution of di-iodotyrosine, although they did not offer experimental evidence. Further reference to the question of the actual oxidizing agent involved will be made below.

Johnson & Tewkesbury proposed in outline a theory of the process involved in the formation of thyroxine from di-iodotyrosine and this was further developed by one of us (Harington, 1944) on the basis of a general consideration of the conditions of the reaction and of additional results obtained by ourselves. The present paper contains an account of these results and of further work which strengthens the basis upon which the theoretical treatment referred to was founded.

Formation of thyroxine by incubation of di-iodotyrosine

The theory of the oxidative conversion of di-iodotyrosine into thyroxine to which reference has been made was developed on the assumption that the process involved the amino-acid in the form of the phenoxide ion; such an assumption seemed justified by the fact that both under the conditions employed by von Mutzenbecher and under those obtaining *in vivo*, where the same transformation presumably occurs, the phenolic group of di-iodotyrosine will be largely ionized. (According to Winnek & Schmidt (1935) this group has pK 6.5.) The first matter to be considered therefore is the effect of pH on the formation of thyroxine under von Mutzenbecher's conditions, i.e. on incubation of di-iodotyrosine at 38° for 14 days. The results of our experiments on this point are summarized in Table 1. From this it appears that there is no

formation of thyroxine in acid solution, whilst on the alkaline side the formation attains a maximum at about pH 10 and falls off sharply at higher alkalinities. In considering these results it must be borne in mind that the pH values recorded are those at the beginning of the incubation; during the reaction acidic products are formed which cause a gradual fall of the pH.

Table 1. *Formation of thyroxine by incubation of di-iodotyrosine for 14 days at 38°*

Initial pH	Destruction of di-iodotyrosine (%)	Net yield of thyroxine (%)
1.0	22	0
7.8	36	0.74
8.8	22	1.53
10	22	2.82
13	17	0
14	13	0

Clearly the fact that formation of thyroxine becomes significant at pH 7.8, which results from addition of 1 molecular equivalent of sodium hydroxide to di-iodotyrosine, and reaches its maximum at pH 10, that is to say after 1.95 equivalents of alkali have been added, is in accordance with the idea that it is the phenoxide ion which is being oxidized. Discussion of the reason for the absence of thyroxine formation at higher alkalinities is best postponed until further experiments have been described.

Effect of added oxidizing agent on the formation of thyroxine

If the formation of thyroxine from di-iodotyrosine is in fact an oxidative process it should be possible to accelerate it and to obtain greater yields by addition of an appropriate oxidizing agent. It was pointed out by Johnson & Tewkesbury that the process seems to bear close resemblance to certain reactions observed by Pummerer, Melamed & Puttfarcken (1922) and Pummerer, Puttfarcken & Schopflocher (1925) in the course of their study of oxidation of phenols by alkaline ferricyanide; we therefore first investigated the action of potassium ferricyanide on di-iodotyrosine under the conditions employed by Pummerer *et al.* for the oxidation of simple phenols such as *p*-cresol; in no experiment of this type, however, were we successful in isolating thyroxine, or indeed any other pure oxidation product. Under the mildest conditions which we could devise, ferricyanide caused extensive disruption of the di-iodotyrosine with the production of intractable tars. A similar experience has recently been reported by Barkdoll & Ross (1944).

We therefore considered alternative oxidizing agents and selected hydrogen peroxide, largely

because the action of this so often simulates the over-all effects of biological oxidations.

In Table 2 is recorded a series of experiments which show that hydrogen peroxide is indeed effective in accelerating the formation of thyroxine from di-iodotyrosine at 38°, although the ultimate yields are somewhat lower. Comparison of Tables 1 and 2 shows that the figures for the destruction of di-iodotyrosine and the yield of thyroxine are of the same order after 15 hr. in presence of hydrogen peroxide as they are after 14 days of simple incubation; moreover, the effects of pH on thyroxine formation are similar in both cases, the maximum at pH 10 being clearly defined both with and without peroxide; an exception to this statement is provided by the low yield obtained at pH 8.8 in presence of peroxide; this may be related to intrinsic instability of thyroxine (see below).

Table 2. *Oxidation of di-iodotyrosine by hydrogen peroxide (0.5 atom. equiv. O) for 15 hr. at 38°*

Initial pH	Destruction of di-iodotyrosine (%)	Net yield of thyroxine (%)
1.0	18	Trace
7.8	26	0.75
8.8	21	0.36
10	17	1.42
13	15	0.24
14	13	0

Effects of elevation of temperature and of increasing amounts of hydrogen peroxide on yield of thyroxine

The favourable effect of hydrogen peroxide on the formation of thyroxine from di-iodotyrosine as indicated by the acceleration of the process led to further experiments under more drastic conditions. In Table 3A are shown the results of oxidations of di-iodotyrosine with hydrogen peroxide at varying pH values and at elevated temperature (90–100°); under these conditions the reaction was, as might be anticipated, still further accelerated, being complete within an hour; the destruction of di-iodotyrosine was much more extensive than at the lower temperature whilst the maximum yield of thyroxine was approximately the same. Moreover, the effect of pH on the reaction remained unaltered.

At this point it became clear to us that we were unlikely to make much further progress in increasing the yield of thyroxine without introducing a new condition, since it was apparent that such thyroxine as might be formed would itself be subject to the destructive effect of hydrogen peroxide which is so obvious in the case of di-iodotyrosine. We therefore decided to attempt to make use of the observation of Leland & Foster (1932) that from an alkaline solution containing both di-iodotyrosine and thyroxine the latter is preferentially extracted by

butanol. Accordingly we made further experiments in which the solution of di-iodotyrosine undergoing oxidation with hydrogen peroxide was shaken with butanol, the butanol layer being separated and renewed from time to time during the reaction. As is shown by the figures in Table 3B this method met with some success, the maximum yield of thyroxine obtained being higher than that recorded in any previous experiment.

Table 3. *Oxidation of di-iodotyrosine with excess hydrogen peroxide (5 atom. equiv. O) at 90–100°*

Initial pH	A. Without butanol		B. With butanol	
	Destruction of di-iodotyrosine (%)	Net yield of thyroxine (%)	Destruction of di-iodotyrosine (%)	Net yield of thyroxine (%)
1.0	34	Trace	—	—
7.8	100	Trace	41	0.23
8.8	90	0.13	—	—
10	46	1.63	26	3.19
13	20	0	18	2.13
14	—	—	16	0

In all the experiments recorded in Table 3 the amount of peroxide employed was greatly in excess of the theoretical, corresponding in fact to 5 atomic equivalents of oxygen. This particular proportion of peroxide had been adopted empirically as the result of a number of orientating experiments, and it became desirable to find whether the yield of thyroxine could be improved by using the butanol technique in conjunction with smaller or greater amounts of oxidizing agent at the optimum pH of 10. Table 4 shows that the yield of thyroxine is surprisingly little affected by increasing the amount of peroxide; this result is probably to be explained by formation of products in the course of reaction which catalyze the decomposition of hydrogen peroxide, so that successive additions of the latter become progressively less effective; such an effect is indeed visually obvious.

Table 4. *Effect of increasing amounts of hydrogen peroxide on formation of thyroxine from di-iodotyrosine at 90–100° at pH 10 (butanol technique)*

H ₂ O ₂ added (atom. equiv. of O)	Destruction of di-iodotyrosine (%)	Net yield of thyroxine (%)
1.7	27	2.8
5	30	3.2
15	39	3.4
25	42	4.0

Effect of pH

It is now necessary to give some further consideration to the effect of pH on the formation of thyroxine from di-iodotyrosine. It will have been noticed that this effect is the same under all the conditions which have been studied and that, in so far as it involves absence of formation of thyroxine in the acid range and maximum formation at the point where disso-

ciation of the acidic groups of di-iodotyrosine becomes complete, it is in accordance with expectation on theoretical grounds. The outstanding question is the reason for the failure of thyroxine formation at higher alkalinities.

The first point to be noted in this connexion is the peculiar behaviour of di-iodotyrosine itself, which is illustrated by the curve in Fig. 1.* This

Fig. 1. Effect of pH on destruction of di-iodotyrosine by heating in solution

curve shows the relative proportions of di-iodotyrosine which are destroyed (with the formation of amorphous products) by heating for 1 hr. at 100° in aqueous solution at different pH values; a range of instability of the amino-acid is revealed which we are unable to explain, although it is perhaps noteworthy that this instability is low at pH values where the ionizable groups of di-iodotyrosine are completely dissociated; whatever the explanation this property of di-iodotyrosine must evidently influence the interpretation of our results. It is clear that under the conditions of our experiments with hydrogen peroxide two processes at least must be going on, namely, the destruction of di-iodotyrosine due to its intrinsic instability and the specific oxidation to thyroxine, and that the final result observed in the form of the yield of thyroxine represents the summation of these two and probably other reactions. It will be noticed that in the region of high alkalinity, where formation of thyroxine fails completely, di-iodotyrosine is extremely stable; at pH 10, where oxidative formation of thyroxine is maximal, di-iodotyrosine is also stable, but, as already pointed out, acid production during the course of oxidation will cause a fall of pH during the reaction and the region of instability will be

* Since this was written a paper by Kraft & Denzel (*Hoppe-Seyl. Z.* 1942, 272, 147) has come to our notice where similar observations have been made in the alkaline range.

entered. It is also, perhaps, of significance that di-iodotyrosine is in its region of greatest instability at the reaction of the body tissues. It is unfortunate that no conclusive information can be obtained concerning the formation of thyroxine at reactions in the neighbourhood of the isoelectric point of di-iodotyrosine (pH 4.3: Winnik & Schmidt, 1935), but the insolubility of the amino-acid and the inapplicability of the butanol technique make experiments at this pH impracticable.

A further question which must be considered in relation to the pH effect is the nature of the oxidizing agent actually responsible for the conversion of di-iodotyrosine into thyroxine. As we have seen, the optimum reaction for this conversion is pH 10. When di-iodotyrosine is incubated at this reaction iodine is slowly liberated and can be detected in solution; it will naturally be converted in part into hypiodite. The action of hydrogen peroxide on di-iodotyrosine also brings about liberation of iodine, and the possibility exists that it is this effect rather than its direct oxidizing action on intact di-iodotyrosine which is responsible for the favourable effect of peroxide on formation of thyroxine.

Whilst there seems to be no certain means of determining whether hydrogen peroxide in our experiments is effecting the conversion of di-iodotyrosine into thyroxine by direct oxidation or by the liberation of iodine, which then brings about the reaction, it appears to us that there are two cogent reasons in favour of the latter supposition; these reasons are the significant degree to which the reaction proceeds in absence of peroxide, and the effect of pH. If peroxide were itself the effective oxidizing agent it would be natural to expect a greater increase than we have been able to observe in the yields of thyroxine obtainable in its presence, as compared with those resulting from simple incubation of di-iodotyrosine under aerobic conditions; moreover, it is not obvious why the effect of peroxide, if this were indeed to produce thyroxine directly, should fall off so sharply between pH 10 and 13.

Accepting the probability that in presence as in absence of peroxide the liberation of iodine is the decisive event, we may consider what this implies as to the nature of the actual oxidizing agent concerned. The two oxidants to be considered are iodine and hypiodite. It was stated by von Mutzenbecher (1939) (without experimental details) that the formation of thyroxine on incubating di-iodotyrosine in alkaline solution was unaffected by the presence of thiosulphate, although it was prevented by sulphite. If this statement were correct it would exclude both iodine and hypiodite as the effective agent since both are reduced by thiosulphate; in fact, however, we have ourselves found that the formation of thyroxine is almost completely inhi-

bited by the presence of thiosulphate, so that as far as this experiment is concerned both iodine and hypiodite remain possibilities. It has already been mentioned that Johnson & Tewkesbury (1942) claimed support for their theory, that hypiodite was the effective agent, by the increased yields of thyroxine obtainable after addition of hypiodite to the reaction mixture. Even if this claim had been substantiated, however, it would not constitute conclusive evidence; the equilibrium between iodine and hypiodite is governed by pH and thus, unless the pH were simultaneously raised, addition of 'hypiodite' would in fact mean the increase in concentration both of hypiodite itself and of free iodine. Assuming the correctness of the claim of Johnson & Tewkesbury therefore, their result could be explained equally well by the supposition that it was the increase in the concentration of free iodine which was effective. We actually describe below an experiment in which iodine is added to a solution of di-iodotyrosine at pH 10 with resulting formation of thyroxine in significant amount; this is, however, no more conclusive than the experiment of Johnson & Tewkesbury since we are merely approaching the same equilibrium mixture of iodine and hypiodite from the other side.

We ourselves indeed incline strongly to the view that it is iodine and not hypiodite which effects the oxidation of di-iodotyrosine to thyroxine. If hypiodite were responsible there is no apparent reason why the reaction should fail above pH 10, whilst if free iodine were needed such a failure would be expected; it is indeed just in the pH range where iodine is completely converted into hypiodite that failure of the oxidation occurs. The supposition that free iodine is the effective oxidizing agent also offers a reasonable conception of the events in the biological formation of thyroxine; there is now ample evidence that this formation occurs from di-iodotyrosine and the experiment of Keston (1944) makes it likely that iodine participates in the reaction. At the pH of the body fluids hypiodite can exist only in minute concentration; on the other hand, in presence of an enzymic oxidizing system capable of liberating iodine from iodide, all the conditions which we have supposed to be necessary for thyroxine formation would be operative; the free iodine produced would have the dual function of reacting with tyrosine to produce di-iodotyrosine and of oxidizing the latter to thyroxine. The conditions existing in the thyroid gland on this supposition would indeed be closely similar to those employed by Ludwig & von Mutzenbecher (1939) in their experiments on the iodination of proteins.

DISCUSSION

It appears to us that our results leave no doubt of the oxidative nature of the conversion of di-iodo-

tyrosine into thyroxine. Proof of this rests on the failure of the conversion in absence of oxygen and on the significant increase of yield and acceleration of the reaction which can be effected by the addition of an oxidizing agent.

The system as we have studied it is, however, greatly complicated by the peculiar intrinsic instability of di-iodotyrosine in the range of pH with which we are concerned, and by the fact that changes in pH not only affect the degree of ionization of di-iodotyrosine but may also influence the concentration of the effective oxidizing agent. It must further be reiterated that in speaking of pH we refer only to the initial value; a fall of 0.5–1 pH unit may occur during the oxidation. In these circumstances an adequate interpretation of the reaction cannot be given but, taking all the observed facts into account, we would offer the following tentative conclusions.

The fact that formation of thyroxine is not observed at pH values below that at which the phenolic group of di-iodotyrosine is significantly dissociated supports the hypothesis that the oxidation involves the phenoxide ion. If this assumption is accepted the failure of thyroxine formation at high alkalinity is to be explained (a) by increased stability of di-iodotyrosine and (b) by ineffectiveness or removal of the oxidizing agent. Reality is given to the possibility (b) by the supposition that the effective oxidizing agent is iodine, which is known to be liberated by simple aerobic oxidation of di-iodotyrosine and more rapidly by oxidation of the latter with hydrogen peroxide. The increasing yields of thyroxine over the range pH 7.8–10 would thus be explained by the favourable effect of increasing ionization of the acidic groups of di-iodotyrosine while free iodine is still present, whilst the failure of thyroxine formation at pH > 13 would be due to the absence of free iodine from the system.

Admittedly this interpretation leaves much to be desired in precision and completeness. Other unknown factors may come into play such as an intrinsic instability of thyroxine, similar to that observed for di-iodotyrosine but operative over a different range of pH. The interpretation given is, however, in accordance with the facts as we have observed them, and it is perhaps worthy of consideration in its biological implications.

EXPERIMENTAL

Formation of thyroxine by incubation of di-iodotyrosine at 38°

The following experiment is typical of the series, the results of which are shown in Table 1. The example chosen for detailed description is that which imitates as closely as possible the conditions described by von Mutzenbecher (1939).

l(+)-Di-iodotyrosine dihydrate (11.75 g.) was dissolved in 0.1N-NaOH (25 ml.); the solution was adjusted to pH 8.7 by addition of N-NaOH (3 ml.) and incubated at 38° for 14 days. At the end of this time 12.5N-NaOH (20 ml.) was added and the mixture was twice extracted with its own volume of *n*-butanol; the combined extracts were washed once with an equal volume of 2N-NaOH and evaporated under diminished pressure. The residue was taken up in water (50–60 ml.) and the solution was acidified at the boiling point.

The precipitate was collected from the cooled solution and dried in a desiccator (152 mg.); it was then extracted with ether and the ether-insoluble portion (78 mg.) was dissolved in the minimum of boiling 0.1N-K₂CO₃ (about 5 ml.); the filtered solution was kept overnight in the ice-chest. The white and crystalline potassium salt which had separated was converted into free crystalline thyroxine by solution in aqueous-ethanolic alkali and acidification at the boiling point with acetic acid. Yield, 32 mg.

The original aqueous solution was freed from butanol by evaporation under diminished pressure, and strongly acidified at the boiling point; a tarry precipitate was removed by rapid filtration of the hot solution and the pH was adjusted to about 5. The di-iodotyrosine which separated on cooling, together with a second crop obtained by concentrating the mother liquor, amounted to 9.22 g. or 79.5% of the starting material.

This means that the net yield of thyroxine, calculated on the di-iodotyrosine destroyed, was 1.5%; the gross yield was 0.33%. von Mutzenbecher records a gross yield of 0.24%; he does not state the amount of di-iodotyrosine recovered unchanged so that his net yield cannot be calculated. Block (1940) and Barkdoll & Ross (1944), working under conditions similar to ours, obtained gross yields of 0.12 and 0.2–0.35% respectively.

Effect of thiosulphate. An experiment on the lines just described but starting at the more favourable initial pH of 10 (cf. Table 1) was carried out in presence of sodium thiosulphate equivalent in amount to the iodine content of the di-iodotyrosine used. The period of incubation was reduced in this case to 9 days and a control experiment was run simultaneously for the same period under the ordinary conditions; the amount of di-iodotyrosine used in each experiment was 9.4 g. of the dihydrate. In the control the thyroxine isolated amounted to 46 mg. and in the thiosulphate experiment to 1 mg.; allowing for the recovery of unchanged di-iodotyrosine these figures correspond to net yields of 1.6 and 0.04% respectively; it is apparent therefore that the presence of thiosulphate almost entirely suppresses the formation of thyroxine.

Effect of exclusion of oxygen. In the preceding experiment a third similar quantity of di-iodotyrosine was incubated for the same length of time under completely anaerobic conditions (McIntosh-Fildes jar). No thyroxine could be isolated and 92% of the di-iodotyrosine was recovered unchanged.

Experiments with added oxidizing agents

(1) *Potassium ferricyanide.* Oxidation of di-iodotyrosine in alkaline solution with potassium

ferricyanide under the conditions employed by Pummerer and his co-workers (1922, 1925) for the oxidation of *p*-cresol led only to the formation of intractable tarry products of acidic nature. An attempt was made to apply the butanol technique subsequently successfully used with peroxide (see below).

Di-iodotyrosine dihydrate (11.75 g.) was dissolved in NaOH (47 ml. of 1.015N; 95% neutralization) and 3 ml. water and the solution was chilled in ice. *n*-Butanol (50 ml.) was added and the mixture, continuously shaken in an ice-bath, was treated with a solution of potassium ferricyanide (8.22 g.) in water (25 ml.) added over 30 min. The mixture was shaken at room temperature for an hour and the layers were separated; the aqueous solution was re-adjusted to pH about 8.5 (20 ml. of N-NaOH required); more butanol (100 ml.) was added and the oxidation continued at 0° with a further 8.22 g. of ferricyanide in 25 ml. water, the reaction being kept alkaline by periodic addition of NaOH (total 30 ml. N solution). The butanol extracts were washed with N-NaOH and evaporated. The residue was taken up in water and precipitated at the boiling point with HCl. The material thus obtained, after drying, was almost completely soluble in ether; no thyroxine could be isolated from the small ether-insoluble residue, nor was any di-iodotyrosine recoverable from the original aqueous solution.

(2) *Hydrogen peroxide at 37°.* An experiment typical of those recorded in Table 2 was as follows. Di-iodotyrosine dihydrate (18.8 g.) was dissolved in N-NaOH (76 ml.; 95% neutralization) to give a solution having pH about 10. Hydrogen peroxide (0.5 atomic equiv. O) was added and the mixture was incubated at 38°. After 9 hr. very little peroxide was detectable. After about 16 hr. the solution was worked up as described under the simple incubation experiments. The thyroxine isolated amounted to 36 mg. and the di-iodotyrosine recovered to 3.52 g.; the net yield of thyroxine was thus 1.5% and the gross yield 0.23%.

(3) *Hydrogen peroxide in excess at higher temperature* (Table 3A). Di-iodotyrosine dihydrate (9.4 g.) was dissolved in N-NaOH (38 ml.) and the solution was heated on the boiling water-bath. Hydrogen peroxide (20 vol. solution; 5 atomic equiv. O in all) was added in fifteen portions over about 1½ hr.; the reaction was maintained alkaline to phenolphthalein throughout. The solution, which had become very dark, was worked up as usual. Of the di-iodotyrosine 54% was recovered; the thyroxine isolated was 58 mg. (1.6% net yield; 0.6% gross).

(4) *Hydrogen peroxide in excess at high temperature with butanol* (Table 3B). The most successful experiment by this technique, the rationale of which has been discussed in the introductory portion of the paper, was the following.

Di-iodotyrosine dihydrate (23.5 g.) was dissolved in N-NaOH (95 ml.); 100 ml. of *n*-butanol, previously saturated with water, were added and the mixture was heated on the boiling water-bath. Additions of 10 ml. portions of 20 vol. hydrogen peroxide were then made, with continued heating and vigorous shaking, at intervals, such that peroxide did not accumulate in the solution. After each five additions

of peroxide the butanol layer was separated and replaced with fresh solvent, the amounts being increased in approximate parallelism with the increase in volume of the aqueous layer; the pH of the latter was readjusted to approximately 10 at frequent intervals during the experiment by addition of the requisite amount of N -NaOH. In all, 75 additions of hydrogen peroxide were made, i.e. approximately 25 atomic equivalents of oxygen/mol. of di-iodotyrosine; the entire procedure occupied about 5 hr. In the early stages the butanol extracts were fairly highly coloured, but the last few were almost colourless.

The combined butanol extracts were concentrated under diminished pressure to 1500 ml. and extracted once with 1000 ml. of N -NaOH; the latter extract was washed with 500 ml. of butanol and this washing, together with the main butanol solution, was evaporated to dryness. The residue was taken up in water (400 ml.) and after removal of traces of butanol by a brief vacuum distillation the solution was brought to the boil and acidified (about pH 4) with dilute H_2SO_4 . Next day the precipitate was collected and dried in a vacuum desiccator. The dried material (1.55 g.) was ground up with dry ether and collected, this process being repeated three times; the last ethereal extract was almost colourless and the insoluble residue amounted to 0.87 g. This was dissolved in the minimum amount (64 ml.) of boiling $0.1N$ - K_2CO_3 ; after 22 hr. at 0° the precipitate of potassium salt was collected and converted into thyroxine in the usual manner by solution in alkaline aqueous ethanol followed by precipitation at the boiling point with acetic acid; the pure crystalline thyroxine thus obtained amounted to 278 mg. Further crops of thyroxine, amounting in all to 45 mg., were isolated by collecting all the aqueous acid-insoluble material from the mother liquors of the potassium salt and of the thyroxine crystallization and purifying it by repeating the extraction from alkaline solution into butanol and separation as potassium salt. The total yield of thyroxine was thus 323 mg.

The original aqueous solution was combined with the alkaline washing of the butanol extracts and concentrated somewhat under diminished pressure; it was then brought to the boil, acidified strongly with concentrated HCl and filtered hot from a tarry precipitate; the filtrate was adjusted to pH 4 by addition of $10N$ -NaOH and the di-iodotyrosine which separated on cooling was collected (10.66 g.). Two further crops of di-iodotyrosine were obtained by concentration of the mother liquor, the total amount recovered being 13.71 g.

The di-iodotyrosine which had been oxidized was therefore 9.79 g. and the net yield of thyroxine was 4%.

The thyroxine which was isolated in this and other experiments crystallized in the characteristic sheaves of bent needles and had m.p. 240° (decomp.); it gave a positive ninhydrin reaction and a clear red colour after heating with nitrite and HCl in ethanol followed by treatment with excess of ammonia. The product obtained in the experiment just described was completely analyzed with the following results. Found: C, 23.5; H, 1.7; N, 1.8; I, 65.3%. (Calc. for $C_{15}H_{11}O_4N_4I_2$: C, 23.2; H, 1.4; N, 1.8; I, 65.4%.) It also showed activity in

the tadpole metamorphosis test equal to that of authentic thyroxine.*

(5) *Iodine*. Di-iodotyrosine dihydrate (9.4 g.) was dissolved in N -NaOH (38 ml.); iodine (2.65 ml. of 3.78*N* solution; 0.5 atomic equiv. O) was added, followed by *n*-butanol (60 ml.). The whole was shaken for 3 hr. at 37° after which no free iodine could be detected; the aqueous layer was then made N with respect to NaOH, the butanol was separated and, together with a second similar extract, worked up as usual; 15 mg. of thyroxine were obtained, whilst 72% of the di-iodotyrosine was recovered from the aqueous solution. The net yield of thyroxine was therefore 0.4%.

Optical activity of l(+)-di-iodotyrosine and l(-)-thyroxine

The di-iodotyrosine which was used in the experiments described in this paper was prepared from optically pure *l*-(-)-tyrosine having $[\alpha]_D^{25} = -12.4^\circ$ in 5% solution in N -HCl; the iodination was carried out either by the method of Harington (1928) or that of Block & Powell (1943) and in all cases the product had $[\alpha]_D^{25} = +1.5^\circ$ in N -HCl, the concentration being about 5%. This rotation is considerably lower than that previously recorded (cf. Abderhalden & Guggenheim, 1908; Harington, 1928), but our material was undoubtedly optically pure, since on catalytic deiodination it gave a 94% yield of tyrosine having $[\alpha]_D^{25} = -12.2^\circ$ in N -HCl. We feel therefore that $+1.5^\circ$ must be regarded as the correct figure for the rotation of *l*(+)-di-iodotyrosine under the prescribed conditions, although we can offer no explanation of the higher figures previously recorded.

There is no reason to suppose that any racemization is involved in the oxidative conversion of di-iodotyrosine into thyroxine, so that the rotation of the thyroxine isolated in our present experiments should be that of the optically-pure natural substance. We have found for our preparation $[\alpha]_D^{25} = -4.3^\circ$ ($c = 3.03$ in a solution made up of 24 g. 0.5*N*-sodium hydroxide and 56 g. of ethanol); this compares with $[\alpha]_D = -4.4^\circ$ observed by Foster, Palmer & Leland (1936) under the same conditions for a sample of thyroxine isolated from thyroid by acid hydrolysis. The figure now obtained is larger than that observed by Harington (1928) for *l*(-)-thyroxine prepared by resolution of *dl*-3:5-di-iodo-tyronine and subsequent iodination; it must be concluded that this resolution was not quite complete. It may be noted that the rotation of thyroxine, like that of other amino-acids, is much influenced by the nature of the solvent; thus our present preparation, dissolved at a concentration of 3.28% in a mixture of 1 vol. N -NaOH and 2 vol. ethanol, had $[\alpha]_D^{21} = -5.04^\circ$.

* We are indebted to Dr R. Deanesly for carrying out this test.

Intrinsic instability of di-iodotyrosine (see Fig. 1)

Quantities of 470 mg. (1 mmol.) of di-iodotyrosine dihydrate were dissolved rapidly by boiling in 25 ml. water containing the requisite amount of HCl or NaOH to give the desired pH. The solutions were then heated for 1 hr. on a boiling water-bath, cooled, treated with 12.5N-NaOH to make a final concentration of N and extracted once with 25 ml. of butanol. The aqueous layer was separated quantitatively and submitted to a brief vacuum distillation to remove butanol; it was then transferred to a boiling tube, being made up to 30 ml. with washings, and acidified when hot with 3 ml. glacial acetic acid.

At the end of the period of heating a great difference in the behaviour of di-iodotyrosine at the various pH values was already apparent; the solutions in the pH range 6–9 became dark brown, and at the stage of acidification threw down a dark-coloured amorphous precipitate which was removed by filtration before crystallization of residual di-iodotyrosine set in. In all cases crystallization was completed by keeping in the ice-chest overnight; the residues of unchanged di-iodotyrosine were then collected, dried *in vacuo* over P_2O_5 and weighed.

SUMMARY

1. The work of von Mutzenbecher (1939) on the formation of thyroxine by incubation of di-iodo-

tyrosine in alkaline solution is confirmed. It is shown that this conversion is prevented by exclusion of atmospheric oxygen or by addition of thiosulphate to the solution; the reaction is also profoundly influenced by the pH during incubation.

2. The effect of added oxidizing agents on the conversion of di-iodotyrosine to thyroxine has been studied; with the aid of hydrogen peroxide increased yields of thyroxine can be obtained in a much shorter time, the effect of pH being similar to that observed on simple aerobic incubation. The conversion can also be accelerated by addition of iodine. Reasons are given for supposing that in all cases observed the actual agent effective in bringing about the oxidation of di-iodotyrosine to thyroxine is free iodine, and attention is called to the biological implications of this hypothesis.

3. The thyroxine obtained by oxidation of *l*(+)-di-iodotyrosine, itself derived from *l*(-)-tyrosine, is shown to have an optical rotation identical with that of the natural product; this provides direct chemical proof of the configurative relationship of natural tyrosine and natural thyroxine.

4. Di-iodotyrosine in aqueous solution has been found to exhibit a remarkable degree of instability towards heat over the pH range 4–9; at higher acidity or alkalinity the amino-acid is comparatively stable.

REFERENCES

- Abderhalden, E. & Guggenheim, M. (1908). *Ber. deutsch. chem. Ges.* **41**, 1238.
 Barkdoll, A. E. & Ross, W. F. (1944). *J. Amer. chem. Soc.* **66**, 898.
 Block, P., Jr. (1940). *J. biol. Chem.* **135**, 51.
 Block, P., Jr. & Powell, G. (1943). *J. Amer. chem. Soc.* **65**, 1430.
 Foster, G. L., Palmer, W. W. & Leland, J. P. (1936). *J. biol. Chem.* **115**, 467.
 Harrington, C. R. (1928). *Biochem. J.* **22**, 1429.
 Harrington, C. R. (1944). *J. chem. Soc.* p. 193.
 Harrington, C. R. & Barger, G. (1927). *Biochem. J.* **21**, 169.
 Harrington, C. R. & Neuberger, A. (1936). *Biochem. J.* **30**, 809.
 Johnson, T. B. & Tewkesbury, L. B. (1942). *Proc. nat. Acad. Sci., Wash.*, **28**, 73.
 Keston, A. S. (1944). *J. biol. Chem.* **153**, 335.
 Leland, J. P. & Foster, G. L. (1932). *J. biol. Chem.* **95**, 165.
 Ludwig, W. & von Mutzenbecher, P. (1939). *Hoppe-Seyl. Z.* **258**, 195.
 von Mutzenbecher, P. (1939). *Hoppe-Seyl. Z.* **261**, 253.
 Neuberger, A. (1934). *Biochem. J.* **28**, 1982.
 Pummerer, R., Melamed, D. & Puttfarcken, H. (1922). *Ber. deutsch. chem. Ges.* **55**, 3116.
 Pummerer, R., Puttfarcken, H. & Schopflocher, P. (1925). *Ber. deutsch. chem. Ges.* **58**, 1808.
 Winnek, P. S. & Schmidt, C. L. A. (1935). *J. gen. Physiol.* **18**, 892.

(An Addendum has been added on p. 212.)

Note on the Nutritive Value of the Nitrogenous Substances contained in Dried Yeast (*Torulopsis lipofera*)

BY H. CHICK AND E. B. SLACK, *Division of Nutrition, Lister Institute, Roebuck House, Old Chesterton, Cambridge*

(Received 6 February 1945)

The yeast was a dried specimen of *Torulopsis lipofera*, a species having a high fat content, provided by Dr A. Kleinzeller (1944). Its composition was: moisture less than 1, fat 18.6, nitrogen 5.85, 'crude' protein (N × 6) 35.1%. Analyses by Stutzer's

method showed that about 10% of the nitrogen was non-protein in character.

The following experiments were designed to show the nutritive value of the nitrogenous substances, protein and non-protein, contained in the yeast, as

washing with water. The washed solution was dried over anhydrous sodium sulphate and reabsorbed on calcium carbonate. The completed chromatogram exhibited three definite bands, the uppermost of which was faintly yellow (violaxanthin). The middle orange-yellow band, which contained the main portion of the colouring matter, was eluted, and the eluate crystallized first from methanol and subsequently from carbon disulphide, the pigment being obtained as orange-red crystals, m.p. 194°; absorption maxima in carbon disulphide 511, 478, 446 m μ , in chloroform 487, 456, 427 m μ . The solution in concentrated acetic acid developed a yellow colour and in dichloroacetic acid the pigment dissolved to a greenish solution. This pigment is therefore lutein. The third pigmented zone at the base of the column was spectroscopically identical with the pigment contained in the bottom zone of the original chromatogram, to which it was therefore added.

BOTTOM ZONE. The eluate from this, which contained very little colouring matter, was reabsorbed on calcium carbonate with the result that a separation from a further quantity of lutein was effected, but the quantity of pigment obtained was not sufficient for the isolation of any colouring matter; absorption maxima in carbon disulphide 516, 483, 451 m μ . It was probably zeaxanthin.

Extract A. The light-petroleum extract was washed free from methanol, dried over anhydrous sodium sulphate and adsorbed on aluminium oxide (Brockmann), the chromatogram being developed with light petroleum (b.p. 40–60°). The deep orange-red zone at the top of the column was eluted with benzene-ethanol and the solution concentrated to small bulk. After standing in the ice-chest for 24 hr. the precipitated pigment was repeatedly extracted with boiling ethanol to remove colourless impurities. Recrystallization of the residue from benzene gave reddish metallic-glancing prisms, m.p. 184°; absorption maxima in CS₂, 520, 484, 452 m μ and in petroleum ether, 485, 452, 424 m μ . This pigment is therefore β -carotene.

Ether extract. The deep red solution, obtained by extraction with diethyl ether of the methanolic mother liquors remaining after precipitation of the pigments by water, was washed free of alkali, dried over anhydrous sodium sulphate and the solvent removed *in vacuo*. The residual dark red oil product was dissolved in benzene and adsorbed on aluminium oxide, the chromatogram being developed with a 1:1 mixture of benzene and chloroform and finally with pure chloroform. The chromatogram contained two orange-red bands on top and a reddish diffuse band underneath. This latter band had no distinct absorption maxima and was not further examined. The top band, on elution, gave a dark red waxy solid; it showed absorption maxima in CS₂ at 497, 466, 437 m μ , in close agreement with the spectroscopic data for the unidentified lipochrome pigment isolated from *Viola tricolor* by Kuhn & Winterstein (1931).

The middle band of the above chromatogram had absorption maxima in CS₂ at 502, 470, 440 m μ . The pigment from this was dissolved in carbon disulphide and reabsorbed on calcium carbonate. It gave two yellow bands very near the top of the column, the upper of which had absorption maxima in CS₂ at 497, 466, 437 m μ (identical with those of the waxy material above). The second and main band had absorption maxima at 501, 469, 440 m μ . The eluate from this gave the characteristic violaxanthin reaction with 25 % hydrochloric acid.

SUMMARY

1. *Iris pseudacorus* has been examined and found to contain β -carotene and the xanthophylls violaxanthin and lutein.

2. A third xanthophyll, present in minute quantities, was spectroscopically identical with zeaxanthin.

3. In addition an uncrystallizable waxy carotenoid was present, similar in properties to that isolated by Kuhn & Winterstein (1931) from *Viola tricolor*.

REFERENCES

- Kuhn, R. & Winterstein, A. (1931). *Ber. dtsh. chem. Ges.* **64**, 326.
 Kuhn, R., Winterstein, A. & Lederer, E. (1931). *Hoppe-Seyl. Z.* **197**, 143.
 Molisch, H. (1896). *Ber. dtsh. bot. Ges.* **14**, 16, 27.
 O'Connor, W. F. & Drumm, P. J. (1941). *Nature, Lond.*, **147**, 58.
 Robinson, G. M. & Robinson, R. (1934). *Biochem. J.* **28**, 1712.
 van Wisselingh, C. (1917). *Flora batava*, **177**, 371.

Addendum to paper by C. R. Harington & Rosalind V. Pitt Rivers (see p. 164)

The following information about the hydrates of *l*(+)-di-iodotyrosine is worth recording. Di-iodotyrosine crystallizes from water, either by separation from the hot solution or by neutralization of a solution in aqueous acid as the dihydrate which forms glistening plates, which retain their water of crystallization on air-drying. (Found: I, 53.9%; calc. for C₉H₉O₃NI₂·2H₂O: I, 54.1%.)

On drying the above dihydrate over P₂O₅ *in vacuo* the water of crystallization is lost. (Found: I, 58.0%; calc. for C₉H₉O₃NI₂: I, 58.6%.)

Separation of di-iodotyrosine by evaporation of an ammoniacal solution at a bath temperature $\pm$ 40° yields long hair-like needles of the monohydrate. (Found: I, 56.1%; calc. for C₉H₉O₃NI₂·H₂O: I, 56.3%.)

When di-iodotyrosine separates slowly from 60 % acetic acid it forms large prisms, containing one molecule of acetic acid, which are stable to desiccation *in vacuo* over P₂O₅. (Found: I, 51.5%; calc. for C₉H₉O₃NI₂·C₂H₄O₂: I, 51.5%.) When the separation is rapid the form obtained is the monohydrate described above.