In the particular field of separation of the products of partial hydrolysis of proteins, the use of ionophoresis several times successively at different pH values on the same fractions promises to effect separation into a number of classes; and though it is improbable that its use alone will serve for the isolation of any single substance, its use as part of the separation will probably prove essential.

With the apparatus we have constructed, not more than one or two millimoles electrolyte can be used in the sample to be analyzed. These quantities are too small for convenience for many purposes, but could be increased by increasing the area of the apparatus. The thickness of the jelly cannot profitably be increased very far. The use of thinner glass and water cooling of the lid would increase the permissible thickness by a factor of about three. Thus, there seems no reason why several grams of materials could not be treated in an apparatus of laboratory size.

Further work would be required to make this method fully quantitative. So far we have obtained recoveries ranging from 80 to 100 %. It is also inherently probable that by running the bands sufficiently far apart, mutual contamination can be reduced to any desired degree. In the experiments described above, the method of test employed would have revealed a contamination of more than 1 or 2 %, but we believe that the actual figure was much lower than this.

SUMMARY

1. A method for the separation of substances of different mobility by ionophoresis in a slab of silica jelly has been developed.

2. A theory of the movement of charged substances in jelly containing a swamping concentration of electrolyte is presented, and a chart showing the relationship of net mobility, ion mobility and dissociation constant is given.

3. A wool hydrolysate has been separated into basic, neutral and acidic fractions at neutral pH.

4. The following pairs of substances have been separated by taking advantage of their different net mobilities at the optimum pH for separation. Glycine and serine; alanine and methionine; lysine and histidine; glycine and glycyglycine; glutamic, aspartic, and cysteic acids.

5. Glutamic and aspartic acids have been separated at neutrality by taking advantage of their differing ion mobilities.

6. Colour reactions, applicable to paper prints have been developed which reveal the presence and position of bands of arginine, histidine; methionine and serine.

Thanks are due to the Director and Council of the Wool Industries Research Association and to the International Wool Secretariat for permission to publish this work. We also thank Mr R. C. Palmer for discussion and criticism and Mr V. Newhouse for assistance with the diagrams.

REFERENCES

- Consden, R., Gordon, A. H. & Martin, A. J. P. (1944). *Biochem. J.* **38**, 224.
 Coolidge, T. B. (1939). *J. biol. Chem.* **127**, 551.
 Gawrilow, N. I., Paradashvili, A. I., Balabouha-Popzova, W. S. & Ljapounzowa, S. W. (1938). *Bull. soc. chim. Pays-Bas* (5), **5**, 973.
 Gordon, A. H., Martin, A. J. P. & Synge, R. L. M. (1943). *Biochem. J.* **37**, 79.
 Ikeda, K. & Suzuki, S. (1909). B.P. 9440/09.
 Kendall, J. (1942). *Nature, Lond.*, **150**, 136.
 MacInnes, D. A. (1939). *Principles of Electrochemistry*, p. 60. New York: Rheinhold Publishing Corporation.
 Martin, A. J. P. & Synge, R. L. M. (1941). *Biochem. J.* **35**, 1358.
 Martin, A. J. P. & Synge, R. L. M. (1945). *Advances in Protein Chemistry*, **2**. New York: Academic Press, Inc.
 Pauly, H. (1904). *Hoppe-Seyl. Z.* **42**, 508.
 Sakaguchi, S. (1925). *J. Biochem., Tokyo*, **5**, 13.
 Synge, R. L. M. (1945a). *Biochem. J.* **39**, 355.
 Synge, R. L. M. (1945b). *Biochem. J.* **39**, 363.
 Williams, R. J. (1935). *J. biol. Chem.* **110**, 589.

The Determination of Nitrogen and of Protein in Pooled Samples of Human Plasma

By R. P. COOK, *Department of Physiology, University College (University of St Andrews), Dundee*

(Received 26 October 1945)

The determination of plasma proteins has assumed importance in recent years on account of the value of the result in dietary surveys and in clinical medicine. The main methods available are: (i) the determination of protein-N (total N minus non-

protein-N) and subsequent conversion of N into protein by use of the factor 6.25 assuming that the total proteins contain 16 % N; (ii) gravimetric estimation. Here the acetone-treated, heat-coagulated proteins are washed, dried and weighed. A good

account of the method is given by Robinson & Hogden (1941); (iii) densitometric methods. These are necessarily indirect but are very convenient for a quick and approximate method of determination (Phillips, Van Slyke, Dole, Emerson, Hamilton & Archibald, 1945; Lloyd, Cheek, Sinclair & Webster, 1945). A comparison of other methods is described by Plimmer & Lowy (1945). Whilst the gravimetric method gives a direct assay, indirect methods dependent on N content are very convenient and should be accurate. During investigations on the determination of plasma proteins in a group of 45 boys (Cook, Keay & McIntosh, 1944, 1945) it was found that individual discrepancies existed between the gravimetric and micro-Kjeldahl methods of estimation, and it was felt that the protein factor used for the conversion of N was possibly at fault. To investigate the subject further it was necessary to use pooled samples of human plasma so that more extensive determinations could be made.

METHODS

Collection of blood and preparation of plasma. This subject is dealt with in detail by Cook *et al.* (1945), but the following points may be emphasized. At least 20 ml. of whole blood were collected from each of the donors who were all apparently healthy; potassium oxalate (90 mg./20 ml. blood) was used as anticoagulant. After the plasma was separated from the individual samples the material was pooled.

Description of pools. Pool A1 was made up by withdrawing 1 ml. from each sample of plasma derived from 45 boys. Pool A2 was prepared by mixing together all the remainders of plasma, 120 ml. being obtained. These two pools were prepared after the individual plasma samples had been stored in some cases for 6 months at -4° . Apart from a slight precipitation of fibrin in some samples there was no obvious change, a semi-sterile technique having been adopted throughout in the earlier sampling. Pool B was prepared from the combined plasmas of 12 adult males, pool C from 6 adult males and pool D from 4 adult females. In the cases of pools B, C and D the period elapsing between collection and analysis was at most 4 hr.

Pool A1 was used for the determination of the distribution of the protein fractions and for a comparison of the methods. Pool A2 was used solely to prepare total plasma proteins. The other pools were used for the protein distribution, comparison of methods and preparation of total plasma proteins. Pools B, C and D were also used to prepare a 'globulin' fraction, and pools C and D an 'albumin' fraction.

Preparation of total coagulated plasma proteins. To 1 vol. of the pooled plasma 10 vol. of acetone were added with constant mixing. The mixture was allowed to stand at room temperature for 12 hr. with occasional mixing and was then centrifuged. The supernatant fluid was poured off and a further 10 vol. acetone added, the precipitate of protein being well broken up. After standing for $\frac{1}{2}$ hr. the material was centrifuged. The washing with acetone was repeated. The precipitated proteins were then treated with 10 vol. 0.6% (w/v) NaCl, when only a small amount of material remained undissolved. The dissolved and suspended material was placed in a beaker and the pH adjusted to 5.0

with *N*-acetic acid. The beaker and contents were then placed in a boiling water-bath and kept constantly mixed. Flocculation readily occurred. The heating was continued for 1 hr., the mixture centrifuged and, after decanting the first supernatant fluid, a washing with 5 vol. of hot distilled water was carried out in the centrifuge cups. After this washing the protein coagulum was suspended in 5 vol. distilled water in a beaker which was placed in a boiling water-bath for 15 min., being kept well mixed during this process. The material was then poured through a sintered glass filter (Jena no. 17 G4) and washed with hot distilled water until the filtrate no longer gave a reaction for chloride. The coagulum was then washed with portions of ethanol equivalent in all to 5 vol. and with a similar amount of anhydrous ethyl ether. The fluffy, white to greyish material was dried in a shallow dish at 100° in an air-oven when it shrank to a horny yellowish mass. The yields from the four pools as g./100 ml. plasma were 6.9, 7.0, 6.6 and 6.8.

Preparation of 'globulin'. This fraction must be regarded only as a clinical entity, as in addition to the separate globulin components (see Discussion) it contains fibrinogen. To 1 vol. of plasma were added 30 vol. of 22% (w/v) Na_2SO_4 solution, and after standing at 37° for 4 hr. the mixture was filtered at 37° through a sintered filter as above. Filtration was rapid. The precipitate was washed with the Na_2SO_4 solution. The 'globulin' cake was then suspended in 0.6% saline and subjected to the same treatment as described above for the total proteins. The washed and dried 'globulin' finally obtained was whiter and less horny than the total coagulable protein. Yields as g./100 ml. plasma were: 2.9, 2.7 and 2.3.

Preparation of 'albumin'. From pools C and D an 'albumin' fraction was prepared thus. The filtrate from the globulin fraction was saturated at room temperature with solid ammonium sulphate. The precipitate so obtained was then filtered off, resuspended in 0.6% saline and treated as for the total coagulable proteins. Yields as g./100 ml. were 3.6 and 4.6. The isolated materials gave negative tests for the presence of ammonium sulphate.

Analytical methods

Before analysis all proteins were finely ground in an agate mortar. Moisture was determined by drying in thin layers in weighing bottles *in vacuo* over P_2O_5 at 100° . The tops of the weighing bottles were replaced immediately on removal from the drying oven. Some samples were dried in an air-oven at 105° . No difference in result was obtained. The values reported by Drs Weiler and Strauss (micro-quantities) were obtained by 'prolonged drying at 100° in high vacuo'. Ash was determined by igniting in an electric furnace at 1000° for 15 min. The micro-Kjeldahl method was used for determining the partition of N in the plasma and the N content of the isolated proteins, these being treated thus. The material, oven dried at 100° , was weighed direct into the digestion flask and digested for 16 hr. using the digestion mixture described by Chibnall, Rees & Williams (1943). It was then made up to volume, and the distillation carried out on triplicate samples, a deviation of $\pm 1\%$ in the titration being the only permissible variation. Repetition was rarely necessary. For pool A2 proteins the Parnas-Wagner type of distillation apparatus was used, but with the proteins from the other pools the more satisfactory Markham (1942) apparatus was employed. The NH_3 was trapped in boric acid and determined by titration

with 0.01 N-HCl. Frequent control experiments testing the apparatus and standard solutions for recoveries of NH_3 from a standard solution of ammonium sulphate were carried out. One sample (total proteins from pool A2) was kindly analyzed by Mr M. W. Rees and was used as an additional check.

The micro-Dumas method was used by Drs Weiler and Strauss for the values shown in Table 1.

Fractionation of plasma-N. This followed the methods given by Peters & Van Slyke (1932), N in the fractions being determined by the micro-Kjeldahl method, the preliminary digestion being continued for 16 hr. after clearing.

Gravimetric method. This was carried out by method A of Robinson & Hogden (1941) on 3 ml. portions of plasma. One important difference in technique was that the heating to bring about coagulation was continued for only 15 min. instead of 1 hr. The washings from the estimations of pools B, C and D were collected and the N content and biuret reaction determined. The coagulated proteins were dried *in vacuo* at 100° over P_2O_5 to constant weight.

RESULTS

Table 1 gives the percentage N (micro-Kjeldahl and micro-Dumas methods) after correction for moisture and ash content in the total coagulable proteins derived from four pools and in the 'globulin' derived from three pools. The following points are emphasized. The dried material constituting the proteins is a difficult and unsatisfactory substance to analyze, there being evidence of a sampling error for the individual samples. As would be expected for a complex mixture there are variations in the analyses of the different pools, the N content of the total plasma proteins ranging from 14.7 to 15.9%, with a mean value of 15.2%. From the three samples of 'globulin' a mean value of 14.3% may be taken. Values obtained for two samples of 'albumin' (corrected for moisture and ash) were 15.2 and 15.6%.

Table 1. *Analyses of total coagulable plasma proteins and 'globulin' fraction from the various pools*

(The individual N values on the pools by both micro-Dumas and micro-Kjeldahl methods are given after correction for moisture and ash content.)

	g./100 g. of protein				
	Pool A2	Pool B	Pool C	Pool D	Mean of pools
I. Total coagulable plasma proteins					
Moisture	3.8	0.8	4.1	2.6	—
Ash	1.1	0.3	1.2	1.6	—
N (corrected):					
Micro-Dumas	15.25, 15.91	15.03	15.23, 15.24	13.29, 15.70, 15.96	15.20
Micro-Kjeldahl	15.40, 15.41, 15.54	15.37	14.97	14.71	15.23
II. 'Globulin'					
Moisture	—	1.6	2.5	7.5	—
Ash	—	3.4	2.9	6.7	—
N (corrected):					
Micro-Dumas	—	14.04	13.99, 14.33	14.63	14.19
Micro-Kjeldahl	—	14.54	14.40	13.45, 15.13	14.44

Table 2. *The distribution of plasma-N (nos. 1-5), gravimetric estimation of total coagulable plasma proteins (no. 6), N in washings therefrom (no. 7), and the values obtained for total plasma-N $\times$ 6.25 (no. 8), and for protein-N $\times$ 6.25 and 6.6 respectively (nos. 9 and 10) for four separate pools of human plasma*

(No. 11 gives the values for total plasma-N $\times$ 6.25 expressed as % of the gravimetric determination. Nos. 12 and 13 give the values for protein-N $\times$ 6.25 and by 6.6 respectively, as % of the gravimetric determinations. Otherwise values as g./100 ml. plasma.)

No.		No. of determinations on each pool	Pool A1	Pool B	Pool C	Pool D	Mean of all pools
1	Total plasma-N	A	3	1.169	1.105	1.106	1.119 $\pm$ 0.038
2	Non-protein-N	B	3	0.036	0.026	0.016	0.024 $\pm$ 0.0067
3	Protein-N	A-B	3	1.133	1.079	1.056	1.087 $\pm$ 0.044
4	Globulin-N		3	0.414	0.399	0.395	0.396 $\pm$ 0.0020
5	Albumin-N		3	0.719	0.680	0.661	0.691 $\pm$ 0.0024
6	Gravimetric determination	C	2	7.42	7.28	7.72	7.35 $\pm$ 0.37
7	N in washings from C		2	—	0.058	0.024	0.042
8	Total plasma-N $\times$ 6.25	D	—	7.31	6.91	6.91	7.00 $\pm$ 0.18
9	Protein-N $\times$ 6.25	E	—	7.08	6.75	6.60	6.79 $\pm$ 0.30
10	Protein-N $\times$ 6.6	F	—	7.48	7.12	6.97	7.17 $\pm$ 0.20
11	D/C $\times$ 100	—	—	98.5	94.8	98.4	95.3 $\pm$ 3.51
12	E/C $\times$ 100	—	—	95.5	92.7	85.5	92.5
13	F/C $\times$ 100	—	—	102.2	97.8	90.3	98.0 $\pm$ 4.72

These observations are too limited to derive any definite conclusions, but they indicate that the N content of 'albumin' is greater than that of 'globulin'.

Table 2 gives the results as g. per 100 ml. for the distribution of the main N-containing fractions in plasma (nos. 1-5), the gravimetric estimation (no. 6), and N contained in the acetone, saline and water washings from the gravimetric determinations (no. 7). The number of determinations done on each pool is indicated in Table 2, the values given being the mean for each pool. It may be remarked that the duplicate estimations of the gravimetric determinations agreed within 2%. The values for N in the washings are the mean of two closely agreeing results in each case. Solely in the case of pool B was a biuret reaction obtained and even here this was only slightly positive. No. 8 shows the values obtained by using the conversion factor 6.25 for total plasma-N; nos. 9 and 10 give values which were obtained by multiplying 'protein-N' by 6.25 and 6.6 respectively. Nos. 11, 12 and 13 give the values of the N determinations multiplied by the several factors as percentages of the gravimetric determination. These values are referred to in the discussion of the results.

DISCUSSION

If the conversion factor 6.25 is to be used for the determination of plasma proteins by methods based on the N content it is necessary to find whether the proteins contain a constant 16% N, and if not, whether they contain a fixed amount of N such that a definite conversion factor may be used. The results shown in Table 1 indicate that the N content of total plasma proteins from different pools and involving in all 67 persons varies considerably. A mean value of 15.2% N may be taken, however,

which would give a conversion factor of approximately 6.6.

That the N content of human total plasma proteins is less than 16% is indicated in Table 3 where values collected from the literature are given. Values for the plasma proteins, either mixed or in various fractions, of other mammals are given by Adair & Robinson (1930 *a, b*), Robinson & Hogden (1941) and Block & Bolling (1945). In addition to the values given for total plasma or serum proteins some values, albeit few, are given for the N content of 'pure' plasma protein fractions. It will be seen that β -globulin, which forms approximately 13% of the plasma proteins (Cohn, Oncley, Strong, Hughes & Armstrong, 1944) contains only 14.8% N. The possibility of the presence of other protein fractions, yet to be isolated, must also be borne in mind.

For practical purposes and to obtain a 'fit' with the gravimetric determinations Table 2 shows that total plasma-N multiplied by 6.25 (nos. 8 and 11) or 'protein-N' multiplied by 6.6 (nos. 10 and 13) gives a reasonable correlation. The use of the classic factor 6.25 for protein-N gives too low a value (nos. 9 and 12). The correlation stated above would be even better if the result from pool C was neglected. This result is much higher than the others and it is interesting to observe that the N content of the washings is very much less than in the case of the other pools. That the protein factors for 'globulin' and 'albumin' are not the same is indicated in the results found here and from the values given in the literature (see Table 3). It is interesting that a good correlation between gravimetric and micro-Kjeldahl methods

Table 3. *Values reported for % N in human total plasma (P), or serum (S) proteins and in protein fractions derived therefrom*

(The values given are, in so far as can be ascertained, corrected for moisture and ash content. Jacobson's (1939) results are for serum samples derived from pathological cases (no mean value given).)			
g./100 g. of protein			
	Range	Mean	Reference
I. Total coagulable proteins			
P Direct coagulation	13.75-15.15	14.3}	Bierry & Vivario (1923)
Acetone treated	15.03-15.45	15.3}	
P Acetone treated	13.29-15.96	15.2	Present
S Direct coagulation and/or acetone treated	15.5- 16.0	15.7	Robinson & Hogden (1941)
S Washed with ethanol and ether	14.1- 15.9	15.2	Block & Bolling (1945)
S Acetone treated	14-15	—	Jacobson (1939)
II. 'Globulin' and its fractions			
'Globulin'	—	14.0	Murill, Block & Newburgh (1940)
'Globulin'	13.45-15.13	14.4	Present
β -Globulin (impure)	—	14.84	Brand, Kassell & Saidel (1944)
γ -Globulin	—	16.03	
Fibrinogen	—	16.9	
III. Albumin			
Crystalline	—	15.7 $\pm$ 0.2	Kendall (1941)
Crystalline	—	15.95	Brand, Kassell & Saidel (1944)
'Albumin'	15.2- 15.6	15.4	Present

(The values given are, in so far as can be ascertained, corrected for moisture and ash content. Jacobson's (1939) results are for serum samples derived from pathological cases (no mean value given).)

is obtained if 'globulin-N' is multiplied by 7 and albumin-N by 6.25. Differences in the relative amounts of 'globulin' and 'albumin' may well account for the individual discrepancies observed by Cook *et al.* (1945). The results found here are at variance with those of Robinson & Hogden (1941) who found a correlation between the micro-Kjeldahl method using the classic factor and the gravimetric method. Among the points of difference between their results and these may be noted the following. Most of their determinations were made on animal sera, two results only being obtained with human sera. They prolonged the coagulation time to 1 hr., and the washings from the proteins gave a positive biuret reaction. Finally on their own evidence the N content of the coagulated serum proteins was 15.7 %, a value giving a conversion factor of 6.36.

Total plasma proteins or even the globulin and albumin fractions as ordinarily isolated are mixtures and it is not to be expected that they would have a constant composition. Whilst the gravimetric method is as yet the most direct and accurate method, it may possess limitations quite apart from its laborious nature; heat coagulation may involve a slight hydrolysis of the protein proper, and a splitting of loosely bound prosthetic groups. A short coagulation time would therefore appear most suitable.

The following general conclusions may be drawn. First, that for all accurate work on the determination of plasma proteins the results should be expressed as 'protein-N' or as N precipitated by some specific reagent, e.g. acid tungstate, trichloroacetic acid or saturated sodium sulphate. The result so obtained would be less equivocal than in the form now usually given. If possible, the use of any specific factor should be avoided. Secondly, for all practical pur-

poses and where normal persons are concerned, total plasma-N multiplied by 6.25 gives a reasonably accurate result and one that is generally accepted as giving a comparative working value (Cook *et al.* 1945).

SUMMARY

1. The N content of four samples of the acetone-treated, heat-coagulated plasma proteins and three samples of 'globulin' from four pools of human plasma derived from 67 normal persons have been determined by the micro-Dumas and micro-Kjeldahl methods.

2. The N content (after correction for moisture and ash) for total plasma proteins had a range 13.3–15.9 %, mean 15.2 %; for 'globulin' the range was 13.5–15.1 %, mean 14.3 %.

3. The distribution of N in various plasma fractions has been determined, and the total proteins estimated by the gravimetric method. The results obtained by the latter method have been compared with the results derived by using different conversion factors for the N content.

4. While fair correlation is obtained by multiplying total plasma-N by 6.25 or 'protein-N' by 6.6, it is suggested that, as the plasma proteins are a complex mixture, values for all accurate work would be best expressed as g. N, in some particular fraction, per 100 ml. plasma.

This work owes much to the co-operation of Drs D. M. Keay and D. G. McIntosh, and Mr H. McDonald, of the staff of the Dundee Public Health Department, in collecting the blood samples and examining the persons involved. To the Medical Research Council the writer is indebted for a grant for expenses and for an honorarium to Miss A. M. Goodall who was responsible for much of the technical work. To her and to Miss M. B. Brown who assisted in the earlier stages he expresses his thanks.

REFERENCES

- Adair, G. S. & Robinson, M. E. (1930*a*). *Biochem. J.* **24**, 993.
 Adair, G. S. & Robinson, M. E. (1930*b*). *Biochem. J.* **24**, 1864.
 Bierry, H. & Vivario, R. (1923). *C.R. Soc. Biol., Paris*, **89**, 15.
 Block, R. J. & Bolling, D. (1945). *The Amino-acid Composition of Proteins and Foods*. Springfield: Thomas.
 Brand, E., Kassell, B. & Saidel, L. J. (1944). *J. clin. Invest.* **23**, 487.
 Chibnall, A. C., Rees, M. W. & Williams, E. F. (1943). *Biochem. J.* **37**, 354.
 Cohn, E. J., Oncley, J. L., Strong, L. E., Hughes, W. L. & Armstrong, S. H. (1944). *J. clin. Invest.* **23**, 417.
 Cook, R. P., Keay, D. M. & McIntosh, D. G. (1944). *Biochem. J.* **38**, xiv.
 Cook, R. P., Keay, D. M. & McIntosh, D. G. (1945). *Brit. med. J.* **2**, 456.
 Jacobson, B. M. (1939). *J. clin. Invest.* **18**, 485.
 Kendall, F. E. (1941). *J. biol. Chem.* **138**, 97.
 Lloyd, B. B., Cheek, E. B., Sinclair, H. M. & Webster, G. R. (1945). *Biochem. J.* **39**, xxv.
 Markham, R. (1942). *Biochem. J.* **36**, 790.
 Murrill, W. A., Block, W. D. & Newburgh, L. H. (1940). *J. biol. Chem.* **133**, 521.
 Peters, J. P. & Van Slyke, D. D. (1932). *Quantitative Clinical Chemistry Methods*. London: Baillière, Tindall and Cox.
 Phillips, R. A., Van Slyke, D. D., Dole, V. P., Emerson, K., Hamilton, P. B. & Archibald, R. M. (1944). *Copper Sulphate Method for Measuring Specific Gravities of Whole Blood and Plasma*. New York: John Macy, jun., Foundation.
 Plimmer, R. H. A. & Lowy, L. (1945). *Biochem. J.* **39**, xviii.
 Robinson, H. W. & Hogden, C. G. (1941). *J. biol. Chem.* **140**, 853.