affected the results with sulphanilamide. While increase in the inoculum within the range tested did not affect the action of 3-hydroxysulphanilamide it

had a marked effect in reducing the action of sulphanilamide both on solid and liquid media (Table 5).

REFERENCES TO APPENDIX 2

- Fleming, A. (1944). *Lancet*, 2, 620.
Harper, G. J. & Cawston, W. C. (1945). *J. Path. Bact.* 57, 59.

The Toxicity of Arsenic Solutions for Tissue Slices

By W. HUGHES AND G. A. LEVY

Departments of Biochemistry and Pharmacology, University of Edinburgh

(Received 12 July 1946)

When arsenic enters the blood stream from the lungs, it is rapidly fixed by the red cells. After a short latent period, haemolysis occurs. It has been generally held that no arsenic entering the body can escape reaction with the erythrocytes, and that poisoning of vital organs is thus due entirely to the substances liberated from the red cells on lysis. In recently published work, Levy (1946), however, has shown that the lethal dose of arsenic for mice, as measured by the arsenic absorbed in exposures to the gas causing 50 % mortality, which varies inversely with the gas concentration, can be as little as 0.67 mg./kg. At this value for the median lethal dose, arsenic is much more toxic than arsenites, the most toxic arsenic derivatives likely to be formed in the reaction between the gas and the erythrocytes.

In the same paper (Levy, 1946) experiments were described in which the arsenic content was determined of tissues from rabbits killed immediately after short exposures to arsenic. Although the tissues were freed from blood before the onset of haemolysis, certain organs were found to contain considerable amounts of arsenic. Liver had the highest arsenic content, while kidney and lung also contained comparatively large quantities. It was concluded that this arsenic represented a fraction of the arsenic which had been removed from physical solution in the plasma before it could be fixed by the red blood corpuscles. None of the arsenic fixed in this way is released until haemolysis commences (Thauer, 1934). Since it seemed possible that the effects of unchanged arsenic on vital organs might explain, partly at least, the high toxicity of the gas for the whole animal, it was decided to determine the toxicity of arsenic, in comparison with that of an arsenite, for surviving liver and kidney slices.

EXPERIMENTAL

To determine the toxicity of arsenic for the tissue slices, it was not considered adequate to shake them in suspension in Ringer solution with a gas phase containing the arsenic derivative, since, to compare the effects with those produced by arsenite solutions, assumptions would have been required as to the distribution of arsenic between the gas and liquid phases. A technique was accordingly developed for treating the surviving tissues with arsenic solutions in absence of a gas phase, under conditions resembling as closely as possible those obtaining *in vivo*.

Method

Rat-liver or kidney slices were treated for exactly 20 min. at 37° with the arsenic solution in a closed vessel completely filled with liquid. The solution was composed of a mixture of oxygenated Ringer solution (50–90 % of the total volume) and Ringer solution (10–50 % of the total volume) saturated with arsenic in hydrogen. Other slices from the same animal were treated with either pure Ringer solution or with a solution of arsenite in Ringer solution, both of which contained quantities of oxygen and hydrogen similar to those present in the arsenic vessel. Sufficient oxygen was always present to supply the tissue slices for 20 min. After this treatment, the oxygen consumption of the slices was measured by the Warburg technique (Dixon, 1934). Phosphate Ringer solution (Krebs, 1933) containing glucose (0.2 %) was used in all stages of the experiments.

Procedure

The gas towers. In all, four towers were required for saturating Ringer solutions with various gases. Each contained 200 ml. fluid. If these towers are numbered 1, 2, 3

and 4, their contents and the gases passed through them were as shown below.

Tower	Solution	Gas
1	Phosphate Ringer	O_2
2	Phosphate Ringer	$AsH_3 + H_2$
3	Phosphate Ringer	H_2
4	Arsenite in phosphate Ringer	H_2

The design of the arsine tower is shown at *A* in Fig. 1, and the other towers were similar, but without the attachments *B* and *C*. Arsine-hydrogen mixture was prepared by the action of zinc on an acidified arsenite solution in presence of copper catalyst, analytical reagents being used throughout, and the gas was collected in the gasometer previously described (Levy, 1946). The concentration was usually of the order of 20–40 mg. AsH_3 /100 ml. Before entering the tower, the gas was passed through 10 N-NaOH solution. Hydrogen was prepared from the same reagents without arsenite, and after washing with alkali was passed through towers 3 and 4, in that order. Oxygen was taken from a cylinder. The solution in tower 4 was prepared by diluting a measured volume (usually 2 ml.) of 0.1 M-sodium arsenite (made from A.R. As_2O_3) to 200 ml. with Ringer solution.

The gas was passed through each tower for 10 min. at a rate of 50 ml./min. and then for 2–3 hr. at 5 ml./min. before using the solutions. Complete saturation of the solution in tower 2 with respect to the pressure of arsine present required about 3 hr., but latterly we did not always wait for equilibrium to be attained, as samples of the solution were withdrawn for analysis immediately before treating the tissues. While liquid was being withdrawn from a tower, the gas exit tube was clipped to prevent entrance of air.

Sampling the solutions. To determine its arsine content, the solution in tower 2 was measured out by means of the burette attachment *B* (Fig. 1), through which the arsine-hydrogen mixture was led after leaving the tower. After the burette had been washed out several times with the solution, a volume of about 5 ml. was run into the sampling tube *C* (Fig. 1). A slow stream of nitrogen was passed for 30 min. through *C* into a wash-bottle containing 10 N-NaOH and three absorption tubes charged with 0.02 N- $AgNO_3$ solution.

Fig. 1. Tower for saturating solutions with arsine (*A*), with attachments for measuring out (*B*) and collecting (*C*) samples for analysis.

In early experiments, the flow of nitrogen was commenced before the arsine solution was measured out, and *C* remained attached to *B* throughout its passage. Latterly, it was found more convenient to detach *C* from *B* before passing nitrogen through the sample. The arsine solution was run into *C* with the nitrogen inlet and outlet tubes (*a* and *b*) closed by clipped rubber tubing, and *C*, complete with rubber stopper, was disconnected from the burette jet, the hole in the stopper being at once closed with a glass rod. This operation was easily performed if the burette tip and glass rod had been lubricated with glycerine. While arsine solution was entering, the pressure in *C* was released when necessary by opening the spring clip on the inlet (*a*) momentarily. The presence of a few drops of water in *C* acted as a trap for arsine in the inlet tube. This sampling procedure was found to yield as accurate and reliable results as that first used.

The arsenite formed in the $AgNO_3$ absorption tubes was titrated, after the addition of excess KI, with iodine in the presence of starch and bicarbonate. After all arsine had been removed with nitrogen, the solutions in the sampling tube *C* were examined for the presence of arsenite, invariably with negative results.

The arsenite content of the solution in tower 4 was, of course, known from its preparation. Titration with iodine showed no change in the arsenite content after passage of hydrogen for several hours.

Treatment of the tissue slices. The design of the vessels used for treating tissue slices with solutions from the gas towers is shown in Fig. 2. Before use, the capacity of each vessel was determined and it was then graduated in tenths of its total volume (60–70 ml.). For mixing, each vessel contained a hollow glass ball perforated at several points. Complete mixing of a coloured with a colourless solution could be achieved by dropping this ball the length of the vessel two or three times, but probably satisfactory mixing could be effected by movement of the tissue slices alone. The top of the vessel and the stopcocks were secured by elastic bands. Three vessels were required, one for treating the tissue with arsine, the second for arsenite treatment and the third for the control slices.

For treatment with arsine, tissue slices were placed in 2 or 3 ml. phosphate Ringer solution in the bottom of one of the vessels by means of long-handled forceps. The ball was then introduced and the top of the vessel attached. Oxygen was passed through the vessel till all air was displaced. With the capillary stopcock (*a*) open, oxygenated Ringer solution from tower 1 was run in through the wide-bore stopcock (*b*) till the solution reached the mark necessary for the desired dilution of the solution from tower 2, the vessel being always filled at least to the 0.5 mark with oxygenated Ringer solution. After samples of the arsine solution had been taken for analysis from tower 2, the burette was disconnected, stoppered and put aside. By means of a short piece of rubber tubing, well washed and

Fig. 2. Vessel for treatment of tissue slices in absence of a gas phase.

filled with solution from the tower, the arsine solution was run through (b) into the vessel till the liquid level just reached the capillary stopcock (a), care being taken to ensure that no bubbles were present before the vessel was completely filled. Both stopcocks were at once closed and the vessel inverted several times so as to mix the contents. Control and arsenite-treated slices went through the same process, the vessels being charged first with oxygenated Ringer solution and then with the solution in tower 3 or 4 respectively.

each experiment, and the figures for the Q_{O_2} (μ l. oxygen absorbed in 1 hr./mg. dried tissue) are in most cases averages for duplicate determinations. Our own figures for the Q_{O_2} of normal rat tissues set up in Warburg flasks directly after slicing were kidney 25 and liver 7.6. It will be seen that treatment of the control slices for 20 min. with a solution of oxygen and hydrogen in absence of an oxygen gas phase does not seem to have greatly affected

Table 1. *Poisoning of tissue slices by arsine and by arsenite*

No. of exp.	Control slices Q_{O_2} (μ l./mg./hr.)	Arsenite poisoned slices			Arsine poisoned slices		
		Concentration (10^{-4} M)	Q_{O_2} (μ l./mg./hr.)	Inhibition (%)	Concentration (10^{-4} M)	Q_{O_2} (μ l./mg./hr.)	Inhibition (%)
Rat kidney							
1	23.1	5	14.4	38	5.0	13.2	43
2	19.0	—	—	—	5.6	10.8	43
3	19.9	2	12.4	38	0.74	15.0	25
4	19.2	2	14.9	22	1.1	19.7	0
5	20.4	5	12.4	39	1.8	14.9	27
6	20.0	5	11.8	41	4.2	13.5	33
7	19.7	4	16.9	14	3.9	16.4	17
8	19.6	4	18.3	7	3.9	16.4	16
9	20.9	5	13.3	36	4.4	15.2	27
Rat liver							
10	7.2	2	4.8	33	1.5	2.3	68
11	5.8	5	3.0	48	2.2	1.7	71
12	7.4	2	5.0	32	0.72	5.4	27
13	6.9	5	3.3	52	1.0	4.0	42
14	6.3	1	4.3	32	1.7	2.8	56

All three vessels were set up in stands and shaken in a bath at 37° for 20 min. At the end of this period, the vessels were emptied, and the tissue slices, after brief washing with Ringer solution, were transferred to Warburg cups containing glucose-Ringer solution. The oxygen uptake was then determined. As a rule, sufficient tissue was placed in each vessel to charge two Warburg cups, and thus allow duplicate determination of the oxygen uptake.

In running it into the mixing vessel, no arsine was lost from solution. This was proved by blowing the contents of charged vessels with nitrogen into conical flasks, from which the nitrogen was led into $AgNO_3$ solution. After passing nitrogen through the conical flasks for 30 min. the figure for arsenite in the $AgNO_3$ solution was found to be within 10% of that calculated for arsine introduced into the mixing vessel. Further, when the oxygen displaced as the solution was run into the mixing vessel was passed through 1.5 ml. 0.02 N- $AgNO_3$ solution, no darkening occurred, although 0.5 μ g. AsH_3 could be readily detected in this way.

RESULTS

Figures obtained for the oxygen uptake of control and poisoned slices of rat liver and kidney are summarized in Table 1. The inhibition of liver oxygen uptake caused by different concentrations of arsine and arsenite is shown graphically in Fig. 3. Slices from the same animal were used throughout

their aerobic metabolism, and in very few cases was there any falling off in the oxygen uptake of the poisoned or control slices during the 70 min. they were shaken in the Warburg apparatus.

Fig. 3. Reduction of liver Q_{O_2} by arsine ($\odot$ — $\odot$) and arsenite ($\times$ — $\times$). Ordinate, concentration (10^{-4} M); abscissa, inhibition (%).

As regards the inhibition of oxygen uptake, arsine and arsenite were equally toxic for kidney slices (Table 1). In the case of liver, however, the molar concentration of arsine required for 50% inhibition in oxygen uptake was only one-quarter that of arsenite, as can be seen from Fig. 3. On treatment with arsine, but not with arsenite, liver slices underwent a marked colour change, assuming a greasy, yellow appearance.

DISCUSSION

Results obtained in previous work (Levy, 1946) suggested that some arsine entering the blood stream may escape reaction with the red cells long enough to be fixed by the body tissues. From the experiments described above, it appears that if it does reach these organs *in vivo*, arsine will have a toxic action on the liver and kidneys. Under the conditions of the experiments, arsine was as toxic as arsenite for kidney tissue, and considerably more so for liver. The action of unchanged arsine in solution in the blood plasma on the liver, and perhaps also other organs, may thus explain, in part at least, the high toxicity of this gas when it is rapidly inhaled. The change in the appearance of liver slices when treated with arsine suggests that a reaction with the gas analogous to the characteristic one which occurs with haemoglobin may take place in the cells of this organ. An examination of the effects of poisoning *in vivo* with arsine on certain enzyme systems, besides yielding valuable information regarding the mode of action, might provide further evidence for the view that it has a specific action on body cells. It is of interest to note that arsenite, which did not change the appearance of liver slices, had the same toxicity for this tissue as for kidney.

If one accepts the postulate that the toxicity of arsine for the whole animal is in part due to effects produced by the gas itself in the body tissues, the decrease in its toxicity with decreasing concentration inhaled (Levy, 1946) becomes readily explic-

able. As the rate of inhalation falls off, more of the gas will be fixed by the erythrocytes before it can be transported round the body, and thus transformed to less toxic arsenic derivatives. The nature of the arsenic-containing products of the reaction in blood has been studied by Graham, Crawford & Marrian (1946).

The technique described above may find general application in experiments in which it is desired to study the effect of known concentrations in solution of substances which are normally gases on the metabolism of surviving tissues.

SUMMARY

1. A technique was devised for treating surviving liver and kidney slices in absence of any gas phase with arsine solutions of known concentration.
2. The effect of arsine in reducing the oxygen uptake of rat kidney was identical to that of arsenite.
3. With liver, a 50% inhibition in the oxygen uptake was obtained with a lower concentration of arsine than of arsenite.
4. The bearing of these findings on the toxicity of arsine for the whole animal is discussed.

The authors are indebted to the Director-General of Scientific Research (Defence), Ministry of Supply, for permission to publish this work, to the Moray Research Fund of Edinburgh University for a grant towards the cost of apparatus, to Dr S. R. Elsdon for instruction in the Warburg technique, and to C. Bell for the manufacture of glassware.

REFERENCES

- Dixon, M. (1934). *Manometric Methods*. Cambridge University Press.
- Graham, A. F., Crawford, T. B. B. & Marrian, G. F. (1946). *Biochem. J.* **40**, 256.
- Krebs, H. A. (1933). *Hoppe-Seyl. Z.* **217**, 193.
- Levy, G. A. (1946). In the Press.
- Thauer, R. (1934). *Arch. exp. Path. Pharmac.* **176**, 531.

Histidine Deficiency in the Rat and its Effect on the Carnosine and Anserine Content of Muscle

By A. T. FULLER, A. NEUBERGER AND T. A. WEBSTER
The National Institute for Medical Research, London, N.W. 3

(Received 2 August 1946)

The original purpose of the work described in this paper was to establish whether histidine was essential to the maintenance of body weight in the adolescent and adult rat and to investigate the symptoms of such deficiency if it occurred. Since our work has been completed, two papers dealing with

the same subject have appeared. Albanese & Frankston (1945) have shown that histidine is required in the diet both by young and adult rats, if body weight is to be maintained. Maun, Cahill & Davis (1946) confirming this finding have noted a slight reduction in haemoglobin and plasma protein