The nucleotide-N and nucleoside- and free purine-N contents of I were respectively about 34 and 5 times those of II. Nucleotides isolated from I, when combined with amino-acids and B-factors found in I, duplicated the proliferation-promoting effect of I.

2. The nucleotide and nucleoside fractions were found to consist principally of adenine and guanine derivatives.

3. Spectroscopic evidence indicated a synthesis of materials absorbing at 2600 A. within the living, injured cells during the course of injury.

The investigations reported here were supported in part by a grant from E. I. du Pont de Nemours and Co. Present address of Dorothea G. Loofbourow, The Massachusetts General Hospital, Boston, and of Constance A. Yeats, U.S. Naval Air Base, Pensacola, Florida.

REFERENCES

- Cook, E. S. & Cronin, A. G. (1941). *Stud. Inst. Divi Thomae*, **3**, 203.
- Cook, E. S., Loofbourow, J. R. & Stimson, M. M. (1939). *Att. X. Congr. Int. Chim.* **5**, 26.
- Davidson, J. N. (1940). *Biochem. J.* **34**, 1537.
- Davidson, J. N. (1943). *Edinb. med. J.* **50**, 70.
- Englert, M. E. (1942). *Rep. Third Conf. Inst. Divi Thomae*. Cincinnati.
- Englert, M. E. (1945). *Stud. Inst. Divi Thomae*, **4**, 59.
- Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
- Kerr, S. (1940). *J. biol. Chem.* **132**, 147.
- Kerr, S. (1941). *J. biol. Chem.* **139**, 121.
- Kerr, S. & Blish, M. E. (1932). *J. biol. Chem.* **98**, 193.
- Hitchings, G. H. (1933). Dissertation, Harvard University.
- Holiday, E. R. (1930). *Biochem. J.* **24**, 619.
- Loofbourow, J. R. (1942a). *Biochem. J.* **36**, 631.
- Loofbourow, J. R. (1942b). *Biochem. J.* **36**, 737.
- Loofbourow, J. R. (1947). *Biochem. J.* **41**, 119.
- Loofbourow, J. R., Cook, E. S. & Stimson, M. M. (1938). *Nature, Lond.*, **142**, 573.
- Loofbourow, J. R., Dwyer, C. M. & Lane, M. M. (1940). *Biochem. J.* **34**, 432.
- Loofbourow, J. R., Englert, M. E. & Dwyer, C. M. (1941). *Nature, Lond.*, **148**, 113.
- Loofbourow, J. R. & Joyce, Lois (1941). *Nature, Lond.*, **148**, 166.
- Loofbourow, J. R. & Stimson, M. M. (1940). *J. chem. Soc.* p. 844.
- Loofbourow, J. R., Webb, A. M., Loofbourow, D. G. & Abramowitz, R. K. (1942). *Biochem. J.* **36**, 513.
- Mitchell, J. F. (1942). *Brit. J. exp. Path.* **23**, 285.
- Reader, V. (1927). *Biochem. J.* **21**, 901.
- Rogosa, M. (1944). *J. Bact.* **47**, 159.
- Stimson, M. M. & Reuter, M. A. (1945). *J. Amer. chem. Soc.* **67**, 2191.
- Webb, A. M. & Loofbourow, J. R. (1947). *Biochem. J.* **41**, 114.

The Quantitative Separation on the Micro-scale of Phenylarsenious Acid, Phenylarsonic Acid, Arsenious Acid and Arsenic Acid in Simple Solution and in Urine

By T. B. B. CRAWFORD

Departments of Biochemistry and Pharmacology, University of Edinburgh

(Received 26 August 1946)

It has been generally assumed that aromatic arsonic acids undergo reduction *in vivo* to the corresponding arsenoxides before producing their parasiticidal or toxic actions. All evidence for this is, however, of an indirect nature (Voegtlin & Smith, 1920; Yorke & Murgatroyd, 1930; Murgatroyd, Russell & Yorke, 1934; Thuret, 1938). It has also been suggested that arsenoxides are oxidized before excretion in the urine (Chance, Crawford & Levvy, 1945). Evidence as to whether or not inorganic arsenic derivatives are formed from organic arsenicals in the body is scanty and inconclusive. Definite information regarding the metabolism of organic arsenicals has awaited the development of suitable microchemical methods.

To investigate the changes undergone in the body by phenylarsenious acid and phenylarsonic acid, a

method was required whereby each constituent of a mixture of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid when present as their salts in biological material could be determined on the micro-scale. It was considered that such a method could be developed incorporating the procedure previously described (Crawford & Levvy, 1946) for the separation of the trivalent and pentavalent arsenic compounds, provided that extraction of the organic but not the inorganic arsenicals from aqueous solution could be effected by an immiscible solvent. The elaboration of a suitable procedure along these lines for pure solutions containing the four arsenicals, and its application, with some modifications, to urine, is described below.

EXPERIMENTAL

Preliminary investigations

Standard solutions of each arsenic-containing compound were prepared and checked as previously described by Crawford & Levvy (1946). All stages of the fractionations were carried out in separatory funnels and total arsenic in the different fractions was determined after digestion to remove organic matter (Levy, 1943). Arsenic determinations on the original solutions employed were carried out concurrently with the analyses of samples from the fractionations and the figures so obtained used in calculating the recovery in each fraction; by this procedure constant errors arising from the arsenic determination were to some extent eliminated. Simultaneously with every experiment controls for traces of arsenic in reagents were done and the experimental results corrected accordingly. The arsenic content of every fraction obtained in each separation was ascertained so as to be able to account for all the arsenic originally added. Unless stated to the contrary, experiments were done in triplicate and the results cited are mean values.

Since emulsions were formed with urine, extractions were carried out on dialysates. In some experiments the arsenicals were added to dialysates of normal urine. Rabbit urine was collected in flasks containing 1-2 ml. chloroform for 24 hr. and then dialyzed through 'Visking' synthetic sausage casing against 1-3 vol. distilled water in the refrigerator for 1 day. After the addition of the arsenicals, the dialysate was allowed to stand for 24 hr. in the refrigerator before use. Samples (50 ml.) of the arsenic-containing dialysate were used in the fractionations.

In other experiments, the arsenicals were added directly to urine. Known volumes (10-20 ml.) were dialyzed in the refrigerator for 24 hr. against 50 ml. distilled water containing a few drops of chloroform. Control experiments in which the dialysate and the dialysis sac together with its contents were analyzed for total arsenic at the end of 24 hr. showed this period sufficient for equilibrium across the membrane.

The development of the procedure for separation of the arsenicals in simple solution

Extraction of phenylarsenious acid from aqueous solution by organic solvents. Peroxide-free diethyl ether was found to remove phenylarsenious acid almost quantitatively from aqueous solution acidified to 6N with respect to H_2SO_4 , while phenylarsonic acid, arsenious acid and arsenic acid were extracted in negligible amounts under identical conditions (Table 1). The partition coefficient of phenylarsenious acid between ether and 6N- H_2SO_4 was found to be approximately 1.5 and thus several extractions with the solvent were necessary. However, even five extractions with equal volumes of ether always left 5-10% of the arsenical in the aqueous phase. This did not appear to be due to oxidation of the phenylarsenious acid since subsequent extraction with ethane-1:2-dithiol (0.4% v/v) in carbon tetrachloride (the dithiol reagent) removed most of the remaining arsenic.

In these experiments, 40 ml. batches of the arsenical solutions were acidified by the addition of 20 ml. 18N- H_2SO_4 and extracted with five 60 ml. portions of ether, shaking vigorously for 2 min. with each portion of solvent. The ether had been rendered peroxide-free by washing with

saturated FeSO_4 and then with water. After drying over CaCl_2 for at least 12 hr. it was distilled just before use. The combined extracts were evaporated down over distilled water (30 ml.), the last traces of ether being removed *in vacuo*.

Table 1. *Extraction of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid from acidified aqueous solution by peroxide-free ether*

As added ($\mu\text{g.}$)	As found in ether extracts ($\mu\text{g.}$)	Added As in first ether extract (%)	As added ($\mu\text{g.}$)	As found in ether extracts ($\mu\text{g.}$)
Phenylarsenious acid			Phenylarsonic acid	
77	71.6	56	91	2.7
78	73.4	59	87	1.7
84	75.6	58	93	1.9
83	75.0	63	90	1.8
85	76.5	51	8.8	Nil
86	79.2	61	8.0	Nil
85	78.2	—	Arsenious acid	
85	78.2	—	42	Nil
80	73.5	—	10	1.6
82	77.9	—	7.8	1.2
81	73.0	—	Arsenic acid	
8.8	7.7	—	45	0.9
7.2	5.5	—	11	0.9
			11	0.9
			Mixture of phenylarsonic acid, arsenious acid and arsenic acid	
			110	3.3
			110	Nil

If the ether was not freed from peroxide, partial oxidation of the arsenical occurred as could be shown by its failure to react with the dithiol reagent.

Strong acidification of the aqueous solutions containing phenylarsenious acid in amounts equivalent to 40-80 $\mu\text{g.}$ As appeared to favour the extraction of the arsenical by ether. From 0.2M-phosphate buffer, pH 7, 73% was extracted as against 82 and 93% from 0.01N and 9N- H_2SO_4 respectively.

Other solvents were examined under varying conditions in preliminary experiments. None of these, however, was suitable for separation of phenylarsenious acid from the other arsenicals, phenylarsonic acid, arsenious acid and arsenic acid. Benzene, chloroform, carbon tetrachloride and light petroleum (b.p. 40-60°) removed little or no phenylarsenious acid from aqueous solution. *n*-Butanol and caprylic alcohol, while removing phenylarsenious acid from acid solution, also extracted considerable amounts of the other three arsenic compounds.

Failure to extract phenylarsonic acid quantitatively from aqueous solution by organic solvents. Of the solvents examined, carbon tetrachloride, chloroform and ethyl acetate failed to extract this arsenical from acid solution. 2-Methylbutan-4-ol (primary *iso*-amyl alcohol), while extracting phenylarsonic acid also removed arsenic acid. Caprylic alcohol and *n*-butanol were unsuitable for separation purposes. Contrary to Morgan (1918), ether did not effect any extraction of this compound from acidified aqueous solution (Table 1).

Reduction of the pentavalent arsenicals. As a result of the preliminary work the following procedure for the quantitative separation of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid when present together in aqueous solution was considered. Phenylarsenious acid could be removed by extraction with ether and arsenious acid by extraction with the dithiol reagent (Crawford & Levvy, 1946) leaving in the aqueous phase the two pentavalent compounds. No organic solvent having been found suitable for the separation of the latter, it was planned to reduce these substances and to re-extract the solution with ether to obtain, in the form of phenylarsenious acid, the fraction of the arsenic originally present as phenylarsonic acid. Determination of the arsenic remaining in the aqueous layer, or in subsequent dithiol extracts of the latter would then give the amount of arsenic acid present. This proposed method required the development of a suitable technique for the quantitative reduction of at least phenylarsonic acid, without interference in the later stages of the fractionation, and which, for preference, could be carried out in a separatory funnel.

Klein & Vorhes (1939) recommend the use of a mixture of KI and $\text{Na}_2\text{S}_2\text{O}_3$ in acid solution for the reduction of arsenic acid. On examination, however, this method was found to be quite unsuitable for present purposes since it led to the formation of ether-soluble derivatives from both phenylarsonic acid and arsenic acid. This phenomenon was finally shown to be due to the presence of the thiosulphate. When approximately 90 μg . As as phenylarsonic acid or arsenic acid (in 6N- H_2SO_4) were treated with three equivalents of $\text{Na}_2\text{S}_2\text{O}_3$, extraction with ether removed almost all the arsenic in both cases.

SO_2 was next considered as a reducing agent. According to Kurtenacker & Fürstenau (1933), the reduction of arsenic acid by this reagent proceeds at maximal velocity in 0.2-0.26N- H_2SO_4 . Experiments showed that SO_2 acting for 45 min. in 0.23N- H_2SO_4 brought about the complete reduction of arsenic acid ($\equiv 100 \mu\text{g}$. As), the product being insoluble in ether but easily removed with the dithiol reagent. Under the same conditions, however, phenylarsonic acid was reduced to the extent of only 10%. Increase of the H_2SO_4 concentration before reduction to 6N, the concentration most suitable for the removal of phenylarsenious acid by ether, did not increase the percentage reduction of phenylarsonic acid and greatly decreased that of arsenic acid. The fact that reduction was partial in the case of phenylarsonic acid ruled out the use of SO_2 as a reducing agent.

Attention was again directed to the use in acid solution (6N- H_2SO_4) of KI, alone or in combination with some reagent other than $\text{Na}_2\text{S}_2\text{O}_3$. KI alone gave with arsenic acid a product insoluble in ether but completely extractable by the dithiol reagent provided sufficient of the reducing agent was present (130 mg. for 80 μg . As). From phenylarsonic acid, however, the completely quantitative formation of an ether-soluble derivative was not obtained with that amount of KI which gave complete reduction of arsenic acid. The iodide was added to the arsenical solution (volume 60 ml.) at an initial temperature of 40-60° in a separatory funnel. After cooling for 60 min. the solution was extracted with ether and, in the case of arsenic acid, with the dithiol reagent.

Quantitative formation of an ether-soluble product from phenylarsonic acid and of a product extractable by the

dithiol reagent from arsenic acid was obtained when sodium metabisulphite ($\text{Na}_2\text{S}_2\text{O}_5$) was added in addition to the KI. In the case of arsenic acid, the criterion that the reduction product should not be soluble in ether was satisfied (Table 2). In the preliminary experiments, in conformity with Klein & Vorhes's (1939) method of reduction,

Table 2. *Reduction of phenylarsonic acid and arsenic acid in acid solution with a mixture of potassium iodide and sodium metabisulphite*

Initial temp. of solution (°)	As added (μg .)	As removed from aqueous layer (%)	Extraction reagent
Phenylarsonic acid			
60	84	96	Ether
60	85	99	Ether
18	80	106	Ether
Arsenic acid			
60	78	Nil 100	Ether Dithiol
60	71	Nil 95	Ether Dithiol
17	71	101	Dithiol

the temperature of the acidified arsenical solution was approximately 60° at the time of addition of the KI. Further work, however, showed that the reduction of the pentavalent arsenicals went equally well at room temperature. The iodide (130 mg.) was added to the arsenical solution (60 ml.) in a separatory funnel 45 min. before the $\text{Na}_2\text{S}_2\text{O}_5$ (50 mg.), both reagents being in fresh aqueous solution. Extraction with ether, in some cases followed by the dithiol reagent, was done 15 min. later. A pinch of solid metabisulphite was added before extraction with each portion of ether. Reducing the amount of KI used in this process to one-tenth prevented complete reduction of either pentavalent arsenical.

Table 3. *Reduction of phenylarsonic acid and arsenic acid in acid solution with a mixture of potassium iodide and l-ascorbic acid*

As added (μg .)	As removed from aqueous phase (%)	Extraction reagent
Phenylarsonic acid		
85	95	Ether
87	102	Ether
Arsenic acid		
78	1 99	Ether Dithiol
73	2 94	Ether Dithiol

l-Ascorbic acid (50 mg.) also proved suitable in conjunction with KI (130 mg.) for the reduction of phenylarsonic acid and arsenic acid in experiments done at 60° (Table 3). *l*-Ascorbic acid alone failed to bring about reduction of either compound.

The product of the reduction of phenylarsonic acid with KI and $\text{Na}_2\text{S}_2\text{O}_5$ or *l*-ascorbic acid was more readily removed from acid solution by ether than was phenylarsenious acid. The addition of the reducing agents to phenylarsenious acid itself was found to render this substance much more readily extractable by ether. It is possible that the arsenic was extracted as phenyldiiodoarsine (Steinkopf & Smie, 1926).

Other reagents examined for the reduction of both phenylarsonic acid and arsenic acid but without success were formaldehyde, hydrazine sulphate and KBr.

Interference of ethane-1:2-dithiol in the separation of the pentavalent arsenicals after reduction. It was the practice, in the preliminary work, to use one 40 ml. and two 20 ml. portions of 0.4% (v/v) ethane-1:2-dithiol in carbon tetrachloride in the extraction of trivalent arsenic derivatives from acidified aqueous solution (60 ml.). This procedure left sufficient dithiol in the aqueous phase to give an ether-soluble compound with the reduction product of arsenic acid. This difficulty could be overcome either by reducing the quantity of the dithiol reagent and repeatedly extracting the aqueous phase with carbon tetrachloride to remove any residual dithiol, or by adding I_2 as a tincture, to convert the residual dithiol to diethylene tetrasulphide (Fasbender, 1888) and removing the excess I_2 by extraction with carbon tetrachloride. The addition of reducing agents in the next stage of the separation procedure did not appear to regenerate ethane-1:2-dithiol from the diethylene tetrasulphide. In practice it was decided to combine both methods.

Procedure for the quantitative separation of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid in simple solution

The whole fractionation is carried out on one sample of the arsenical solution (40–50 ml.) contained in a separatory funnel of suitable capacity, the stopcock and stopper of which give a tight fit without the use of lubricant.

Twenty ml. 18N- H_2SO_4 are added and the solution shaken vigorously for 2 min. on each occasion with

five 60 ml. portions of freshly distilled peroxide-free ether. The combined ether extracts are evaporated over 30 ml. distilled water, the last traces of ether being removed *in vacuo*. The aqueous residue is analyzed for total arsenic, which gives the figure for the phenylarsenious acid content of the original solution.

After the removal of the phenylarsenious acid by ether, the solution is extracted by shaking vigorously for 2 min. on each occasion with two 20 ml. portions of 0.4% (v/v) ethane-1:2-dithiol in carbon tetrachloride, followed by three extractions with 20 ml. portions of carbon tetrachloride. Between each fresh treatment with the dithiol reagent or with carbon tetrachloride, 5 ml. carbon tetrachloride are added to the funnel and run off without any shaking to wash through the stopcock and stem of the funnel. The dithiol and the carbon tetrachloride extracts and washings are combined and the carbon tetrachloride distilled off after the addition of 30 ml. distilled water. The arsenic determination carried out on the aqueous residue so obtained gives the figure corresponding to the arsenious acid content of the original solution.

Before proceeding with the reduction of the pentavalent arsenicals remaining in the aqueous solution, 1 ml. 1% (w/v) iodine in absolute ethanol is added. After 15 min. the solution is shaken with two 20 ml. carbon tetrachloride to remove excess iodine. These extracts may be discarded.

The reduction of the pentavalent arsenicals is then carried out at room temperature by the addition of 1 ml. 13% (w/v) colourless aqueous potassium iodide, followed, after 45 min., by 1 ml. 5% (w/v) freshly prepared sodium metabisulphite solution. Fifteen min. later, the extractions with ether and with the dithiol reagent and carbon tetrachloride are repeated. To ensure the maintenance of complete

Table 4. *Analyses of mixtures of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid in aqueous solution*

Phenylarsenious acid			Arsenious acid			Phenylarsonic acid			Arsenic acid			As found in aqueous residue ($\mu\text{g.}$)	Total As		
Added ($\mu\text{g. As.}$)	Found ($\mu\text{g. As.}$)	Error ($\mu\text{g. As.}$)	Added ($\mu\text{g. As.}$)	Found ($\mu\text{g. As.}$)	Error ($\mu\text{g. As.}$)	Added ($\mu\text{g. As.}$)	Found ($\mu\text{g. As.}$)	Error ($\mu\text{g. As.}$)	Added ($\mu\text{g. As.}$)	Found ($\mu\text{g. As.}$)	Error ($\mu\text{g. As.}$)		Added ($\mu\text{g.}$)	Found ($\mu\text{g.}$)	Recovery (%)
64.4	67.4	+3.0	6.1	14.1	+8.0	66.4	61.6	-4.8	6.3	5.6	-0.7	Nil	143.2	148.7	104
64.4	65.5	+1.1	6.1	13.9	+7.8	66.4	68.3	+1.9	6.3	5.7	-0.6	Nil	143.2	153.4	107
64.4	61.6	-2.8	6.1	13.2	+7.1	66.4	65.9	-0.5	6.3	6.1	-0.2	2.7	143.2	149.5	105
6.5	5.4	-1.1	61.3	66.2	+4.9	6.1	4.9	-1.2	67.2	61.7	-5.5	7.8	141.1	146.0	104
6.5	6.9	+0.4	61.3	65.8	+4.5	6.1	6.3	+0.2	67.2	64.7	-2.5	3.6	141.1	147.3	105
6.5	10.3	+3.8	61.3	65.4	+4.1	6.1	10.8	+4.7	67.2	62.4	-4.8	4.0	141.1	152.9	109
6.5	6.0	-0.5	12.2	13.2	+1.0	19.9	12.7	-7.2	26.9	21.4	-5.5	Nil	65.5	53.3	81
12.9	13.9	+1.0	18.3	20.8	+2.5	26.6	21.9	-4.7	6.3	Lost	—	Nil	64.1	—	—
19.3	22.1	+2.8	24.4	27.0	+2.6	6.1	4.2	-1.9	13.4	8.2	-5.2	Nil	63.2	61.5	97
25.8	21.1	-4.7	6.1	6.7	+0.6	13.2	11.7	-1.5	20.2	17.3	-2.9	Nil	65.3	56.8	87
25.8	21.2	-4.6	Nil	1.2	+1.2	6.1	6.3	+0.2	Nil	Nil	Nil	Nil	31.9	28.7	90
6.5	4.8	-1.7	Nil	1.0	+1.0	26.6	22.3	-4.3	Nil	Nil	Nil	Nil	33.1	28.1	85

reduction during the ether extractions, a small pinch of solid sodium metabisulphite is added to the solution prior to extraction with each portion of ether. The appearance of a brown colour in the ethereal layer, most noticeable when the extractions are carried out in bright sunlight, may be disregarded. Although the coloration is probably due to traces of iodine formed from hydrogen iodide in the ether by peroxides, it does not appear to interfere in the separation. Arsenic determinations on the ether and on the dithiol extracts give the figures for phenylarsenic acid and arsenic acid respectively in the original solution.

The aqueous phase on the completion of the fractionation should prove on analysis to be arsenic-free if only the four arsenicals specified were present in the original aqueous solution. The method may be curtailed slightly by omitting the final extractions with the dithiol reagent if it is assumed that residual arsenic in the aqueous layer after reduction and extraction with ether corresponds to the arsenic acid originally present.

Results for this procedure using aqueous solutions containing known amounts of the four arsenicals are shown in Table 4, each figure being the result for a single extract. The carbon tetrachloride used to remove excess iodine was also analyzed in each case but no arsenic was ever found in this fraction.

Application of the fractionation procedure to urine dialysates

The formation of ether-soluble compounds between pentavalent arsenicals (phenylarsenious acid and arsenic acid) and urine constituents. The fractionation procedure was unsuitable for urine dialysates owing to

Table 5. *Formation of ether-soluble complexes from phenylarsenious acid and arsenic acid in the presence of acidified urine dialysate*

Arsenical added to	As added ($\mu\text{g.}$)	As removed by ether extraction (%)
Phenylarsenious acid		
Urine	35	89
Urine	42	94
Urine dialysate	66	91
Urine dialysate	93	93
Arsenic acid		
Urine	34	85
Urine	42	92
Urine dialysate	66	110

the formation of ether-soluble complexes between the pentavalent arsenicals and urinary constituents as a result of which the arsenic corresponding to these compounds appeared in the phenylarsenious acid fraction (Table 5). Attempts to decompose these complexes without change in the valency of the

arsenic so that the fractionation could be carried out on one sample of solution were unsuccessful. It was found, however, that an intensification of the reduction process decomposed the complexes. Arsenious acid, or arsenic acid after reduction, was not extracted from urine dialysates by ether and therefore the organic arsenicals could be separated from the inorganic by ether extraction after reduction. Further, complex formation by the pentavalent compounds did not lead to interference in the extraction of the trivalent arsenicals from urine dialysates with the dithiol reagent (Crawford & Levvy, 1946). On the basis of these findings, recourse was made to a method of fractionation involving the use of two samples of the solution.

Fractionation procedure as applied to urine dialysates containing phenylarsenious acid, phenylarsenious acid, arsenious acid and arsenic acid. Two 40–50 ml. portions of the arsenic-containing urine dialysates are treated in the following manner. The first portion is analyzed after acidification, according to the method described above for pure solution but omitting the initial extractions with ether. The first dithiol extracts thus contain the arsenic corresponding to arsenious acid together with phenylarsenious acid. After reduction by the modified method described below, the arsenic corresponding to phenylarsenious acid and to arsenic acid is removed by extraction with ether and with the dithiol reagent respectively.

The treatment of the second portion of dialysate after acidification commences with reduction. The organic arsenic (phenylarsenious acid and phenylarsenious acid) is removed by ether extraction while the inorganic arsenic (arsenious acid and arsenic acid) is removed by subsequent extraction with the dithiol reagent. The figures for phenylarsenious acid and arsenic acid having been obtained from the fractionation of the first portion of dialysate, those for phenylarsenious acid and arsenious acid may be found by difference.

This modified method has the advantage that possible errors arising from the incomplete extraction of phenylarsenious acid by ether are eliminated since the arsenical is extracted from the dialysate only after the reduction process in which a more readily ether-soluble derivative is formed.

To effect reduction of the pentavalent arsenicals in urine dialysate 2 ml. 13 % (w/v) potassium iodide solution are added and the mixture heated to 55–65°. This heating can be done conveniently by holding the separatory funnel containing the solution in front of an electric radiator. The solution is allowed to cool for 45 min. and 2 ml. 5 % (w/v) sodium metabisulphite solution are added. The extractions with peroxide-free ether are commenced 15 min. later.

The evaporation of the carbon tetrachloride from dithiol extracts should be carried out in the presence

of 30 ml. 1% Na_2CO_3 solution to prevent loss of arsenic due to the volatility of the phenylarsenious acid derivative in steam. The presence of dilute alkali (10–20 drops 10N-NaOH in 30 ml. water) appears to prevent loss of arsenic during the removal of the solvent from the combined ether extracts.

various fractions agreed with the total added with an accuracy within the range of that of the method used for arsenic determination in each fraction (standard deviation 12%; Levvy, 1943).

Modification of the fractionation procedure was necessary for the analysis of urine dialysates owing

Table 6. *Analyses of mixtures of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid added to urine before dialysis, or to urine dialysate*

(Triplicate samples of the dialysate fractionated; averaged results cited.)

Phenylarsenious acid			Arsenious acid			Phenylarsonic acid			Arsenic acid			Total As		
Added	Found	Error	Added	Found	Error	Added	Found	Error	Added	Found	Error	Added	Found	Recovery
($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	($\mu\text{g. As}$)	(%)
Urine dialysate														
27.8	26.0	-1.8	14.4	18.1	+3.7	29.1	27.3	-1.8	15.2	14.7	-0.5	86.5	86.1	100
46.3	48.4	+2.1	23.2	24.2	+1.0	46.7	41.9	-4.8	115.0	113.2	-1.8	231.2	227.7	99
47.4	43.4	-4.0	17.6	20.3	+2.7	46.1	44.8	+3.7	19.0	17.6	-2.4	130.1	131.5	101
Urine														
35.8	33.3	-2.5	17.7	19.2	+1.5	35.2	35.3	+0.1	18.6	17.3	-1.3	107.3	105.1	98
44.8	43.0	-1.8	24.3	24.5	+0.2	43.5	45.8	+2.3	19.1	18.4	-0.7	131.7	131.7	100
38.3	30.6	-7.7	14.3	14.2	-0.1	37.2	39.0	+1.8	15.7	15.1	-0.6	105.5	98.9	94

Table 6 summarizes the results obtained by this procedure in experiments in which mixtures of the four arsenicals were added to urine or urine dialysates. Each experiment was carried in triplicate and the results were averaged before insertion in the table.

DISCUSSION

The fractionation of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid in simple solution was carried out on one sample of the solution in a separatory funnel. Phenylarsenious acid was extracted from the acidified solution by peroxide-free ether, arsenious acid being then removed by extraction with the dithiol reagent (Crawford & Levvy, 1946). Reduction of the arsenic remaining in the aqueous phase by the addition of potassium iodide along with either sodium metabisulphite or *l*-ascorbic acid was followed by a repetition of the ether and dithiol extractions to remove the arsenic corresponding to phenylarsonic acid and arsenic acid respectively. The method was examined over a range of 5–70 $\mu\text{g.}$ arsenic for each compound and the error in any fraction appeared to be constant at an average value of not more than 3 $\mu\text{g.}$ of arsenic. It is of interest to note that, with one exception, only traces ($> 4 \mu\text{g.}$) of arsenic were found in the aqueous residue from the separation and that the sum of the arsenic contents of the

to the formation in the acidified dialysates of ether-soluble complexes by the pentavalent arsenicals. Complex formation by the pentavalent arsenicals in urine dialysates at normal pH could not be demonstrated by ether extraction but the possibility that the complexes were present as ether-insoluble salts cannot be excluded. The fractionation was carried out with two samples of the dialysate, one of which was submitted to immediate reduction to break down the complexes formed by As^{V} . The error in the fractionation of urine dialysates was certainly not greater than that in the procedure for simple solutions.

SUMMARY

1. A method is described whereby the amounts of each constituent of a mixture of phenylarsenious acid, phenylarsonic acid, arsenious acid and arsenic acid in simple solution may be determined.

2. Some modification of the method was found to be necessary before it could be applied to urine dialysates since both phenylarsonic acid and arsenic acid were found to yield ether-soluble complexes with urine constituents.

The author wishes to express his gratitude to Dr G. A. Levvy for his helpful advice and encouragement. Thanks are also due to Miss M. Clark for technical assistance. A grant for expenses from the Medical Research Council is gratefully acknowledged.

REFERENCES

- Chance, A. C., Crawford, T. B. B. & Levvy, G. A. (1945). *Quart. J. exp. Physiol.* **33**, 137.
- Crawford, T. B. B. & Levvy, G. A. (1946). *Biochem. J.* **40**, 455.
- Fasbender, H. (1888). *Ber. deutsch. chem. Ges.* **21**, 1470.
- Klein, A. K. & Vorhes, F. A. (1939). *J. Ass. off. agric. Chem. Wash.* **22**, 121.
- Kurtenacker, A. & Fürstenau, E. (1933). *Z. anorg. Chem.* **212**, 289.
- Levy, G. A. (1943). *Biochem. J.* **37**, 598.
- Morgan, G. T. (1918). *Organic Compounds of Arsenic and Antimony*. London: Longmans, Green and Co.
- Murgatroyd, F., Russell, H. & Yorke, W. (1934). *Ann. trop. Med. Parasit.* **28**, 227.
- Steinkopf, W. & Smie, P. (1926). *Ber. deutsch. chem. Ges.* **59**, 1453.
- Thuret, M. J. (1938). *J. Pharm. Chim., Paris*, **28**, 22, 60.
- Voegtlin, C. & Smith, H. W. (1920). *J. Pharmacol.* **15**, 475.
- Yorke, W. & Murgatroyd, F. (1930). *Ann. trop. Med. Parasit.* **24**, 449.

The Amino-acid Composition of Certain Bacteria and Yeasts

BY JOAN C. FREELAND AND E. F. GALE

Medical Research Council Unit of Chemical Microbiology, Biochemical Laboratory, Cambridge

(Received 18 September 1946)

The amino-acid composition of bacteria has hitherto received little attention; recently Stokes & Gunness (1946) have reported some analyses on a range of bacteria though detailed results are not yet available. Camien, Salle & Dunn (1945) collected values from previous literature and extended our knowledge with regard to a variety of organisms. The analysis of protein hydrolysates by the use of specific decarboxylases (Gale, 1945, 1946) enables accurate values for certain amino-acids to be obtained on comparatively small amounts of material. The present investigation was undertaken in order to determine whether the amino-acid composition of bacterial protein varies with the growth conditions of the organism and also whether there is any marked difference between the amino-acid composition of Gram + and Gram - species.

To determine the amino-acid composition of the bacterial protein it is necessary to correct values obtained with acid-hydrolysates of intact organisms for free amino-acid carried down from the growth medium. Such free amino-acid may be carried down on the surface of the organism or in the internal environment of the organism (Gale, 1947). Taylor (1947) has shown that only Gram + organisms carry down large amounts of free amino-acid in the internal environment, Gram - species containing negligible amounts of free amino-acid.

EXPERIMENTAL

Organisms. The *Esch. coli* used in these experiments was obtained from the National Collection of Type Cultures (N.C.T.C.), No. 86; a Gram - organism N.C.T.C. No. 6578 was isolated by this Unit

(Gale & Epps, 1944) also the strains of *B. subtilis*, *Staph. aureus* (B) and *Strep. faecalis*; *Proteus vulgaris*, *Aerobact. aerogenes* and *Staph. aureus* (A) were given to us by the Department of Pathology, Cambridge; *B. brevis* was kindly sent by Dr R. J. Dubos; the three strains of yeast were obtained from the Carlsberg Laboratory through Prof. Winge.

Culture media. Yeasts were cultivated in nutrient salt solution containing 0.2% Difco yeast extract and 4% glucose. For the cultivation of bacteria, a tryptic digest of casein was used as basal medium with the addition of glucose and/or marmite when necessary (Taylor, 1947). *Esch. coli*, *Aerobact. aerogenes*, organism No. 6578 and *Proteus vulgaris* were grown in flasks containing 4.5 l. of medium for each experiment; the other organisms were grown aerobically in liquid media in Roux bottles lying flat. An amino-acid free medium was also used for *Esch. coli* and *Aerobact. aerogenes* and consisted of Stephenson's salt-mixture with the addition of either 1.0% sodium lactate or 1.0% glucose; this was incubated in Roux bottles.

Methods. Amino-acids were estimated by the manometric method using specific amino-acid decarboxylase preparations (Gale, 1945, 1946) and results expressed as amino-acid-N as percentage total-N determined as usual by the micro-Kjedahl method. After growth, the organisms were centrifuged out of the medium and thoroughly washed with water on the centrifuge. The washed organisms were then made up into thick suspension in water (30-50 mg. dry weight/ml.) and the suspension divided into two parts. One part was then hydrolyzed with hydrochloric acid (Gale, 1945) and total amino-acid determined on the hydrolysate. The