2. Evidence has been obtained which tends to show that folic acid (Mitchell, Snell & Williams, 1941) is not identical with pteroyl-glutamic acid (Angier *et al.* 1945, 1946) but that vitamin B₉ (Binkley *et al.* 1943) may possibly be more similar to the latter than the former.

3. From a consideration of the effects of two impure preparations, one from liver and the other from poppy-straw, there appears to be a connexion between physiological properties and source, since

the five preparations can be divided into two classes according as their sources are liver or plant tissue.

4. Synergism has been observed between folic acid and both thymine and orotic acid.

The author wishes to thank Roche Products Ltd. (Basle) for gifts of materials and for a grant in aid of the work, and Prof. F. C. Happold for helpful discussion and advice.

REFERENCES

- Angier, R. B., Boothe, J. H., Cosulich, D. B., Fahrenbach, M. J., Hutchings, B. L., Hultquist, M. E., Kuh, E., Mowat, J. H., Northey, E. M., Seager, D. K., Semb, J., Sickels, J. P., Smith, J. M., Stokstad, E. L. R., Subbarow, Y. & Waller, C. W. (1945). *Science*, **102**, 227.
- Angier, R. B., Boothe, J. H., Cosulich, D. B., Fahrenbach, M. J., Hutchings, B. L., Hultquist, M. E., Kuh, E., Mowat, J. H., Northey, E. M., Seager, D. K., Semb, J., Sickels, J. P., Smith, J. M., Stokstad, E. L. R., Subbarow, Y. & Waller, C. W. (1946). *Science*, **103**, 667.
- Binkley, S. B., Bird, O. D., Bloom, E. S., Brown, R. A., Calkins, D. G., Campbell, C. J., Emmett, A. D., Hogan, A. G., O'Dell, B. L. & Pfiffner, J. J. (1944). *Science*, **100**, 36.
- Binkley, S. B., Bird, O. D., Bloom, E. S., Brown, R. A., Emmett, A. D., Hogan, A. G., O'Dell, B. L. & Pfiffner, J. J. (1943). *Science*, **97**, 404.
- Bohonos, M., Hutchings, B. L. & Peterson, W. H. (1941). *J. biol. Chem.* **141**, 521.
- Bohonos, M., Hutchings, B. L., Slobodkin, N. H. & Stokstad, E. L. R. (1944). *Science*, **99**, 371.
- Chattaway, F. W. (1944). *Nature, Lond.*, **153**, 250.
- Chattaway, F. W., Dolby, D. E. & Happold, F. C. (1944). *Biochem. J.* **38**, xviii.
- Chattaway, F. W., Dolby, D. E., Happold, F. C., McMillan, M. & Waters, J. (1945). *Biochem. J.* **39**, xii.
- Dolby, D. E., Happold, F. C. & Sandford, M. (1944). *Nature, Lond.*, **153**, 619.
- Dolby, D. E. & Waters, J. (1944). *Nature, Lond.*, **153**, 139.
- Foster, J. C., Keresztesy, J. C. & Stokes, J. L. (1944). *Science*, **100**, 522.
- Hall, D. A. (1947). *Biochem. J.* **41**, 294.
- Hutchings, G. H., Falco, E. A. & Sherwood, M. B. (1945). *Science*, **102**, 252.
- Johnson, B. C. (1946). *J. biol. Chem.* **163**, 255.
- Johnson, E. R. & Mueller, J. H. (1941). *J. Immunol.* **40**, 33.
- Keresztesy, J. C., Rickes, E. R. & Stokes, J. L. (1943). *Science*, **97**, 465.
- Kreuger, K. & Peterson, W. H. (1945). *J. biol. Chem.* **158**, 145.
- Larsen, A. & Stokes, J. L. (1945). *J. Bact.* **50**, 219.
- Luckey, T. D., Briggs, G. M. & Elvehjem, C. A. (1944). *J. biol. Chem.* **152**, 157.
- Mitchell, H. K. & Snell, E. E. (1941). *Proc. nat. Acad. Sci., Wash.*, **27**, 1.
- Mitchell, H. K., Snell, E. E. & Williams, J. R. (1941). *J. Amer. chem. Soc.* **63**, 2284.
- Snell, E. E., Guirard, B. M. & Williams, J. R. (1942). *J. biol. Chem.* **143**, 519.
- Stokes, J. L. (1944). *J. Bact.* **48**, 201.
- Stokstad, E. L. R. (1943). *J. biol. Chem.* **149**, 573.

The Folic Acid Group of Growth Factors

2. THE EFFECT OF FOLIC ACID ON ACID PRODUCTION BY GROWING CULTURES OF *STREPTOCOCCUS LACTIS* R

By D. A. HALL, *Biochemical Department, School of Medicine, Leeds*

(Received 9 November 1946)

It has been stated by Williams (1944) that folic acid, the growth factor preparation for *Streptococcus lactis* R and some other lactic acid producing bacteria, isolated by him from spinach leaves, is not a single entity but may consist of a group of compounds of similar chemical composition, but

differing sufficiently to possess distinct physiological properties. Since it has also been suggested (Hall, 1947a) that none of the components of this complex is identical with preparations of similar activity isolated from liver, it appeared to be of interest to consider folic acid separately to deter-

mine whether it were unitary in nature, or if complex to examine any differences in activity between the various components.

METHODS

The medium, organism and inocula used in all the experiments reported below were as described previously (Hall, 1947*a*). Two preparations were employed: a 7.7% concentrate of folic acid isolated from spinach and kindly supplied by Prof. R. J. Williams and the synthetic material pteroyl-glutamic acid, claimed as identical with natural liver *L. casei* factor, prepared by the Lederle laboratories (cf. Angier, Boothe, Cosulich, Fahrenbach, Hutchings, Hultquist, Kuh, Mowat, Northey, Seager, Semb, Sickels, Smith, Stokstad, Subbarow & Waller, 1945, 1946). Both preparations were kept in stock aqueous solution containing 10 and 1 $\mu\text{g./ml.}$ respectively, and after adequate dilution, added before autoclaving.

In most of the experiments it was desirable to continue incubation for considerable periods of time in order that stationary conditions of acid production should be reached. It has been stated (Elvehjem & Tepley, 1945) that growth of *Strep. lactis* ceases if the pH value is allowed to drop below 5.2, and to obviate this it was necessary to add alkali to the cultures in order to maintain the pH at 7.4. In view of the different rates of acid production which were encountered during a complete incubation, a constant drop-wise addition was impossible; as a compromise alkali was added approximately every 24 hr. to bring the pH back to 7.4, using bromothymol blue as internal indicator. Sterile 0.05 N-NaOH was added aseptically, thus permitting progressive titration figures for the acid produced in any culture to be determined. Alkali as dilute as this was required in order that an accurate titration figure might be obtained. Although this entailed a threefold increase in volume before the end of the incubation, such a dilution was found to be immaterial since a series to which 40% (w/v) NaOH was added did not show any significant discrepancies in rate of acid production, nor in the ultimate value for the titration. Initially the pH of the medium which dropped to about 6.8 during the period of sterilization, was returned to 7.4 by the aseptic addition of alkali before inoculation; later this was obviated by arranging the reaction at 7.8, when the desired drop occurred during autoclaving without any change in activity of the medium.

The figures given for growth of the organism were obtained by a determination of the viable count by the dilution technique, inoculation of dilutions being made into 'Lab-Lemco' broth fortified by the addition of an acid hydrolysate of yeast. Strictly, this technique is not applicable to streptococci, because of morphological considerations. It was, however, shown that there was no change in average chain length between the beginning and end of a 14 days' incubation period. Hence it can be assumed that if the figures relating to growth are incorrect, they are so by a constant factor, and although absolute growth figures may not have much meaning, the changes in rate of growth will be adequately recorded.

Glucose utilization was estimated by determination of the residual glucose in a sample of the cell-free Seitz filtered medium. For this experiment 1 l. of medium was prepared, and samples withdrawn at intervals. Growth in these samples was immediately stopped by the addition of 0.1 ml.

of 0.1% (w/v) solution of 'merthiolate' (sodium ethyl-mercurithiosalicylate). After removal of the cells by Seitz filtration, glucose determinations were made by Benedict's method.

RESULTS

To determine the mode of action of the folic acid complex, acid production/time curves were obtained for growing cultures containing a wide range of concentrations of the factor. Fig. 1 shows the

Fig. 1. Acid production/time curves for the growth of cultures of *Strep. lactis* R containing various concentrations of folic acid. Curves in order from the left refer to cultures containing respectively 1, 0.1, 0.01, 0.001 and 0 $\mu\text{g.}$ of folic acid/ml. of medium.

type of result obtained. Given adequate incubation the ultimate acid production was independent of the initial concentration of folic acid but was equivalent to a total conversion of the glucose in the medium to lactic acid. The rates at which the acid was produced, however, initially varying in accordance with the concentration, changed after 60–90 hr. and if the glucose supply was not completely utilized, became equal to the rate at which acid was produced after a lapse of 7 days in cultures without added folic acid.

This apparent ability to dispense with added folic acid after 7 days' incubation was a consistent phenomenon. The rate of growth over the initial period in blank culture was dependent on the batch of medium used, possibly being conditioned by the purity of the casein acid-hydrolysate used. As can be seen from Table 1, however, the change to an increased rate of acid production occurred at about 160 hr. whatever the initial rate. The angles between the two portions of the curves are also independent of the initial rate of acid production, being in all three cases quoted in Table 1 equivalent to a change in rate of approximately 0.045 ml. 0.05 N-acid/10 ml. medium/hr.

Table 1. Acid production by cultures of *Strep. lactis R* without added folic acid or pteroyl-glutamic acid

(Acid production measured in ml. of 0.05 N-NaOH required to bring to pH 7.4 (bromothymol blue as internal indicator).)

Time (hr.)	Acid production		
	Culture 1 (ml.)	Culture 2 (ml.)	Culture 3 (ml.)
24	0.55	0.10	0.15
48	1.50	1.50	1.85
72	1.95	2.55	3.25
96	2.40	3.60	4.35
120	2.85	4.75	5.40
144	3.15	5.70	6.70
168	3.80	6.95	8.85
192	5.25	8.95	11.15
216	6.60	11.10	13.75
240	8.10	13.30	15.70
264	9.75	15.50	18.00
288	11.15	17.35	19.00
312	12.55	18.40	19.00

A series of curves, similar to those of Fig. 1, obtained by the addition of pteroyl-glutamic acid (Angier *et al.* 1945, 1946) is shown in Fig. 2. The

Fig. 2. Acid production/time curves for the growth of cultures of *Strep. lactis R* containing various concentrations of pteroyl-glutamic acid. Curves in order from the left refer to cultures containing respectively 0.001, 0.0001, 0.00001, 0.000001, 0.0000001 and 0 μ g. of pteroyl-glutamic acid/ml. of medium.

same independence of ultimate acid production on the initial concentration of the factor can be observed, but there is not the same parallelism between the curves for the lower concentrations and that for the blank culture. In the former series there was a pronounced alteration in rate between 24 and 90 hr. incubation, after which the rate common to the lower concentrations set in. When the added factor is pteroyl-glutamic acid, however, there was a gradual falling-off in rate over the whole period of incubation.

In order to determine whether there was any correlation between growth rate, glucose utilization

and acid production, a further series of experiments were undertaken in which the two former determinations were made in addition to the usual measurement of acid production. Medium (1 l.) was prepared and inoculated with about 5 million organisms in sterile water suspension. This volume of medium was aerated strongly (1 l./hr.) during the whole period of incubation, and samples were withdrawn at 24 hr. intervals. Table 2 gives the figures obtained for the various observations for a typical

Table 2. Growth and metabolism of *Strep. lactis R*

(Acid production measured in ml. of 0.05 N-NaOH/10 ml. culture required to bring to pH 7.4 (bromothymol blue used as internal indicator). Glucose content measured on samples removed after periods stated, and determined by Benedict's method. Viable count measured by serial dilution.)

Time of incubation (hr.)	Glucose remaining in culture medium (mg./ml.)	Acid production	Viable count (log value)
14	4.84	1.05	6.8
18	4.35	1.65	8.7
22	3.90	2.25	10.9
38	3.23	5.00	13.4
63	2.70	6.55	15.6
86	2.42	7.35	17.3

experiment. It will be noticed that between 24 and 90 hr. there was a falling-off in the rate of glucose utilization, which was mirrored by the change in viable count. The acid production figures also showed a change to the secondary constant rate, shown in Fig. 1. It is possible, therefore, that there is one mechanism responsible for carbohydrate metabolism during the early stages of growth, and another which functions later, and that acid production and growth although not strictly proportional to one another are definitely interconnected.

Attempts to separate the early and later activities of folic acid have been relatively successful, due to the discovery that quinoxaline has the power of inhibiting both growth and acid production of the organism. This compound, which may be considered as a xanthopterin nucleus with the nitrogen atoms of the pyrimidine ring replaced by carbons, appears to act as an antagonist to those growth factors containing a pterin system. Table 3 gives a series of acid production figures for cultures of *Strep. lactis R* containing quinoxaline and either folic acid or pteroyl-glutamic acid. It can be seen that both of these compounds, dissimilar in so far as they do not both show xanthopterin-like absorption spectra, were capable of reversing the inhibition caused by quinoxaline, due apparently to the pteridine, although non-xanthopteroic structure of pteroyl-glutamic acid. Large concentrations of quinoxaline were required for complete inhibition, but the

Table 3. *The effect of quinoxaline on acid produced by Strep. lactis R in the presence of folic acid or pteroyl-glutamic acid*

(Acid production is measured in ml. 0.05 of N-NaOH/10 ml. culture required to bring to pH 7.4. (Bromothymol blue used as internal indicator).)

Additions to basal medium			Incubation period (hr.)						
Quin-oxaline	Folic acid	Pteroyl-glutamic acid	18	42	66	90	114	138	162
μg./ml.			Acid production (ml.)						
—	—	—	—	1.2	2.25	3.40	4.60	5.50	6.40
1000	—	—	—	1.15	1.90	2.50	3.05	4.95	6.80
100	—	—	—	1.30	2.55	4.00	5.10	6.00	7.20
10	—	—	—	1.50	2.80	4.10	5.45	6.05	8.10
—	0.1	—	0.7	3.55	5.15	6.95	9.65	11.05	12.45
—	0.001	—	—	1.85	3.25	4.40	5.60	6.80	8.65
—	0.00001	—	—	1.60	2.90	4.25	5.40	6.60	7.85
—	—	0.01	1.0	8.50	13.80	18.45	19.00	19.00	19.00
—	—	0.0001	0.85	3.05	13.65	18.80	19.05	19.05	19.05
—	—	0.000001	0.30	4.65	7.50	9.65	11.50	13.05	14.85
1000	0.1	—	0.50	3.65	5.85	7.70	8.95	9.85	10.55
1000	0.001	—	0.25	0.35	0.75	1.40	2.70	4.70	7.10
1000	0.0001	—	0.20	0.30	0.70	1.35	2.65	4.80	5.85
1000	—	0.01	1.20	8.50	13.60	18.45	19.10	19.10	19.10
1000	—	0.0001	0.60	5.80	9.75	13.25	14.90	16.30	17.05
1000	—	0.000001	0.20	2.10	4.05	5.25	6.25	7.05	7.45
100	0.1	—	0.65	3.75	5.50	6.90	9.90	11.10	12.35
100	0.001	—	0.20	1.50	2.70	3.70	4.65	5.85	8.05
100	0.0001	—	0.15	1.35	2.45	3.55	4.25	5.90	6.40
100	—	0.01	1.20	8.90	14.70	18.50	19.00	19.00	19.00
100	—	0.0001	0.55	6.75	12.25	16.70	17.70	18.00	18.05
100	—	0.000001	0.30	3.75	5.95	7.10	8.70	10.00	11.35

results obtained are striking. Outside certain limits of concentration, either the inhibitory or the reversal effects may be paramount, giving either no growth and acid production or unimpaired activity, but between these limits, quinoxaline merely had the effect of impeding growth and causing a corresponding lag in acid production. Fig. 3 shows a series of curves relating acid production to time of incubation for cultures containing concentrations of quinoxaline and folic acid in this intermediate range. It will be seen that the quinoxaline merely depressed the activity during the period when the first rate of acid production (cf. Fig. 1) was operative. After a period depending on the concentration of both factor and inhibitor, acid production proceeded at an enhanced rate, but this alteration never persisted beyond the period after which, in the absence of quinoxaline, the secondary rate would supervene. Thereafter the acid production proceeded at the normal secondary rate, equal to that at which acid was produced in cultures not containing addenda. The only case in which there was evidence of an alteration in the time of onset of the secondary rate was when the quinoxaline was added to a culture without folic or pteroyl-glutamic acid. Under these circumstances acid production at the new rate began earlier.

Fig. 3. *The effect of the addition of quinoxaline on the acid production of cultures of Strep. lactis R containing various concentrations of folic acid. Additions in terms of mg. of quinoxaline and μg. of folic acid respectively as follows: A, Nil, Nil; B, 2.5, Nil; C, 2.5, 0.00001; D, Nil, 0.00001; E, 2.5, 0.001; F, Nil, 0.001.*

DISCUSSION

There have been various indications of the possibility of folic acid being not a single substance but a group of closely related compounds with similar chemical but differing physiological activity. It has been difficult to reach any decision on this since there has been no standard simple substance with which folic acid could be compared. The preparation of synthetic pteroyl-glutamic acid, claimed to be identical with the liver *L. casei* factor by Angier *et al.* (1945, 1946), provided this criterion, since it may be assumed that the sole active substance present in this synthetic material is pteroyl-glutamic acid.

In view of this possibility of comparing folic acid with the pteroyl-glutamic acid, it is interesting to notice the differences between the acid production/time curves at various concentrations which are apparent from Figs. 1 and 2. The pteroyl-glutamic acid gives a series of curves, the slopes of which gradually fall away but which progress until the whole of the glucose supply is utilized, when a stationary state is reached. There is no abrupt change in rate such as occurs with folic acid after about 90 hr. incubation. One explanation of the change in rate of acid production observed after 90 hr. might be that folic acid is utilized stoichiometrically, and built *per se* into the structure of the developing cells with the concomitant expenditure of energy derived from the dismutation of the carbohydrate supply. This would of course mean that the amount of acid ultimately produced would be directly proportional to the initial concentration of folic acid, and would give rise to curves with plateaux at different levels. It would then have to be assumed that synthesis occurs thereafter at a constant rate causing the curves to rise at equal slopes until the whole of the carbohydrate source is utilized. This hypothetical onset of synthesis at the end of 90 hr., about 70 hr. before a similar effect is observable in the blank cultures, is the weak point in the hypothesis since there is apparently no reason why the presence of folic acid should induce an earlier onset of synthesis. This is borne out by the fact that the pteroyl-glutamic acid, although sufficiently different in physiological activity to act synergistically with folic acid (Hall, 1947a), is active in inducing similar degrees of acid production without a marked change of rate after 90 hr. Hence the possibility of folic acid being stoichiometrically utilized may be disregarded and other hypotheses considered.

Another possibility is that folic acid is in effect a mixture of two compounds, one of which is active during the earlier period of growth, the other later, and that during both these reactions the active

material functions as part of an enzyme system and not as a substrate. In other words, although the rate of acid production is proportional to the concentration, the ultimate degree is entirely independent of this factor. Of these two components, it appears that the organism is only able to synthesize that which is active during the later stages of growth. The question of why the rates are identical during the second phase, no matter what the initial concentration, may be answered by assuming that the concentration of this component is sufficiently above the threshold value required for constant catalytic effect to obviate any proportionality.

Quinoxaline has been shown to have certain power as an inhibitor of the growth of *Strep. lactis* R. Large quantities are required but the results obtained with it are striking. The almost universal inhibition of the first of the two hypothetical reactions of folic acid, without any alteration in the secondary rate of acid production, may be significant. Quinoxaline was chosen as an inhibitor, on the assumption that the observations of both Woolley (1944) and Madinaveitia (1944) regarding the replacement of nitrogen atoms in adenine and riboflavine nuclei to give benzimidazole and a mepacrine-like structure, respectively, could be extended to those compounds containing xanthopterin-like structures. The activity of quinoxaline may not be indicative of the presence of a pterin structure, however, since histidine has a similar power of reversing the inhibition (Hall, 1947b). It seems possible, however, that the converse may apply. In other words any preparation which is active as a growth factor for *Strep. lactis* R but will not reverse the effect of the inhibitor may be assumed not to contain the structure to which the inhibitor is analogous. Hence, while for other reasons it is possible to say that the complete folic acid complex contains a pterin-like structure (Mitchell, 1944), it is also possible to say on the above basis that the second hypothetical component does not, and hence the former must be pterin-like in nature. The evidence may indicate the presence in folic acid of two components, the first of which, associated with the early period of active growth and acid production, may be pterin-like in nature, and hence allied to pteroyl-glutamic acid (although not identical, cf. Hall, 1947a), while the second, associated with the later stages of growth, does not contain a pterin-like structure. It is interesting to notice that the 'folic acid' produced by resting cells of *Strep. lactis* R from the substance obtained by Foster, Keresztesy & Stokes (1944) does not have the same properties as folic acid, being very unstable, and may be identical with the second component suggested above, which it has not been found possible to detect in the cell-free culture filtrate, a fact which may be due to its instability.

SUMMARY

1. Folic acid and pteroyl-glutamic acid have been examined for their ability to affect acid production by *Strep. lactis* R during a long period of incubation.

2. Evidence has been obtained pointing to the existence in folic acid of two factors with activity in the early and later phases of growth and acid-production.

3. Quinoxaline can act as an inhibitor of the growth of *Strep. lactis* R, and both folic acid

and pteroyl-glutamic acid can reverse this inhibition.

4. Examination of acid production in the presence of intermediate concentrations of folic acid and quinoxaline is consistent with the presence of a pterin-like structure in the component responsible for early acid production, and not in the postulated component which is responsible for later activity.

The author wishes to thank Roche Products Ltd. for gifts of materials and for a grant in aid of this work, and Prof. F. C. Happold for helpful discussion and advice.

REFERENCES

- Angier, R. B., Boothe, J. H., Cosulich, D. B., Fahrenbach, M. J., Hutchings, B. L., Hultquist, M. E., Kuh, E., Mowat, J. H., Northey, E. H., Seager, D. K., Semb, J., Sickels, J. P., Smith, J. M., Stokstad, E. L. R., Subbarow, Y. & Waller, C. W. (1945). *Science*, **102**, 227.
- Angier, R. B., Boothe, J. H., Cosulich, D. B., Fahrenbach, M. J., Hutchings, B. L., Hultquist, M. E., Kuh, E., Mowat, J. H., Northey, E. H., Seager, D. K., Semb, J., Sickels, J. P., Smith, J. M., Stokstad, E. L. R., Subbarow, Y. & Waller, C. W. (1946). *Science*, **103**, 667.
- Elvehjem, C. A. & Tepley, L. J. (1945). *J. biol. Chem.* **157**, 139.
- Foster, J. C., Keresztesy, J. C. & Stokes, J. L. (1944). *Science*, **100**, 522.
- Hall, D. A. (1947a). *Biochem. J.* **41**, 287.
- Hall, D. A. (1947b). *Biochem. J.* **41**, 299.
- Madinaveitia, J. (1944). *Biochem. J.* **38**, xxvii.
- Mitchell, H. K. (1944). *J. Amer. chem. Soc.* **66**, 274.
- Williams, J. R. (1944). *Abstr. Pap. Amer. Chem. Soc.* 108th Meeting, p. B 14.
- Woolley, D. W. (1944). *J. biol. Chem.* **152**, 225.

The Folic Acid Group of Growth Factors

3. HISTIDINE AND ITS RELATIONSHIP TO 'FOLIC ACID'

By D. A. HALL, *Biochemical Laboratories, School of Medicine, Leeds*

(Received 9 November 1946)

Since 'folic acid' was first described by Mitchell, Snell & Williams (1941) there have been repeated attempts to identify the activity of the natural preparations with compounds of known composition. On the basis of the absorption spectrum of the material, attention was drawn to the connexion between folic acid and xanthopterin (Mitchell, 1944). The work of Totter & Day (1943), Totter, Shukers, Kolson, Mims & Day (1944) and Totter, Mims & Day (1944), claiming the alleviation of nutritional cytopenia in monkeys, by the use of folic acid and to a certain extent xanthopterin, did much to establish this belief in the identity of the two compounds. Mitchell & Williams (1944), however, showed that xanthopterin could not replace folic acid in the growth of *Strep. lactis* R nor was folic acid as effective as xanthopterin in fish anaemia. It was also shown that of a large group of compounds with a high nitrogen content, only thymine was able to replace folic acid for *Strep. lactis*. This fact has been proved and examined by

both Stokes (1944) for folic acid and Kreuger & Peterson (1945) for vitamin B₁₂, the former claiming that thymine does not directly replace folic acid, but that a thymine-like compound is the product of a reaction in which folic acid plays a part. On the other hand, in view of certain of his results, and those reported by the present author in a previous paper (Hall, 1947a) it appears more likely that if thymine is directly connected with folic acid, it may more readily be supposed to be a precursor than a product of folic acid activity. Chattaway (1944) examined a number of pyrimidine derivatives as growth factors for *L. casei*, and showed that orotic acid was active in this respect. It has also been shown (Hall, 1947a) that orotic acid, although not active alone for *Strep. lactis*, showed synergism with folic acid, as also did thymine. This tended to support the other possible conclusion that the active pyrimidine derivatives do not replace folic acid, but are in effect offering an alternative route for metabolism.