Photo by Lafayette, Ltd.

J. A. GARDNER

Obituary Notice

JOHN ADDYMAN GARDNER, 1867–1946

John Addyman Gardner, who for many years past has been a familiar and beloved figure at nearly all Meetings of the Biochemical Society, passed away on 13 May 1946. The surviving original members of this Society will recall the part played by him in its foundation. Many will appreciate the long and able service he gave to the administration of the Society, and most of the present members will feel the loss of a kind and sympathetic friend.

Born in 1867 at Bradford, Gardner attended that city's Grammar School, now famous for its many distinguished pupils. Those qualities of kindness and geniality, of industry and enthusiasm, which have always been associated with Gardner, seem to have been prominent at Bradford. Maybe the Yorkshireman has something more than an average claim to these. It was not uncomplimentary on the part of his schoolfellows to have nicknamed him Bull Gardner. He appears to have excelled on the classical side, and it was not until after he had been elected to an open scholarship at Magdalen College, Oxford, in 1886, and after a period of residence there that he gave up his classical studies for chemistry, an example perhaps of his general ability and versatility. Amongst his fellow-students studying chemistry in the School of Natural Science, he soon became as popular as he had been at Bradford. Amongst his friends at Oxford there appeared many of the eminent chemists of his day. It was during his undergraduate days that he became attracted towards chemical research, an occupation which undoubtedly claimed by far the greater part of his activities during his lifetime. As an undergraduate he was for some time President of the Oxford University Junior Scientific Club, and was engaged upon other social activities amongst his scientific colleagues.

Gardner was awarded a Demyship while at Magdalen College and also the Daubeny Medal. In 1889 he graduated with first class honours in the School of Natural Science. Proceeding to Heidelberg, he completed some research under Victor Meyer, who, from a report now in the writer's possession, appears to have been yet another impressed with Gardner's ability to carry out research work "*mit grossem Fleiss und Eifer*". In 1891, Gardner returned to Oxford as Demonstrator in Chemistry. For three years, in collaboration with J. E. Marsh, he worked on the constitution of the terpenes (*J. chem. Soc.* 1891–94), a task of no mean order, since in those days the extraction of the required terpenes from suitable plants

was often a necessary preliminary to such investigations.

In 1894 Gardner was appointed Lecturer in Chemistry and Toxicology at St George's Hospital Medical School, where he continued research on the terpenes, studying the bromination and oxidation products of camphene, camphoric and camphopyric acids. In those days the greater part of the hospital's clinical analyses and investigations were carried out under Gardner's supervision. The Medical School of St George's Hospital also provided pre-clinical courses for the London University Preliminary and Intermediate Examinations. Teaching and examining then occupied a large part of his time, and he became actively engaged on different Examination Boards. His frequent secondary school inspections were useful and popular. His interest in pre-clinical education caused him to take an active part in the introduction of a course of study and examination in organic chemistry in the London University 2nd M.B. To have secured such recognition of the growing importance of biochemistry was probably Gardner's first effort to advance this subject.

In 1905 Gardner was placed in charge of the chemical section of the Physiological Laboratory, which was attached to the London University side of the Imperial Institute at South Kensington, and known more generally as the Waller Laboratory, after its Founder and Director. Although retaining his position as Chemist to St George's Hospital, Gardner's interests were now directed more to South Kensington, where laboratory facilities for research were on a liberal scale. In these surroundings, Gardner, in collaboration with Buckmaster, undertook a laborious study on chloroform anaesthesia. An accurate method for the estimation of chloroform in blood was devised, and the anaesthetic and lethal quantities of chloroform were ascertained by a series of experiments on the cat. The rates of uptake and elimination and the function of the red corpuscles during anaesthesia were also investigated (*Proc. Roy. Soc.* 1906–7). It was at this time that Gardner began those researches on the sterols with which his name has become associated. Between 1908 and 1914, there appeared a number of communications under the title of 'The Origin and Destiny of Cholesterol in the Animal Organism'. The first with Dorée (*Proc. Roy. Soc.* 1908) referred to the discovery by Bondzynski that coprosterol, a reduction product of cholesterol, was excreted in human faeces, but

that the horse excreted hippocoprosterol. Pure coprosterol was prepared and a study of the faeces of the horse and other herbivora showed that hippocoprosterol was common to all, and that it originated from the grass on which the animals were fed.

Gardner and Dorée later showed that the cholesterol of the bile in the dog was reabsorbed, and that whether cholesterol or coprosterol was excreted depended largely on the nature of the diet. The coprosterol in faeces was shown to have its origin in the cholesterol of the diet. Coprosterol was not found in herbivora, and it was readily obtained from the faeces of a dog fed on a brain diet rich in cholesterol.

Gardner and Dorée prepared coprostenone and its derivatives, and by the action of sodium amyloxyde obtained the epimer of coprosterol which they described as ψ -coprosterol. Work on the isomeric reduction product of cholesterol, referred to as β -cholestanol, was included. The action of ozone on cholesterol and cholestenone led to the opening up of one of the rings with the formation of an acid of lower carbon content.

With Ellis, Gardner showed that the cat excretes cholesterol and coprosterol, mostly the latter on a meat diet. With Fraser, he showed that the degree of inhibition of the haemolytic effects of saponin was due to the presence of cholesterol or phytosterol (as had been observed by Windaus), and followed the cholesterol content of the diets fed to animals. These authors found in the blood of rabbits fed on bran about 80 mg./100 ml. total cholesterol, of which only about one-sixth was in the form of esters. This amount varied slightly on diets free from or enriched with cholesterol. It was in this communication that Gardner and Fraser first utilized Windaus's discovery of the insoluble compound formed by cholesterol with digitonin; they investigated this compound in its application to cholesterol estimations and verified Windaus's observations on the accuracy attending its use in quantitative work.

Gardner commenced some experiments with a view to ascertaining how far cholesterol could be synthesized by the animal. He showed, with Ellis, that newly hatched chicks contained a little less, certainly not more, cholesterol, than could be accounted for in the original eggs. Some evidence that cholesterol was synthesized in adult subjects was obtained by Gardner and Fox (*Proc. Roy. Soc.* 1921) and subsequent work by other investigators has done much to remove any doubts that such synthesis takes place. In a series of experiments it was noted that a man, weight being constant, excreted slightly less coprosterol than could be accounted for by the cholesterol content of food intake. During an attack of influenza for the periods when the subject was losing weight, an excess of coprosterol was ex-

creted, especially during a period when the loss of weight was heaviest (*Proc. Roy. Soc.* 1912).

At the same time Gardner continued to interest himself in chloroform anaesthesia. His attention was drawn to publications in which it was asserted that carbon monoxide was present in normal blood and especially in the blood of chloroform-anaesthetized animals. His previous elaborate researches with Buckmaster (quoted above) did not countenance these statements. By further careful and exact experiments with Buckmaster (*Proc. Roy. Soc.* 1909) he showed that the carbon monoxide, claimed to be found in the blood of anaesthetized animals, arose from the action of the caustic alkali on the chloroform content of the blood gases during carbon dioxide absorption; and the assertions that carbon monoxide was present in normal blood were also refuted.

Following this work Gardner and Buckmaster recorded some accurate observations on the venous and arterial blood gases of the normally breathing and of the anaesthetized cat. Working with hermetically sealed blood-gas pumps of the Töpler, Barcroft and Bone types, it is probable that some of the most accurate figures ever obtained in this type of analysis were made. Certainly, Gardner himself never undertook more laborious work (*J. Physiol.* 1910; *Proc. Roy. Soc.* 1910), in which one notes disagreement with the findings of the Chloroform Committee of the British Medical Association for 1910, to the effect that during chloroform narcosis the blood retains unimpaired its normal capacity of absorbing oxygen, and that the diminished concentration of this gas was solely due to slowing of respiration. According to Gardner and Buckmaster, diminished oxygen in anaesthetized blood was due to interference by the anaesthetic with the normal function of the red blood corpuscles. Accurate blood-gas analysis showed that the replacement of air by oxygen in respiration gave little increase of oxygen in the blood and no change in average carbon dioxide content.

Gardner was interested in the chemical constitution of cholesterol, and with Dorée (*loc. cit.*) he identified coprosterol as a secondary alcohol by its oxidation to coprostenone. With Williams (*Biochem. J.* 1913) he described further oxidation products of coprosterol and of the ketone. With Ellis (*Biochem. J.* 1918) he described some of the oxidation products of ' β -cholestanol' (cholestanol).

During the 1914-18 war, Gardner's previous acquaintance with the pungent lachrymatory substance chloropicrin led him to recommend its use in gas warfare, as an efficient but not too inhuman weapon. This volatile substance was produced in large quantities, under Gardner's supervision. During the same period he was responsible with Sir F. G. Hopkins for the comparison of the relative digestibilities of breads made from different flours.

Not content with these wartime responsibilities, he took an active share in the volunteer Police Force as a Special Constable.

Soon, after the 1914-18 war the Physiological Laboratory at South Kensington closed.

Appointed Reader in Physiological Chemistry, he was lecturer in Organic Chemistry at the London School of Medicine for Women, but his major activities were still directed towards research, now largely confined to the Biochemical Laboratories at St George's Hospital Medical School. He was critical of methods described by different investigators for the determination of cholesterol and its esters. With Williams (*Biochem. J.* 1921) he showed that cholesterol esters developed colour with the Liebermann reagent at quite different rates compared with cholesterol, and never regarded the colorimetric method as accurate for total cholesterol. Some useful information concerning errors in gravimetric determinations of ester cholesterol, particularly those due to incomplete hydrolysis, was obtained in conjunction with Fox (*Biochem. J.* 1921, 1924).

Communications with Gainsborough (*Biochem. J.* 1927-38) dealt, for the most part, with cholesterol and its esters in plasmas; in these experiments are to be found some of the most accurate determinations recorded for the cholesterol and cholesterol ester contents of normal plasma together with the limit of variations on 50 subjects examined. Sources of error in such determinations were ascertained, and errors due to the difficulty of separating sterols from protein matter were also recorded. With Murray, Gardner and Gainsborough noted an error due to shrinkage of the corpuscles by the addition of certain anti-coagulants, with as much as a 4% increase in the plasma volume.

Gardner and Gainsborough drew attention to the large amount of cholesterol found in the total blood of a man, as compared with the amount to be found in an average daily diet. The whole of the latter, if taken into the blood stream, might effect an increase within the normal variations of the sterol content.

During this period (*Brit. med. J.*, *Lancet*, *Quart. J. Med.* 1930-38) Gardner and Gainsborough were also engaged in studying the sterol content of plasma under different normal and pathological conditions. Hypercholesterolaemia was not produced by a simple cholesterol-rich meal, though it would appear on continuing such diets when there was a greater increase of cholesterol ester than of cholesterol in the plasma. So also could hypocholesterolaemia be occasioned on a prolonged cholesterol-low diet, but this resulted in a lowering of ester and not of free cholesterol. The influence of a simple diet on the plasma cholesterol is more marked by variations in the type of food than by variation in cholesterol content. This may result in an increase or decrease in the plasma content, and these variations are

regarded as due to endogenous metabolic changes in which cholesterol plays a part. Further work demonstrated the close attachment of cholesterol to the euglobulin fraction of plasma proteins, the presence in normal urine of minute quantities of cholesterol which were considerably increased in some forms of nephritis, and the reversal of the normal ratio of ester to free cholesterol in the plasma during pregnancy. The normal ratio of approximately 2:1 became inverted between the 24th and 32nd week of pregnancy but returned to normal before termination; at the same time the hypercholesterolaemia noted by other observers was a less constant finding. Studies were also made on nephritis, hyperthyroidism, myxoedema and on hepatic and biliary disease. In the latter it was demonstrated that partial or complete biliary obstruction was followed by a marked lowering of the ester cholesterol of the plasma, but that later if obstruction was complete hypercholesterolaemia ensued with, however, a reversal of the ester to free cholesterol ratio. In hepatic disease the plasma cholesterol changes were dependent on the flow of bile to the intestine. The changes in the amount of plasma cholesterol in ester form were presumably related to the effect of biliary stasis on the absorption of fat and cholesterol from the intestine.

Reverting to the investigation of faeces, Gardner and Gainsborough analyzed the intestinal contents obtained from a patient with a terminal ileostomy and found that, while there was practically no coprosterol present, the quantity of β -cholestanol was very much higher than in colonic faeces.

During the latter part of this period of work Gardner was handicapped by illness. Although diabetes was well controlled, other affections reduced his capacity, and much other work on faecal sterols, the sterols of fungi, and the role of deoxycholic acid was never completed.

Gardner's interests were wide and varied. While at St George's Hospital, his clinical examinations extended over a period of more than 50 years. He wrote and lectured on the sulphanilamide drugs. With Buckmaster he contributed a comprehensive account of 'Blood' to Allen's *Commercial Organic Analysis*, in which his love of accuracy is to be found in the careful and minute description of the methods for determining spectrographic absorption. Perhaps his only pastime was fishing in the Yorkshire rivers, and perhaps it was for this reason that Gardner was always interested in any investigation concerning the pollution of river waters. He was a member of the Scientific Advisory Committee set up in 1924 to advise the Standing Committee on River Pollution under the Ministry of Agriculture and Fisheries. His considerable experience in the accurate determinations of blood gases led him to make use of his tapless and jointless modification of

the Töpler pump for the difficult investigation of the oxygen consumption of different fresh-water fish. The relatively small gas volume relative to water volume demanded a high degree of accuracy and manipulative skill. Working with Leathes and later with Powers and King (*Biochem. J.* 1914, 1922, 1923) he found that the oxygen consumption of fish approximately doubled for a rise of 10°, and that below 10° the respiratory quotient fell appreciably, suggesting a fundamental change of metabolism. Of the different fish experimented with, trout were asphyxiated by a reduction of normal oxygen tension to 1/3, gold fish would require a reduction to 1/10, and eels still less.

Gardner was one of the founders of the Biochemical Society. At a meeting called at University College, London, in January 1911 Gardner acted as Chairman and explained the reasons for the formation of a Society which should include the many and wide interests of Biochemistry. The audience was very interested and proposed immediate action. Rules were drawn up and the first meeting was held in March 1911. At the subsequent dinner the rules were passed. After discussion it was deemed preferable to call the association The Biochemical Club, mainly for the reason that it did not possess a *Journal*. At that time the *Biochemical Journal* was owned and edited by the late Professor Benjamin Moore and Dr Whitley and published in Liverpool. Professor Moore became a member of the Biochemical Club. Negotiations were initiated for the publication of a *Journal* and ultimately it was arranged with Moore and Whitley that the *Biochemical Journal* should be purchased by the Biochemical Club. Gardner was active in these negotiations. The purchase was completed early in 1913. At the Annual Meeting in March of that year the association became the Biochemical Society. Plimmer was appointed Hon. Secretary and Gardner Hon. Treasurer. The membership of the Society was then under 200, and during the war years, 1914–18, Gardner managed the financing of the *Journal* so that the Society was never in debt. Though the work of the Treasurer increased year by year Gardner was reluctant to relinquish his duties and remained

Treasurer for over 30 years. Thanks to his careful management he saved a considerable surplus of funds which were placed in the hands of Trustees. Gardner acted throughout as convenor and secretary of the Trustees. The present firm financial foundation of the Biochemical Society is entirely due to Gardner.

In 1903, Gardner married Mrs Henrietta White, who died in 1926. This charming and highly accomplished lady was hostess at many pleasant gatherings in their Kensington house. During the 1914–18 war Mrs Gardner was appointed by the War Office as interpreter for the lesser known Nordic languages. During the last war Gardner was an active Warden. Cool under all emergency, his presence in the neighbourhood of High Street, Kensington, during all heavy raids was invaluable.

On retirement from St George's Hospital in 1942 with no abatement of his enthusiasm for research, he continued to work in the writer's laboratory at St Mary's Hospital Medical School. While so engaged he received by telephone during the V-rocket period, the message that his sister, Miss Gardner, had been injured and their flat destroyed. This seemed to forecast the cessation of Gardner's work. Fortunately, Miss Gardner was not seriously hurt. Failing to find accommodation in London, they left for Bradford, hoping to return later.

Gardner was a connoisseur of good food and good wine. He was also a connoisseur of good men, never happier than when in their company. Full of kindness towards his fellow-men, he was hurt by the absence of such in others. A beloved member of the Old Bradfordians, a Knight of the Round Table, and a member of a number of Masonic Lodges in London and at Bradford, he was the friend of all. His punctilious sense of responsibility, and no doubt his desire not to lose touch with so many friends, brought him back to London in all weathers to preside at his Lodge functions. He died at Bradford from heart failure following bronchopneumonia. Since the death of Mrs Gardner he had been under the constant care of his sister, Miss Mary Gardner, whose devotion to her brother is well known to many members of the Biochemical Society.

G. W. ELLIS