- Matsumura, K. (1927). *J. Amer. chem. Soc.* **49**, 810.
 Mellor, D. P. & Craig, D. P. (1941). *J. roy. Soc. N.S.W.* **74**, 475.
 Michaelis, L., Hill, E. S. & Schubert, M. P. (1932). *Biochem. Z.* **255**, 66.
 Mills, W. H. & Watson, W. H. (1910). *J. chem. Soc.* **97**, 741.
 Moeller, T. (1943). *Industr. Engng Chem.* (Anal. ed.), **15**, 346.
 Morgan, G. & Burstall, F. (1936). *Inorganic Chemistry*. Cambridge: W. Heffer & Sons.
 Musajo, L. & Minchilli, M. (1941). *Ber. dtsh. chem. Ges.* **74**, 1839.
 v. Niementowski, S. & Sucharda, E. (1916). *Ber. dtsh. chem. Ges.* **49**, 12.
 Nölting, E. & Trautmann, E. (1890). *Ber. dtsh. chem. Ges.* **23**, 3654.
 Pfitzinger, W. (1902). *J. prakt. Chem.* (2), **66**, 263.
 Rosenmund, K. W. & Karst, G. (1941). *Arch. Pharm., Berl.*, **279**, 154.
 Schlosser, A. & Skraup, Z. (1881). *Mh. Chem.* **2**, 530.
 Skraup, Z. (1882). *Mh. Chem.* **3**, 535.
 Wrede, F. & Strack, E. (1929). *Ber. dtsh. chem. Ges.* **62**, 2051.
 Zentmyer, G. A. (1944). *Science*, **100**, 294.

The Potentiation of the Toxicity of some Vesicants to the Pyruvate Oxidase System *in vitro* by certain Dithio-compounds

BY R. A. PETERS AND R. W. WAKELIN, *Department of Biochemistry, Oxford*

(Received 23 September 1946)

One of the more puzzling features of the *in vivo* action of mustard gas has always been the small amount which can prove toxic to animals. In this laboratory, for instance, it was reported (Holiday, Ogston, Peters, Philpot & Stocken, 1940) that 2 mg. of mustard gas placed upon the skin of most rats weighing 200 g. would kill. Allowing for loss by evaporation, hydrolysis and fixation in the skin, this means that the toxicity is < 10 µg./g. of rat, and is of a different order from that associated with the *in vitro* action of mustard gas upon certain enzymes and enzyme systems which is c. 300 µg./ml. (Peters & Wakelin, 1940; Dixon & Needham, 1941a; van Heyningen, 1941); it is also quite different from the toxicity of arsenicals and of divinyl sulphone and 2:2'-dichlorodiethyl sulphone, which approaches *in vitro* more closely the concentration of *in vivo* toxicity. There is much evidence now of causal connexion between lewisite action and enzyme poisoning (see review by Peters, Stocken & Thompson, 1945). It would be satisfactory also to interpret the activity of mustard gas in terms of an enzyme poisoning; Dixon, van Heyningen & Needham (1942) have indeed developed the theory that mustard gas and other vesicants owe their action to toxic effects upon hexokinase. In the course of this work, in which the pyruvate oxidase system was used as a test for possible thiol-containing therapeutic agents (Holiday *et al.* 1940), it was found that *N:N*-diethyldithiocarbamate or *N:N*-di-(2-hydroxyethyl)-dithiocarbamate had a special action. Instead of protecting the pyruvate oxidase system against mustard gas it much increased its toxicity; this effect was called 'potentiation'. The important point was made that in the presence of some dithio compounds, mustard gas becomes almost 20 times more toxic, and some evidence was produced that the action was due to

a compound of the dithio substance and mustard gas. While there is no reason to think that the potentiating action of these dithio compounds occurs *in vivo*, it was felt that a closer study was advisable. We were reinforced in our suspicion that there might be unsuspected ways in which the action of mustard gas was altered *in vivo* by our finding that mustard gas was often toxic to the succinic dehydrogenase enzyme in a brain *brei*, whereas it was not so in the same enzyme in ground-muscle preparations (Peters & Wakelin, 1941). The opposite is true of phosphocreatine phosphokinase, which is sensitive to mustard gas in the isolated state (Cori, Colowick, Berger & Slein, 1945), whereas it has not been found to be poisoned by treating skin with mustard gas *in vivo* (Needham, quoted by Dixon, 1943).

Following our initial observations, Dixon & Needham (1941b), using the pyruvate oxidase of *E. coli*, confirmed the effect of diethyldithiocarbamate, and the toxicity of the incubated product; they added the further observation that addition of diethyldithiocarbamate (not itself toxic), after hydrolysis of mustard gas was complete, could still cause an inhibition. Hence they thought that the potentiations could not be explained simply by the formation of a more toxic addition compound.

In this report, modified from the original (Peters & Wakelin, 1943b), we present a closer study of the potentiating effect upon the pyruvate oxidase system of diethyldithiocarbamate and diethanol-dithiocarbamate; no essential difference in the action of the two compounds has been found. The toxic effects are due to the formation of a compound of mustard gas with the diethyldithiocarbamate ion, which is more toxic than the sum of the toxicities of the two compounds.

EXPERIMENTAL

Brei from pigeon brain has been used as described previously (Peters & Wakelin, 1946). We have determined the amount of the compound giving a 50% reduction in O_2 uptake in presence of pyruvate over the respiration period 30–120 min. from the start of the readings. When diethyldithiocarbamate (*D*) or diethanoldithiocarbamate (*Dol*) was used, this was put into the respiration bottles before the poison. Both *D* and *Dol* have some toxicity; with *brei*, we used 1 mg./ml. and with dispersions, 0.5 mg./ml. Where dialyzed dispersions were employed, the dialysis was carried out for 4–5 hr. at 2° against 0.4% KCl. Q_{O_2} values have been calculated assuming that pigeon brain contains 80% water. In the case of *brei*, during the respiration period of 30–120 min., the residual respiration becomes much reduced and is therefore relatively less important as the experiment proceeds; further, since the presence of pyruvate inhibits lactate oxidation (Green & Brosteaux, 1936), there will be less residual respiration in the bottles containing pyruvate; any traces of pyruvate formed from residual lactate will not increase the O_2 uptake as the system is already saturated with pyruvate. In our earlier experiments we always set up controls for the residual respiration; but based upon the considerations above and upon experimental observations, we dispensed with these in this series and changes from the results with pyruvate alone were calculated. Much gain in time and saving of animals result when pyruvate is added to all twelve of the bottles, as one control and five duplicate observations are possible on one brain. Any toxicity of the dithiocarbamate has been allowed for by subtracting this from the total toxicity figures.

The procedure originally reported (Holiday *et al.* 1940) for obtaining maximal increases in toxicity has been used. Diethyl- (or diethanol-) dithiocarbamate (1 mg./3 ml. of Ringer-phosphate) was added after division of the *brei* or after adding the dispersion to the bottles; after shaking for 1–2 min., the mustard gas emulsion in lecithin was then added. Under these conditions the extra toxicity will be observed within 10–15 min. and will be at a maximum after 30 min. at 38°. The reasons for using lecithin and the arguments indicating that it has no adverse influence on the results are given fully elsewhere (Peters & Wakelin, 1941).

Mono- and bis-compounds of mustard gas and diethyldithiocarbamate were tested. The preparation of the mono-compound (I) 2-chloroethyl 2-(*N*-diethyldithiocarbamato)-ethyl sulphide and bis-compound (II) bis-[2-(*N*-diethyldithiocarbamato)-ethyl] sulphide is described elsewhere (Peters & Wakelin, 1947).

RESULTS

Potentiation

Table 1 shows the potentiating effect of *Dol* upon the toxicity brought about by mustard gas and some other compounds. It will be seen that there is

Table 1. *The effect of diethanoldithiocarbamate (Dol) upon the concentrations of poisons required to produce 50% inhibition of O_2 uptake in the pyruvate oxidase system*

(Pigeon brain *brei*; substrate, Na pyruvate, 0.018 M; results the average for respiration period 30–120 min., 38°.)

		With <i>Dol</i> (μ M)	No <i>Dol</i> (μ M)
Arsenical	Lewisite	16	16
Non-arsenical	Iodoacetamide	(60)*	60
	Divinyl sulphone	(67)†	67
	Mustard gas	(63)‡	>1000
	2:2'-Dichlorodiethyl sulphone	110§	110
	2:2':2"-Trichlorotriethylamine	130	750
	Allyl isothiocyanate	195¶	242¶

* One determination gave for 18 μ M, –24% change in respiration.

† One determination gave for 33.5 μ M, –28% change in respiration.

‡ Based upon 14 determinations (10 different brains) at different concentrations of mustard gas.

§ One determination.

|| Three determinations.

¶ Three determinations; these values are approximate but they indicate the order of toxicity. The difference of results with and without *Dol* is not significant.

(Results were calculated on total oxidation with pyruvate as substrate, without subtraction of the residual respiration.)

approximately a twenty-fold increase in the toxicity of mustard gas, and smaller increases also occur with other compounds. In Table 2, the result of addition of *Dol* upon several compounds having reactive Cl atoms as in mustard gas itself are compared at a concentration of 200 μ g./3 ml. At this concentration the effect of mustard gas added alone is very slight. It will be noticed that the effect is larger with the compounds containing the two reactive Cl atoms.

Conditions for potentiation. Table 3 gives a list of all substances tested and shows that only substances containing the dithio grouping have this action; it was not found with all of these. The list in Table 3 shows that potentiation was confined to substances which were capable of alkylation reactions similar to mustard gas. The absence of potentiation with the arsenical tested, or with the sulphones, is interesting, and also the fact that, in the presence of *Dol*, the toxicity of mustard gas to the pyruvate oxidase system approaches that of divinyl sulphone.

Nature of the potentiation. This has been investigated in the case of mustard gas only and diethyldithiocarbamate (*D*). The toxicity is due to some compound other than mustard gas, for the following reason. If an emulsion of mustard gas (c. 1 mM) be shaken in Ringer-phosphate for approx. 45 min. at 38°, it is then practically non-toxic, owing to hydrolysis to thiodiglycol and HCl; but an emulsion of

Table 2. *The effect of diethyldithiocarbamate (D) upon the toxicity of sulphides containing Cl as in mustard gas compared at concentrations of 200 µg./3 ml.*

(Substrate Na pyruvate.)				
Compound	Remarks	Change in respiration (%)		Change due to <i>D</i> (%)
		No <i>D</i>	With <i>D</i>	
2-Chloroethyl ethyl sulphide	One Cl	-28	-38	-10*
2-Chloroethyl propyl sulphide	One Cl	-5	-42	-37†
2-Chloroethyl phenyl sulphide	One Cl	-10.5	-40.5	-30‡
2:2'-Dichlorodiethyl sulphide	Two Cl	-13	-80	-67
Di-(2-chloroethylthioethyl) ether	Two Cl	-(0.7)	-(52-56)	-(45-56)§
1:1'-Dichlorodiethyl sulphide	Two Cl	-17	-7	+10
3:3'-Dichlorodipropyl sulphide	Two Cl	-23	-38.4	-15

* 78 µg. alone gave -2% change in respiration.
† 86 µg. alone gave -16%.
‡ 108 µg. gave nil to 11.5%; +*D* -22%.
§ 50 µg. (63 µM) alone gave -23% and -36.5%; 37 µg. gave -35%.

Table 3. *Substances which potentiate or do not change the effect of mustard gas, when added together with this to a brei or dispersions of pigeon brain*

Potentiating	No change
Diethyldithiocarbamate	2:3-Dimercaptopropanol (BAL)
Diethanoldithiocarbamate	Thiosalicylate
Diethanedithiophosphonate	p-Thiol benzoate
Butanedithiophosphonate	Monothiophosphonates
	(Na salts)
	Dithiophosphonates
	(Na salts)
	Na ethyl monothiocarbamate
	Thiomalic ester
	Allyl isothiocyanate
	Na thiosulphate

Both improve respiration: Methane, ethane, n-butane, toluene, diethane, n-hexane

Table 4. *Substances potentiated by diethyl- (or di-ethanol-) dithiocarbamate*

Effect	Potentiated
Large	Mustard gas
	Di-(2-chloroethylthioethyl) ether
Intermediate	2:2':2"-Trichlorotriethylamine
Small	2-Chloroethyl ethyl sulphide
	2-Chloroethyl propyl sulphide
	2-Chloroethyl phenyl sulphide
	2-Chloro-2'-hydroxydiethyl sulphide
Slight	3:3'Dichlorodipropyl sulphide

mustard gas added to D in Ringer-phosphate is still toxic after 2 hr. and will retain this toxicity after storage in the cold. Exp. 2373 (Table 5) shows this; exp. 2374 shows that the compound is much more toxic than mustard gas itself, and that emulsification is necessary for full development of the toxicity. The toxic action develops quickly, and the greatest toxicity is obtained by warming the lecithin mixture of the compound and centrifuging in Ringer-phosphate at 38°.

By shaking mustard gas in an aqueous solution of D under conditions simulating those of the potentiation experiment, a mono-substitution product (I) is formed, which has been isolated as crystalline needles m.p. 40.5° (Peters & Wakelin, 1943a,

1947). The compound is soluble in c. 60% ethanol, and by kinetic methods (Peters & Walker, 1923) the rate of acid liberation is the same as that of mustard gas in the same solvent.

Table 6 shows the action of the highly purified compound (I) when compared with an amount of D equivalent to that contained in the concentration of (I) tested.

Table 7 shows that compound (I) does not further potentiate mustard gas and in its presence the latter shows no added activity; also that the bis-compound (II) of mustard gas and 2 molecules of diethyldithiocarbamate is not toxic under these conditions with a brei, and that mustard gas exerts its usual toxic effect in its presence.

Causes of potentiation

Since the mono-compound (I) is toxic, and the bis-compound not so, and since (I) still contains a Cl capable of substitution, it was felt first that potentiation must be due to the formation of the tissue compound (III) from (I).

This view was not entirely tenable because (a) the crude but Cl-free OH compound (IV) found by hydrolysis of (I) was still toxic and (b) the mono-Cl analogues of mustard gas showed potentiation,

Table 5. *Toxicity of compounds (not isolated) formed by shaking mustard gas with diethyldithiocarbamate (D), and effects of emulsifying with lecithin*(Pigeon *brei*; substrate, Na pyruvate.)

Exp. 2373. Toxicity of mustard gas (lecithin emulsion) after shaking with diethyldithiocarbamate for 70 min. at 38° in respirometer bottles.

Addition	Q_{O_2}	
	30-120 min.	Decrease (%)
Equivalent of <i>D</i> alone (150 μ g.)	10.72	—
Product of <i>D</i> + 200 μ g. mustard gas	6.41	40

Exp. 2374. Toxicity of mixture of *D* + mustard gas (*c.* equimolar) when presented to the brain tissue as a suspension or as an emulsion in lecithin. (Lec. = lecithin. Na pyruvate in all bottles and diethyldithiocarbamate added in equivalent amounts as control where necessary. The mixture of Lec. + *D*, Lec. + *D* + mustard gas, *D* + mustard gas, and *D* alone were shaken respectively for 1 hr. at 38° in Ringer-phosphate solution. Samples (0.4 ml. $\equiv$ 200 μ g. mustard gas) were then added to the experimental bottles either suspended or re-emulsified with lecithin.)

Addition	Q_{O_2}		
	(0-30 min.)	(30-120 min.)	Change (%) (30-120 min.)
Suspended { <i>D</i> alone	14.55	10.57	—
Product (<i>D</i> + 200 μ g. mustard gas)	12.70	9.90	- 6
Product (<i>D</i> + 800 μ g. mustard gas)	12.50	9.13	- 14
Emulsified { Lec. + <i>D</i>	13.42	9.67	—
Lec. + product (<i>D</i> + 200 μ g. mustard gas)	8.24	5.94	- 38.5

Table 6. *Comparison of toxicity of crystalline compound (I) of mustard gas and diethyldithiocarbamate with an amount of the latter equivalent to that combined in the amount of (I) used for test*(Brain *brei*. Na pyruvate, 0.018M.)

Addition	Q_{O_2} (30-120 min.)
Lecithin only	10.22
Lecithin + <i>D</i> (191 μ g.)	9.37
Lecithin + (I)* (300 μ g.)	5.58

* (I) emulsified with lecithin at 38° by mixing and stirring into warmed Ringer-phosphate solution.

though to a much less extent than the bis-Cl compound.

We think that the non-toxicity of compound (IV) is largely due to its failure to reach the active centre; even when emulsified with lecithin, it is largely present as crystals at 38° and remains suspended. This view is supported by other work, because very toxic effects have been obtained (Table 8) with a non-crystalline compound of diethanoldithiocar-

Table 7. *Comparison of toxicity of mono- and bis-compounds (I) and (II) in presence and absence of added mustard gas (MG)*(Brain *brei*. Na pyruvate, 0.018M.)

Addition	Q_{O_2}		
	(0-30 min.)	(30-120 min.)	Change (%) (30-120 min.)
Exp. 2382. Lec. only	14.0	10.03	—
Lec. + 0.6 mg. (I)	9.01	3.94	- 61
Lec. + 0.6 mg. (I) + 0.33 mg. MG	9.06	3.72	- 63
Lec. + 2 mg. (II)	14.13	10.51	+ 5
Lec. + 2 mg. (II) + 0.33 mg. MG	12.22	7.59	- 27
Exp. 2383. Lec. only	13.02	10.93	—
Lec. + 0.6 mg. (I)	7.85	3.55	- 67
Lec. + 0.6 mg. (I) + MG	8.39	3.19	- 71
Lec. + 1 mg. (II)	13.02	10.38	- 5
Lec. + 1 mg. (II) + MG	12.71	7.89	- 24

Lec. = lecithin.

The Potentiation of the Toxicity of some Vesicants to the Pyruvate Oxidase System
in vitro by certain Dithio-compounds

By R. A. Peters and R. W. Wakelin

Volume 41 (1947), No. 4, p. 548, col. 2, line 5

for of compound (IV) *read* of compound (II)

bamate and mustard gas free from Cl, which, though impure appeared to be mainly the bis-Dol compound, and which was much more soluble than the diethyl compound.

Table 8. *Toxicity of bis-product of Dol + mustard gas*
(Brain brei. Na pyruvate, 0.018 M.)

Addition	Q_{O_2} (30–120 min.)	Change (%)
Lecithin only	9.80	—
Lecithin + 300 μ g. (Dol only)	8.85	-10
Lecithin + 300 μ g. (Dol + MG)	2.61	-73

Hence the conclusion is reached that it is the $\text{=N.C}\begin{smallmatrix} \text{S} \\ \diagup \diagdown \end{smallmatrix}$ grouping in compound (I) which is inherently toxic; this would be in agreement with the lesser toxicity of dithiocarbamates themselves and of allyl isothiocyanate. Substitution in the SH thus increases the reactivity of the thio (=S) group.

In the potentiation experiment, the effect will be due to a combination of (a) the toxicity of compound (I) or its OH derivative and (b) formation of tissue compounds as (III) by substitution of Cl either directly or through compound (I) as intermediary. The latter provides a possible explanation of Dixon & Needham's (1941b) experiments; in their case, the initial treatment with mustard gas may have left unhydrolyzed Cl half-products such as (V) available for combination with added D. We have shown already that such products are present in mustard gas-kerateine (Peters & Wakelin, 1941).

Compound (V):

In reaching the above conclusions, we have discarded some other possibilities, which may be mentioned briefly.

Discarded possibilities. We have produced evidence for the presence of SH groups in brain pyruvate dehydrogenase (Peters & Wakelin, 1946; circulated privately in 1941) from experiments with S—S compounds, and by showing that sodium maleate is inhibitory in very low concentration. It might be thought that potentiation was produced by blocking some of the SH groups necessary for pyruvate dehydrogenase action, and that this might render the whole system more sensitive. In the experiments of which one is quoted (Table 9) with various concentrations of maleate no potentiation was observed. This does not suggest that potentiation is due to partial blocking of an SH group.

Again, it was thought that the SH groups might be partly oxidized to S—S by oxidized diethyldithiocarbamate, as Keilin & Hartree (1940) have demonstrated an inhibition of this nature with succinic dehydrogenase. In a test, however, we could obtain no inhibition in this way. Again, potentiation cannot be reversed either with glutathione or with kerateine (the latter is known to stabilize respiration, see Peters & Wakelin (1941)).

Table 9. *Effects of Na maleate and mustard gas upon Q_{O_2} of brain brei with pyruvate*

Addition/3 ml.	Decrease in Q_{O_2} (%)	Total decrease (%)
200 μ g. mustard gas	- 8	- 46
400 μ g. Na maleate	- 38	
200 μ g. mustard gas + 400 μ g. Na maleate	- 46.5	- 46.5

Toxicity to animals

When impure, compound (I) is toxic to the rat, both as applied on the skin and by injection, and kills with the signs of mustard gas poisoning, except that there is a tendency for ulceration of the skin, which is unusual for mustard gas. Dr Holiday tells us that he once observed this also with a crude mixture. It is undoubtedly due to residual traces of mustard gas or its half-hydrolysis product dissolved in the oil. We have found these crude oils very irritant to mustard gas-sensitive men. When pure we have injected comparatively large doses, emulsified in lecithin, intraperitoneally into several rats and one guinea-pig. In two cases, the rats died after 1 and 2 months respectively, but we hesitate to regard this as significant. If placed in the eye of a rabbit, no ill effects have followed (Mann & Pirie, 1943).

Since the compound is toxic to the pyruvate system *in vitro*, it is evident that its physical properties prevent penetration to the active centres *in vivo*, probably owing to predominant lipid solubility. A similar conclusion was reached for the difference in toxicity between some acriflavines (Manifold, 1941).

DISCUSSION

The much greater potentiation shown with the di-Cl compounds is the first distinction between mono-Cl and di-Cl consistent with competition reactions (Holiday, Ogston, Philpot & Stocken, 1940), for which there is the explanation that the greater activity may be due to the possibility that more than one toxic compound can be formed with the di-Cl compound. In regard to the action of the substituted dithiocarbamate compound, one possible view is that it diverts an attack upon some other grouping into one upon SH groups in the pyruvate oxidase system.

So far there has been little exploration of the enzyme field. We have found that inhibition of the succinoxidase system by mustard gas, when it occurs, is not potentiated by D. As the cytochrome system is common to both this and the pyruvate system, this means that it is not the cytochrome component of the pyruvate system which is attacked. The potentiation can be observed also with brain dispersions.

SUMMARY

1. Some dithiocarbamates and dithiophosphonates potentiate the toxic action of mustard gas upon the pyruvate oxidase system; the effect of diethyl- (or diethanol-) dithiocarbamate has been closely examined.

2. Only compounds having a Cl atom which behaves kinetically as in mustard gas are potentiated; the effect is quickly established.

3. Potentiation is due to the formation of the half product of mustard gas with diethyldithiocarbamate (compound (I), 2-chloroethyl-2-(*N*-diethyldithiocarbamate)-ethyl sulphide), prepared crystalline under the conditions of the experiment. This is much more toxic than either of its components.

4. The toxicity of (I) is not due to the Cl group; the hypothesis is advanced that the $=N.C \begin{smallmatrix} S \\ \diagup \\ S- \end{smallmatrix}$ group is inherently toxic.

5. The bis-substitution product was only slightly toxic, which is explained by its inability to reach the active enzyme centre in the brain *brei*.

6. (I) is not toxic when injected into rats. This is interpreted as due to failure to reach the pyruvate system *in vivo*.

This work formed part of a programme of research carried out for the Ministry of Supply and reported in 1943; we are indebted to the Chief Scientific Officer, Ministry of Supply, for permission to publish. We are also grateful to other members of the team for help; and to Mr Philpot and Dr Stocken for thiophosphonates.

REFERENCES

- Cori, F. C., Colowick, S. P., Berger, L. & Slein, M. W. (1945). By communication.*
- Dixon, M. (1943). Report to Ministry of Supply by Dixon, no. 19.*
- Dixon, M. & Needham, D. M. (1941*a*). Report to Ministry of Supply by Dixon, no. 1.*
- Dixon, M. & Needham, D. M. (1941*b*). Report to Ministry of Supply by Dixon, no. 2.*
- Dixon, M., van Heyningen, R. & Needham, D. M. (1942). Report to Ministry of Supply by Dixon, no. 11.*
- Green, D. E. & Brosteaux, J. (1936). *Biochem. J.* **30**, 1489.
- Holiday, E., Ogston, A. G., Peters, R. A., Philpot, J. St L. & Stocken, L. A. (1940). Report to Ministry of Supply by Peters, no. 15.*
- Holiday, E., Ogston, A. G., Philpot, J. St L. & Stocken, L. A. (1940). Report to Ministry of Supply by Peters, no. 1.*
- Keilin, D. & Hartree, H. (1940). *Proc. Roy. Soc. B*, **129**, 277.
- Manifold, M. C. (1941). *Brit. J. exp. Path.* **22**, 111.
- Mann, I. & Pirie, A. (1943). Personal communication.
- Peters, R. A., Stocken, L. A. & Thompson, R. H. S. (1945). *Nature, Lond.*, **156**, 616.
- Peters, R. A. & Wakelin, R. W. (1940). Reports to Ministry of Supply by Peters, nos. 2 and 8* (and in the Press).
- Peters, R. A. & Wakelin, R. W. (1941). Report to Ministry of Supply by Peters, no. 39.*
- Peters, R. A. & Wakelin, R. W. (1943*a*). Report to Ministry of Supply by Peters, no. 71.*
- Peters, R. A. & Wakelin, R. W. (1943*b*). Report to Ministry of Supply by Peters, no. 72.*
- Peters, R. A. & Wakelin, R. W. (1946). *Biochem. J.* **40**, 513.
- Peters, R. A. & Wakelin, R. W. (1947). *J. chem. Soc.* p. 849.
- Peters, R. A. & Walker, E. (1923). *Biochem. J.* **17**, 260.
- van Heyningen, R. (1941). Report to Ministry of Supply by Dixon, no. 3.*

* Information available on application to Ministry of Supply, London.

Observations upon a Compound of Mustard Gas and Kerateine

By R. A. PETERS AND R. W. WAKELIN, *Department of Biochemistry, Oxford*

(Received 20 December 1946)

The fact that mustard gas will combine with some proteins is known from the work of Berenblum & Wormall (1939) on serum proteins, Berenblum (1940) on nucleoproteins, Pirie (1941) on collagen, and may be inferred also from the work of Fell & Allsopp (1940-1). Since enzymes are proteins, such combination is indicated from enzyme studies in Oxford (Peters & Wakelin, 1940) and Cambridge (see summary by Dixon & Needham, 1946), as also from kinetic considerations of the reactions of

mustard gas (Holiday, Ogston, Philpot & Stocken, 1940; Ogston, 1941).

Earlier work in this laboratory (Peters & Walker, 1924-5) had shown that mustard gas did not abolish the fixed SH reaction (Hopkins & Dixon, 1922) of extracted muscle tissue though dichlorodiethyl sulphone and phenyldichloroarsine readily did so. This is rather puzzling because mustard gas so readily combines with simple monothiols (Holiday *et al.* 1940; van Heyningen, 1941). The original object of