- Ennor, A. H. (1942). *Aust. J. exp. Biol. med. Sci.* **20**, 73.
- Ennor, A. H. & Stocken, L. A. (1947). *Nature, Lond.*, **159**, 308.
- Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
- Fiske, C. H. & Subbarow, Y. (1929). *J. biol. Chem.* **81**, 629.
- Flock, E., Bollman, J. L. & Mann, F. C. (1936a). *J. biol. Chem.* **115**, 179.
- Flock, E., Bollman, J. L. & Mann, F. C. (1936b). *J. biol. Chem.* **115**, 201.
- Higgins, G. M., Berkson, J. & Flock, E. (1932). *Amer. J. Physiol.* **102**, 673.
- Higgins, G. M., Berkson, J. & Flock, E. (1933). *Amer. J. Physiol.* **105**, 177.
- Kalckar, H. M. (1945). *Ann. Rev. Biochem.* **14**, 283.
- Kaplan, N. O. & Greenberg, D. M. (1943). *J. biol. Chem.* **150**, 479.
- Kaplan, N. O. & Greenberg, D. M. (1944a). *J. biol. Chem.* **156**, 525.
- Kaplan, N. O. & Greenberg, D. M. (1944b). *J. biol. Chem.* **156**, 543.
- Kaplan, N. O. & Greenberg, D. M. (1944c). *J. biol. Chem.* **156**, 511.
- Kosterlitz, H. W. & Ritchie, C. M. (1943). *Biochem. J.* **37**, 181.
- Lehninger, A. L. (1944). *J. biol. Chem.* **154**, 309.
- Lehninger, A. L. (1945a). *J. biol. Chem.* **161**, 437.
- Lehninger, A. L. (1945b). *J. biol. Chem.* **157**, 363.
- Le Page, G. A. (1946). *Amer. J. Physiol.* **146**, 267.
- Le Page, G. A. & Umbreit, W. W. (1945). In *Manometric Techniques and Related Methods for the Study of Tissue Metabolism*, by W. W. Umbreit, R. H. Burris & J. F. Stauffer. Minneapolis: Burgess Publishing Co.
- Lipmann, F. (1941). In *Advances in Enzymology and Related Subjects*, **1**, 99, ed. by Nord, F. F. & Werkman, C. H. New York: Interscience Publishers Inc.
- Lipmann, F. & Tuttle, L. C. (1944). *J. biol. Chem.* **153**, 571.
- Lipmann, F. & Tuttle, L. C. (1945). *J. biol. Chem.* **159**, 21.
- Markham, R. (1942). *Biochem. J.* **36**, 790.
- Meier, R. & Thoenes, E. (1933). *Arch. exp. Path. Pharmac.* **169**, 655.
- Meijbaum, W. (1939). *Hoppe-Seyl. Z.* **258**, 117.
- Munoz, J. M. & Leloir, L. F. (1943). *J. biol. Chem.* **147**, 355.
- Nelson, N., Rapoport, S., Guest, G. M. & Mirsky, I. A. (1942). *J. biol. Chem.* **144**, 291.
- Potter, V. R. & Elvehjem, C. A. (1936). *J. biol. Chem.* **114**, 495.
- Quastel, J. H. & Wheatley, A. H. M. (1935). *Biochem. J.* **29**, 2773.
- Rapoport, S. (1945). *J. biol. Chem.* **161**, 429.
- Rapoport, S., Leva, E. & Guest, G. M. (1943a). *J. biol. Chem.* **149**, 57.
- Rapoport, S., Leva, E. & Guest, G. M. (1943b). *J. biol. Chem.* **149**, 65.
- Rapoport, S. & Nelson, N. (1945). *J. biol. Chem.* **161**, 421.
- Stone, W. E. (1943). *J. biol. Chem.* **149**, 29.
- Wieland, H. & Rosenthal, C. (1943). *Liebigs Ann.* **554**, 241.

The Estimation of Creatine

By A. H. ENNOR (Wellcome Research Fellow) AND L. A. STOCKEN
Department of Biochemistry, University of Oxford

(Received 1 October 1947)

During the course of an investigation into the metabolism of fatty livers (Ennor & Stocken, 1948) it became desirable to identify, and subsequently to estimate, amounts of a very labile phosphorus compound present in the livers of normal guinea pigs and in others treated with carbon tetrachloride. Although phosphate liberated from organic phosphate compounds in acid molybdate is often considered to arise from phosphocreatine, there are other possibilities, and we endeavoured therefore to correlate the labile phosphorus with the creatine liberated following mild hydrolysis. It was decided to estimate the creatine as such rather than after conversion to creatinine, and an attempt was made to employ the method of Eggleton, Elsdon & Gough (1943). These authors have pointed out that trichloroacetic acid (TCA) extracts of liver contain inhibitory substances which prevent the estimation of creatine by the Barritt (1936) modification of the Vosges & Proskauer (1898) reaction. The suggestion

was also made that these inhibitors were the ω -amino aliphatic acids.

In the fractionation of TCA extracts of guinea-pig liver by means of barium (Le Page & Umbreit, 1945) it was found that both the inhibitor and phosphocreatine were precipitated in the ethanol-insoluble fraction. An examination of this fraction indicated that the inhibitory substance or substances was more likely to be glutathione than an amino-acid as suggested by Eggleton *et al.* (1943). Such a supposition, because of the wide distribution of the tripeptide, necessarily casts some doubt upon the validity of the application of the α -naphthol-diacyetyl reaction to the estimation of creatine in tissues.

This communication is concerned with some experiments on the inhibition of the Barritt reaction by sulphhydryl compounds and suggests a modification of the procedure of Eggleton *et al.* by means of which a more accurate determination of the creatine content of tissues may be made.

EXPERIMENTAL

Reagents

Creatine hydrate. An A.R. specimen was recrystallized according to Hunter (1928) and its purity checked by N analysis.

Diacetyl. A stock solution of approx. 1% was prepared from A.R. dimethylglyoxime as described by Walpole (1911). This was diluted and used as described by Eggleton *et al.* (1943). No apparent loss in activity has been observed following 3 months' storage at -1° .

α -Naphthol. A faintly coloured A.R. sample has been used throughout. It was considered unnecessary to effect further purification by steam distillation. The solution of this compound and the volume used was, in all cases, as described by Eggleton *et al.* (1943).

Glutathione (GSH). This was a commercial specimen.

Oxidized glutathione (GSSG). This was prepared by bubbling O_2 through a slightly alkaline solution of GSH of appropriate concentration until a negative nitroprusside reaction was obtained.

Benzyl mercaptan and methionine. These were gifts from the Ministry of Supply.

Cysteine ester hydrochloride. This compound was mentioned casually by Friedmann (1903) who gave no details of the synthesis. A mixture of 20 g. cystine, 60 ml. conc. HCl (36.5% w/v), 20 ml. water, 200 ml. ethanol and 20 g. tin foil were gently refluxed until all the tin had dissolved. After removal of the tin with H_2S , the solution was evaporated to dryness *in vacuo* and the residue dissolved in 150 ml. ethanol and saturated with dry HCl at 0° . Crystals of cysteine ester hydrochloride separated on standing overnight. Yield 24 g. (79%). M.p. 128° . SH by I_2 titration, 17.7%; calc. for $C_3H_{12}O_4SNCl$, 17.8%.

Cystine ester hydrochloride. This was prepared according to the method of Friedmann (1903).

p-Chloromercuribenzoic acid was synthesized according to Whitmore & Woodward (1932).

2:3-Dimercaptopropanol (BAL) was obtained as described by Stocken (1947).

Procedure

This, apart from one minor modification, has followed the method described by Eggleton *et al.* (1943). Greater reproducibility is obtained if, after adjusting the volume of the creatine- α -naphthol-diacetyl solution to 10 ml. in a narrow tube, the contents are transferred to a 6×1 in. tube for mixing and development of the colour. After 30 min. development at room temperature (18°) the intensity of the colour was measured in 1 cm. cells in the Hilger Spekker absorptiometer with Ilford filter no. 604 (spectrum green).

Influence of monothiols on the Barritt reaction

Cysteine ester hydrochloride. For preliminary tests and because of the small amount of GSH available, the initial experiments were carried out with cysteine ester hydrochloride. Varying amounts of this compound were added to tubes each containing

1.0 ml. of $5 \times 10^{-4}M$ -creatine ($65.5 \mu g.$). The percentage inhibition of colour development was then calculated and plotted against the molar ratio, cysteine ester : creatine.

The influence of cysteine ester on the colour produced by creatine, diacetyl and α -naphthol is very marked, 60% inhibition being observed with a molar ratio of cysteine ester : creatine of 0.8 : 1.

Fig. 1. Inhibition of the Barritt reaction by varying amounts of cysteine ester hydrochloride and glutathione. $65.5 \mu g.$ of creatine throughout. To all tubes were added 2.0 ml. α -naphthol solution (1.0 g. α -naphthol in 100 ml. of a solution containing 6.0 g. NaOH + 16 g. Na_2CO_3) + 1.0 ml. of a 1 in 20 dilution of stock diacetyl + varying amounts of thiol. Adjusted to 10 ml. with water and colour measured after 30 min. at room temperature. □ denotes cysteine ester hydrochloride; △ denotes glutathione.

Glutathione (GSH). To test the inhibitory effect of GSH three points on the cysteine ester curve were chosen and the colour produced by creatine in the presence of a GSH : creatine ratio of 0.5, 0.8 and 1.2 mol. to 1 was determined. The points marked △ in Fig. 1 were obtained in this manner and are sufficiently close to the cysteine-ester inhibition curve to suggest that it is primarily the thiol group which is responsible for the inhibition of the Barritt reaction.

Benzyl mercaptan and 2:3-dimercaptopropanol. Two simple thiols were used in an attempt to show that the thiol group is the determining factor in the inhibition of the colour.

Table 1. Inhibition of Barritt reaction by simple thiols

Molar ratio thiol : creatine	Inhibition (%)	
	Benzyl mercaptan	BAL
0.5	47	11.5
0.8	66	17.5
1.2	84.5	24.0
2.0	—	46.5

The results (Table 1) obtained with benzyl mercaptan are similar to those found with cysteine ester and GSH and support the idea that the inhibition seen with the latter compounds is due primarily to the reactivity of the thiol group and not to the amino group. Further confirmation of this is given in the section dealing with the neutralization of the inhibitor. It will be noticed that the degree of inhibition given by BAL is very much less than that produced by the other compounds. It is likely that the inhibition is due to addition of the thiol group across some reactive double bond formed in the sequence of reactions leading to the coloured complex. If this is the case, it is not to be expected that BAL would be as efficient an inhibitor as benzyl mercaptan even if it be supposed that only one of the thiol groups in BAL is capable of such an addition.

It is clear from the foregoing experiments that thiol-containing compounds present in tissues will produce considerable interference with the estimation of creatine, and that the magnitude of the error introduced will depend on the ratio of thiol to creatine. Thus, in those tissues where the creatine content is low and the thiol content high, it would be anticipated that the method discussed would not even detect creatine.

Effect of other sulphur-containing compounds

It was considered unlikely that the oxidation of the SH group would eliminate the inhibition because of the well known tendency towards cleavage of the disulphide linkage in alkaline solution. Experiments with GSSG and cystine ester (Table 2) confirm this, although it will be noticed that GSSG produces a much greater degree of inhibition, and is apparently, therefore, much more susceptible to hydrolysis than is cystine ester.

Table 2. *Disulphide inhibition of the Barritt reaction*

Molar ratio disulphide : creatine	Inhibition (%)	
	Cystine ester	GSSG
0.25	—	22.0
0.4	—	34.0
0.6	—	50.0
1.0	3.4	—
2.0	5.0	81.0
4.0	9.1	—
8.0	18.1	—
16.0	31.9	—

It is clear from these experiments that not only thiols, but any compound which generates a thiol on treatment with alkali, will seriously interfere with the estimation of creatine in tissue extracts. It is to be regretted that the only other sulphur compound of biological importance which we have had the opportunity of testing in this reaction was methionine. This produced an inhibition of about 5% when present in a molar ratio of 4 methionine : 1 creatine.

Protection of the Barritt reaction from thiol inhibition

It is apparent that when the molar ratio of SH to creatine exceeds 0.2–0.4 (7–15% inhibition), the successful application of the Barritt reaction will depend on the removal of SH groups. Compounds known to combine with thiol groups include iodoacetic acid, maleate, trivalent arsenicals and organic mercurials. Of these we preferred the mercurials because of the high rate of formation and stability of the resultant *R*-Hg-SR' compound in alkaline solution. *p*-Chloromercuribenzoic acid was chosen as a possible protective agent in order to ensure non-precipitation of the Hg derivative.

Procedure. 1.0 ml. of a 5×10^{-4} M-solution of creatine (65.5 μ g.) was added to a series of tubes, followed by 1.0 ml. of 4×10^{-3} M-cysteine ester. This gave an initial ratio of cysteine ester : creatine of 8 : 1 (a ratio of 7 : 1 will produce 100% inhibition under these conditions, see Fig. 1). Varying volumes of an appropriate dilution of a 10^{-2} M-solution of *p*-chloromercuribenzoic acid (CMBA) were then added to produce CMBA : cysteine ester ratios from 0 to 2.0. α -Naphthol and diacetyl were then added as before and the resultant colours estimated after 30 min. All determinations were carried out in duplicate and the figures in Table 3 are the means of the values obtained.

Table 3. *Protection afforded by p-chloromercuribenzoic acid (CMBA) on the thiol inhibition of the Barritt reaction*

(Degree of protection = μ g. creatine found/ μ g. creatine present $\times 100$.)

Molar ratio CMBA : cysteine ester	Degree of protection
0.1	0
0.2	1.5
0.3	2.0
0.4	2.8
0.5	2.9
0.6	5.4
0.7	11.8
0.8	31.5
0.9	73.7
1.0	89.6
2.0	92.2

Increasing amounts of CMBA, as would be expected, lead to an increased removal of the inhibition produced by cysteine ester (Table 3). Complete removal of the inhibition, however, was not obtained even when the molar ratio of CMBA : cysteine ester was 2 : 1. Any of the three following factors could be responsible for this effect: (a) dissociation of the resultant *R*-Hg-SR', (b) the *R*-Hg-SR' itself, (c) the excess of the CMBA.

The degree of protection afforded by CMBA against the effects of GSH, GSSG, benzyl mercaptan and BAL was similar to that obtained in the case of cysteine ester.

Effect of ethyl α -amino- β -(*p*-carboxyphenylmercurithio)-propionate and *p*-chloromercuribenzoic acid on the Barritt reaction. The ethyl α -amino- β -(*p*-carboxyphenylmercurithio)-propionate was not isolated, but prepared in solution by mixing equivalent amounts of *p*-chloromercuribenzoic acid and cysteine ester. Convenient amounts of suitable dilutions were then added to 1.0 ml. 5×10^{-4} M-creatinine (65.5 μ g.). The subsequent procedure was as described earlier.

Table 4. Inhibition of the Barritt reaction by *p*-chloromercuribenzoic acid (CMBA) and ethyl α -amino- β -(*p*-carboxyphenylmercurithio)-propionate (R-Hg-SR')

Molar ratio compound : creatine	Inhibition (%)	
	R-Hg-SR'	CMBA
0.5	0	0
1.0	0	0
2.0	0	1.7
4.0	4.9	4.5
8.0	5.3	6.1
10.0	—	7.4
16.0	8.2	—
32.0	11.1	—

The results of this experiment (Table 4) indicate that some inhibition is to be expected when the ratio of either the complex, or of the CMBA, to creatine exceeds 2 : 1. Since, however, a ratio of 4 : 1 only produces an inhibition of 5% and a ratio of 2 mol. CMBA to 1 mol. cysteine affords 92% protection, it is clear that the error introduced is small compared with that obtained when free thiol compounds are present.

Stoichiometric relationship of the cysteine ester inhibition. It was considered worth while to determine the colour developed by varying amounts of creatine in the presence of varying amounts of cysteine ester when the relative amounts of each were maintained at a constant ratio. A molar ratio of cysteine ester to creatine of 0.6 : 1 was chosen since at this value the inhibition expected is 44.8% (Fig. 1).

The strictly linear relationship between the amount of creatine and the optical density is shown in Fig. 2. In the presence of cysteine it will be noticed that there is a distinct diminution in the inhibition at low concentrations of the thiol (Fig. 3).

If it is true that interference with the production of the coloured compound is due to a preferential reaction of some intermediate compound with the thiol then it would be expected, under the conditions of this experiment, that the percentage inhibition would remain constant. This, however, is not the case, and the explanation is probably to be found in the small absolute amount of cysteine ester which is present at the lower creatine levels. Thus at 8.18 μ g. of creatine there are only 5.6 μ g. of cysteine

ester in a volume of 10.0 ml., and it must be considered unlikely that such a small amount would remain in the reduced state. The actual ratio of cysteine ester to creatine would thus be less than 0.6 : 1. Reference to Fig. 1 shows that a reduction of this ratio to 0.4 : 1 would account for the low inhibition seen at the 8.18 μ g. level of creatine.

Fig. 2. Comparison of the colour produced by varying amounts of creatine in the absence ○ and the presence Δ of a molar ratio of cysteine ester : creatine of 0.6 : 1.

Fig. 3. Variation in the inhibition of the Barritt reaction in the presence of a constant ratio of cysteine ester : creatine of 0.6 : 1.

Application of the method to the estimation of creatine in tissue extracts

Well fed, adult, male rats of about 300 g. body weight were stunned by a blow on the head and the neck vessels severed. Samples of the various tissues were weighed in a mortar of the Potter & Elvehjem (1936) type, and homogenized with 5 ml. of 10% (w/v) trichloroacetic acid (TCA). After centrifugation, the residue was re-extracted with 10 ml. of 5% TCA. The combined supernatant fluids were made up to 25.0 ml. with water and filtered prior to the removal of samples for estimation.

Procedure. The creatine was determined in suitable samples of the *TCA* extracts both in the presence and absence of *CMBA*. To obtain some idea of the absolute amounts of reducing substances present, and hence the amount of *CMBA* necessary to combine with the SH present, samples of the *TCA* extracts were titrated with 0.01*N*-iodine after the addition of 0.5 vol. *N*-HCl (to facilitate comparison with the previous experiments, the iodine titres are expressed as equivalents of cysteine ester). If it be assumed that the reducing substances are entirely

CMBA. Thus small intestine, kidney, lung and spleen with ratios (obviously too high, cf. Fig. 1), of 2.6–7.5 : 1 show increases from 70 to 250 %, and the other tissues with ratios of 0.05–0.5 : 1 have increases of 2–16 %. In the case of liver tissue, a mean ratio of 37.2 : 1 will prevent the estimation of creatine in the absence of *CMBA* and thus the addition of this compound must lead to an infinite increase.

Table 5. *Creatine content of rat tissues found in the presence and absence of CMBA*

Tissue	Cysteine ester equivalent (mg./100 g.)	Creatine (mg./100 g.)		Molar ratio cysteine ester equivalent : creatine	Increase with <i>CMBA</i> (%)
		<i>CMBA</i> absent	<i>CMBA</i> present		
Liver (1)	185	0	4.6	28.8	∞
(2)	244	0	3.8	45.7	∞
Spleen (1)	161	5.9	15.8	7.2	197
(2)	181	3.9	15.3	8.4	292
Kidney (1)	139	7.7	20.5	4.8	166
(2)	159	9.2	22.9	4.9	150
Lung (1)	124	6.0	13.0	6.8	117
(2)	149	5.8	11.8	9.0	104
Smooth muscle (small intestine) (1)	124	24.2	40.4	2.2	67
(2)	147	20.7	34.8	3.0	68
Testis (1)	172	248.0	295.0	0.4	16.5
(2)	178	260.0	299.0	0.4	15
Brain (1)	84	114.5	130.0	0.5	14
(2)	136	115.0	133.0	0.7	15
Cardiac muscle (ventricle) (1)	110	170.0	192.0	0.4	13
(2)	147	197.0	212.0	0.5	7
Skeletal muscle (1)	46	516.0	520.0	0.06	2
(2)	39	500.0	511.0	0.05	2

thiol in character the addition of an equivalent amount of *CMBA* will ensure an excess of the reagent. The figures quoted in Table 5 are the means of at least two determinations on each sample.

It will be noticed that in the case of kidney, spleen, lung and small intestine, the addition of *CMBA* produced a considerable increase in the amount of creatine estimated (Table 5). In the case of liver, the addition of *CMBA* now renders possible the determination of the creatine present, zero values being recorded in its absence. With the remaining tissues investigated, voluntary muscle, testis, brain and cardiac muscle, the increases are not as great, a maximum mean increase of only 15.7 % being present in the case of testis.

The magnitude of the increase in the amount of creatine measurable in the presence of *CMBA* will depend on the relative amounts of thiol and creatine. The method employed to obtain the cysteine ester equivalent, and thus the ratio of this to creatine, will lead to a high result. Nevertheless it is of some interest to note that there is a fair correlation between the ratios so obtained and the percentage increase in the creatine measured in the presence of

Although the figures for the creatine content of the various tissues examined are the maximum obtainable in the presence of *CMBA*, they do not necessarily represent the true creatine values because of the possible inhibition by substances other than the *R*-Hg-SR' compound formed and the excess *CMBA* present. This possibility was examined by attempting to recover 20 µg. of creatine which were added to samples before analysis.

Appropriate samples of the *TCA* extracts were taken in quadruplicate, and to each set of tubes 2.0 ml. of 5×10^{-4} *M*-*CMBA* (1426 µg.) were added. To two of the tubes in each series 20 µg. of creatine were added. α -Naphthol and diacetyl were added as before and the colour measured after 30 min. In view of the inhibition produced by the excess of *CMBA*, the amount of creatine was obtained by reference to a standard curve prepared by measuring the colour produced by known amounts of creatine in the presence of 1426 µg. of *CMBA*.

The recoveries of 20 µg. of added creatine (Table 6) indicate that this procedure will give values for the creatine contents of tissues which are reasonably close to those actually present.

Table 6. *Recovery of 20 μ g. creatine added to trichloroacetic acid extracts of various rat tissues in the presence of CMBA*

Tissue	Creatine found (μ g.)		Recovery of added creatine (%)
	No addition	+ 20 μ g. creatine	
Muscle	28.4	48.4	100
Testis	18.0	37.4	97
Lung	13.6	31.6	90
Brain	22.2	41.0	94
Cardiac muscle (ventricle)	20.2	41.0	104
Liver	18.2	36.2	90
Spleen	13.2	30.8	88
Kidney	43.4	60.8	87
Smooth muscle (small intestine)	41.3	59.9	93

As a matter of interest the behaviour of the α -naphthol-diacetyl method for the estimation of creatine in whole blood has also been studied (Table 7).

Table 7. *Creatine content of whole blood found in the presence and absence of CMBA*

(10.91 g. of rabbit blood + 10 ml. of 10% (w/v) trichloroacetic acid, proteins separated by centrifugation. The insoluble protein residue re-extracted with a further 10 ml. of 5% trichloroacetic acid, centrifuged and the supernatant fluids combined and made up to 25.0 ml. with water. Filtered. 2.0 ml. samples taken in all cases.)

Creatine in sample (μ g.)			Increase in presence of CMBA (%)	Recovery (%)
CMBA absent	CMBA present	+ 20 μ g. creatine, CMBA present		
14.6	18.6	36.2	27.4	88

Creatine, mg./100 g. blood = 2.4 (calculated from creatine measured in sample in the presence of CMBA).

Here again the recovery of added creatine may be carried out satisfactorily and is comparable in magnitude with that observed in tissue extracts. The influence of SH is again well illustrated by the 27% increase in the amount of creatine measured in the presence of added CMBA.

Some experiments have been carried out with urine from normal males, young children and pregnant females. These have been neither sufficiently numerous nor comprehensive to quote in detail, but some of the preliminary results are of interest. It has been found that some degree of inhibition is present in 1.0 ml. of urine as judged by the increased values in the presence of CMBA. These increases were variable and have not been correlated with other factors, but were of the order of 10%. 100% recoveries of added creatine were obtained by the addition of CMBA, and it therefore appears possible that this modification may prove useful for the estimation of creatine in urine.

DISCUSSION

It has been suggested by Eggleton *et al.* (1943) that ω -amino-acids may be responsible for the failure of the Barritt reaction to estimate creatine in TCA-

liver extracts and in urine. They further state that extracts of most animal tissues, with the exception of liver, 'appear to contain insignificant amounts of interfering substances'. An examination of the ethanol-insoluble barium salts obtained from TCA extracts of liver led us to the conclusion that GSH was more likely to be the agent responsible for the inhibitory effect, and further that it is the SH group which is the determining factor.

The inhibitory influence of SH-containing compounds on the α -naphthol-diacetyl reaction for the estimation of creatine is very striking and depends on the molar ratio of SH: creatine. In the estimation of creatine in spleen, kidney, lung and intestine, the SH: creatine ratio is sufficient to produce errors ranging from 60 to 250%.

The inhibitory action of SH can be removed almost completely by the addition of *p*-chloromercuribenzoic acid, and the figures obtained in the presence of this compound suggest that the method, so modified, is now capable of giving creatine values

which are close to the true values. An effect, similar to that obtained with this substance, could probably be gained by the addition of other organic mercurials and, in fact, by any substance reacting with SH groups to give a complex stable to alkali; but since *p*-chloromercuribenzoic acid behaved satisfactorily in our hands this has been used exclusively.

The results of experiments with pure SH compounds and SH-containing tissue extracts in the presence of *p*-chloromercuribenzoic acid strongly suggest that SH compounds are the only inhibitors of importance as far as the application of the Barritt reaction to tissue extracts is concerned. This conclusion is further strengthened by the fact that, in most instances, recoveries of known amounts of added creatine were 90% or greater. Although Eggleton *et al.* (1943) have observed a 50% reduction of the colour produced by 30 μ g. of creatine in the presence of 4 mg. of glycine it seems improbable that such amounts of any amino-acid could be present in the usual samples taken for estimation of creatine in TCA-tissue extracts.

The only body fluids examined have been whole blood and urine, and in both cases the addition of CMBA appears to overcome the inhibitors which

are present. The suggestion of Walpole (1911) that it is the NH_3 which interferes with the reaction in normal urine has not been confirmed.

It follows from the high degree of sensitivity of this reaction to the presence of sulphydryl compounds that use might be made of this fact for the detection, and possibly the estimation, of small amounts of SH and alkali-labile SS substances. Experiments are in progress to test this possibility.

As Barritt (1936) has pointed out, the reaction between diacetyl, α -naphthol and creatine to form a coloured product is also given by other guanido compounds. Eggleton *et al.* (1943) have stated that arginine, guanidine and glycoeyamine produce about one ninth of the colour produced by equivalent amounts of creatine. All results quoted in this communication have been calculated on the assumption that creatine alone produces the colour measured. It is believed that the error introduced by such an assumption is small. However, in the application of the revised method, the possibility of increased amounts of other guanido compounds in pathological states must be borne in mind.

SUMMARY

1. The α -naphthol—diacetyl reaction for the estimation of creatine has been investigated.

2. The inhibition, which has been noted by other workers to occur when the estimation of the creatine content of trichloroacetic acid extracts of liver is attempted, has been shown to be due to the presence of sulphydryl compounds and, by inference, to the presence of glutathione.

3. The inhibition by sulphydryl compounds depends on the molar ratio of the sulphydryl compound to creatine, and is therefore most marked in the liver where the creatine content is low and the glutathione content high.

4. The ratio of sulphydryl compounds to creatine, in several tissues investigated, is sufficiently high to produce very gross errors in the estimation of creatine by this method. Exceptions are found in voluntary and cardiac muscle, brain and testis, but even here the method is subject to error.

5. The presence of disulphide compounds such as GSSG also produces an inhibition due, presumably, to the tendency of the disulphide linkage to split in alkaline solution.

6. The inhibition produced by sulphydryl compounds can be almost completely removed by the addition of *p*-chloromercuribenzoic acid; both this compound and that formed by its reaction with the sulphydryl compounds possess weak inhibitory powers.

7. The creatine content of various tissues, blood and urine has been determined both in the presence and in the absence of *p*-chloromercuribenzoic acid and an assessment made of the errors introduced by the omission of this compound.

8. The recoveries of creatine added to extracts of various tissues, blood and urine are mostly in excess of 90 %, thus indicating the general applicability of the modified method.

We wish to express our thanks to the Chief Scientist of the Ministry of Supply for gifts of methionine and benzyl mercaptan and to Messrs E. Facer and S. Moorbath for skilled technical assistance.

REFERENCES

- Barritt, M. M. (1936). *J. Path. Bact.* **42**, 441.
Eggleton, P., Elsdon, S. R. & Gough, N. (1943). *Biochem. J.* **37**, 526.
Ennor, A. H. & Stocken, L. A. (1948). *Biochem. J.* **42**, 549.
Friedmann, E. (1903). *Beitr. chem. Physiol. Path.* **3**, 1.
Hunter, A. (1928). *Creatine and Creatinine*. London: Longmans Green.
Le Page, G. A. & Umbreit, W. W. (1945). *Manometric Techniques and Related Methods for the Study of Tissue Metabolism*, by W. W. Umbreit, R. H. Burnis & J. F. Stauffen. Minneapolis: Burgess Publishing Co.
Potter, V. R. & Elvehjem, C. A. (1936). *J. biol. Chem.* **114**, 495.
Stocken, L. A. (1947). *J. chem. Soc.* p. 592.
Vosges, O. & Proskauer, B. (1898). *Z. Hyg. InfektKr.* **23**, 20.
Walpole, G. S. (1911). *J. Physiol.* **42**, 301.
Whitmore, F. C. & Woodward, G. E. (1932). *Organic Synth. Collective* **1**, 153.