XII. THE DISTRIBUTION OF NITROGEN IN BEER.

By JOHN SMITH SHARPE.

(From the Department of Physiology, Glasgow University.)

(Received May 25th, 1917.)

The value of beer as a food has been for long a subject of controversy. There can be no doubt that the alcohol and sugars have a certain value as sources of energy, but whether the nitrogenous compounds are of any value has never been fully investigated.

In the analysis of beer it is customary to determine the nitrogen content, to multiply this by 6.25 and to express the result as protein—a totally fallacious procedure.

In the Report of the Committee of the Royal Society on the Food Supply of the United Kingdom it is stated “that the nitrogen compounds of beer have been little studied,” and so incomplete is our knowledge as regards their nature that the committee merely made a guess in assuming that something between 50 and 100 % of the nitrogen is in the form of protein or amino-acids which are utilisable in the body.

The object of the present paper is to throw further light upon the nature of these nitrogenous compounds.

Fate of the Proteins of Barley during the Brewing Process.

As a result of the malting, part of the protein in the grain is acted on by proteolytic enzymes which break it down into simpler substances. These along with a little soluble protein pass into solution during the mashing but the greater part is left in the brewers' grain. During the processes of boiling, hopping and subsequent cooling of the wort some of this protein becomes precipitated.

The amount of nitrogen thus finally left in solution is termed by the brewer “permanently soluble nitrogen.” By the action of the yeast during fermentation a further quantity of soluble nitrogen is removed by these organisms using a certain proportion as food.

A further amount of protein is removed after the beer is made and this process is carried out to ensure that the beer may remain "bright."

The following table is taken from analyses by Bungener and Fries [quoted by Chapman, 1912, p. 14]. It shows the increase in the soluble nitrogen in the production of malt from barley.

TABLE I.

	Barley	Malt
Total nitrogen	1.690 %	1.580 %
Nitrogen as albumin	0.161	0.230
„ peptone	0.040	0.060
„ amides	0.206	0.534
Total soluble nitrogen	0.355	0.642

Thus it will be seen that in barley there is about 20 % of the total nitrogen in a soluble form, while in malt almost 40 % exists in this form. Of this 40 %, which during the period of mashing goes into solution, a certain proportion as already stated remains in the wort as "permanently soluble nitrogen." A large amount of research work on these bodies has been undertaken by Horace Brown [1907].

He accounts for almost 66 % of this "permanently soluble nitrogen" as the following table details:

TABLE II.

Ammonia nitrogen ...	...	3.5 %
Malt-albumose nitrogen ...	...	20.0
Malt-peptone nitrogen ...	...	31.0
Amide and amino-nitrogen ...	...	8.5
Nitrogen due to organic bases ...	...	4.0
Balance (unclassified) ...	...	33.0
		<hr/> 100.0

The 4 % of nitrogen due to organic bases comes from betaine and choline, and allantoin (glyoxyldiureide) was also present on the wort (Horace Brown).

The proportion of nitrogen capable of being assimilated by the yeast was ascertained by Horace Brown. By successive croppings of yeast and the estimation of the nitrogen content of the wort before and after fermentation he found that almost 60 % could be accounted for by the yeast action. This was a laboratory experiment and the action of the yeast was continued much longer than it is in practice. However the same test was carried out under brewing conditions with the result that the assimilated nitrogen varied between 20 and 36 % of the permanently soluble nitrogen.

From the foregoing it will readily be seen that the protein content of the beer or ale will not be appreciable when it reaches the consumer.

Application of Protein Tests.

1. The biuret test was tried both directly and on the isolated protein bodies. In every case negative results were obtained. The same result has also been obtained by Horace Brown.

2. Millon's reagent gave a negative result.

3. The ninhydrin test was also negative.

4. The xanthoproteic test alone gave a positive result.

Table III gives the percentage of so-called protein in different ales, calculated from the total nitrogen as given in the second column [Chapman 1912].

TABLE III.

	Proteins	Total nitrogen
London Mild Ale	0.33 %	0.052 %
Burton Pale Ale	0.50	0.080
English Strong Ale	0.91	0.145
Dublin Stout	0.75	0.120
American Ale	0.57	0.091

The only figures for the distribution of nitrogen in beer which I have been able to find are by Voltz, Förster and Baudrexel [1910] who give an analysis of North German dark beer used by them in their investigation of the absorption and general action of beer.

TABLE IV.

Alcohol ...	...	...	...	4 %
Extract ...	...	...	...	5.7

Analysis of Extract.

Ash ...	...	...	...	3.44
Organic ...	...	...	...	96.56
Nitrogen $\times 6.25$...	...	...	...	6.36 = 1.01 % N.
Ethereal extract	...	...	...	0.13
Nitrogen-containing substances:				
Protein ...	...	...	...	42.7
Amide-N. ...	...	...	...	57.3

METHODS.

(1) Total Nitrogen. This was determined by the Kjeldahl method using 10 cc. of the sample. The Kjeldahl-Jodlbauer [see Sutton, 1904, p. 87] modification was tried but no advantage was gained since no nitrogen seems to exist as nitrates or nitrites.

(2) Protein Nitrogen. 25 cc. of the sample were evaporated to half bulk and then cooled to about 10°, 10 % tannic acid solution containing 2 % phosphoric acid (about 4 vols. to 1 of sample) was added, shaking the while,

until no further precipitate formed. The protein mass was allowed to flocculate, then filtered off on nitrogen-free filter paper and washed with the precipitating solution. The nitrogen in the precipitate was then determined by the Kjeldahl method.

Nitrogen found $\times 6.25$ = Total Protein.

Tannic acid solution used by itself as a precipitant, may bring down bases along with the protein but this is prevented by the solvent action of the phosphoric acid. By too lavish use of tannic acid an error may be incurred, (it is a case of the mutual precipitation of two colloids) with consequent loss of protein [Barger 1914, p. 117].

Other methods of taking out the proteins were tried but it was found that the above gave the most satisfactory results. The lead acetate precipitation was investigated. This salt has a well recognised power of throwing proteins out of solution, especially if a little of the tribasic acetate is added. The results with this were far from satisfactory as the following figures will show :

Protein nitrogen by tannic acid method	0.021 %
„ „ acetate method	0.011 %

The failure of lead acetate to precipitate all the protein is difficult to explain, unless it be that the metallic salts of these plant proteins are more soluble than those in the corresponding animal series. Further light on this point would be interesting.

The use of (a) copper acetate or (b) phosphotungstic acid as a precipitating agent is open to serious objection as both precipitate purine and basic bodies. These were therefore rejected.

(3) Amino-acid Nitrogen. The estimation of the nitrogen combined in this form was effected by the nitrous acid method of van Slyke [1912]. 5 cc. of the ale or beer were run into the deaminising flask and this was acted on by 20 cc. of the nitrous acid mixture. Numerous duplicate estimations were performed and an average of the whole was taken. Amide-nitrogen is included in the figures given.

A full description of the process employed by previous workers for the separation of the amino-acid and amide-nitrogen is given by Horace Brown [1903].

(4) Purine Nitrogen. (a) *Qualitative Tests*. The presence of purine bodies was proved by the following tests :

1. The murexide test.
2. Precipitation by ammoniacal silver nitrate and magnesia mixture.
3. Precipitation after adding copper sulphate and sodium bisulphite.

4. The intense blue colour produced on adding phosphotungstic acid reagent (Folin) to the material made alkaline with sodium carbonate. This Folin considers characteristic of uric acid (trioxypurine).

(b) *Quantitative Tests.* The estimation of the uric acid content was carried out by the new method of Folin and Denis [1913] for its estimation in urine. It involves the precipitation as a silver magnesium compound with subsequent treatment by the phosphotungstic acid reagent and comparison of the colour produced with a standard. This method was used in the case of dark coloured beer and stout where the original colour of the material would have interfered greatly with the delicacy of the reaction. It was however found that in light ales or beers quite accurate results were obtained by direct treatment with Folin's reagent, but that care had to be exercised not to allow too much time to elapse after the addition of the reagent owing to the liability of the colour to fade on standing longer than five minutes. Of the beer under examination 2 cc. were therefore drawn off and placed in a test tube; 5 cc. of saturated sodium carbonate solution were next added and then from ten to twenty drops of the phosphotungstic acid. The mixture was shaken together and slightly heated to facilitate the reaction. This was washed into a Nesslerising-tube to about 25 cc.

The standard was prepared in a similar tube by mixing together 20 cc. of water, 5 cc. of saturated sodium carbonate solution and about 20 drops of the phosphotungstic acid, a lithium carbonate solution of uric acid (1 cc. = 1 mg. uric acid) was then delivered from a burette into the tube containing the standard until the depth of colour in each was alike. Both volumes were then made the same. These were further compared in a Duboscq colorimeter and any necessary adjustment made. The number of cc. of the standard solution used multiplied by 0.333 gives the equivalent in trioxypurine nitrogen¹.

(c) *Xanthine-like purines.* These were determined by a modification of Salkowski's [1895] method for urine. The proteins were removed by lead acetate, the lead removed by H_2S gas. The phosphates were removed from the filtrate by magnesia mixture. Ammoniacal silver nitrate was then added and the resulting precipitate was filtered off and suspended in water, decomposed with H_2S and filtered. The filtrate was evaporated to a syrup and the residue treated with a little water containing 2 % of sulphuric acid which dissolves the xanthine bases leaving the uric acid practically insoluble.

The purine bases were reprecipitated by ammoniacal silver nitrate and

¹ A table giving the purine content of some beers and ales is given by C. von Noorden [1907].

either estimated by the quantity of silver in the precipitate or by the quantity of nitrogen given by the Kjeldahl method. For this work the nitrogen in the precipitate was estimated.

Xanthine seems to form the greater bulk of these more soluble purines. Confirmatory tests were given by the base liberated from the silver compound. The following may be quoted:

(a) Strecker's test. On evaporating the residue on a watch glass with a drop of nitric acid a yellow residue remains which turns reddish yellow on addition of potassium or sodium hydroxide.

If ammonia be substituted for the fixed alkali, no violet coloration is obtained. This is a distinction from uric acid.

(b) The residue was soluble in nitric acid and on crystallising, characteristic nitrate crystals were formed.

(c) Solutions of the residue gave a yellow crystalline precipitate with sodium picrate; at first it seemed to be amorphous, then assumed the form of round bunches of crystals.

Hypoxanthine, adenine and guanine were tested for, but none of these was found. Methylpurines were in every case absent.

The proportion of nitrogen existing in the different forms of combination found by the above processes, *b* and *c*, varies greatly. Thus the nitrogen as trioxypurine may be as much as 90 % of the total purine nitrogen or as low as 15 or 20 %, the sum being completed by the nitrogen combined in the xanthine-like bodies.

The "purine nitrogen" reported in Tables V to VIII is the sum of the nitrogen found by the above two processes *b* and *c* (p 105). This gives the total purine nitrogen.

(5) Total solid matter. This was estimated by drawing off 10 cc. from a good average sample of the beer or ale in question. It was then placed in a tared evaporating basin and evaporation continued on a water bath until dry. It was then transferred to a vacuum desiccator and dried until constant in weight.

(6) Carbohydrate. The amount of carbohydrate was determined after precipitation of the protein using Wiley's mercuric nitrate method [see Allen 1914, p. 157]. 25 cc. of the material were placed in a 100 cc. flask, 10 cc. of mercuric nitrate solution were added and the whole thoroughly mixed. When the froth settled, water was run in up to the mark on the neck and the contents again shaken. No allowance need be made for the precipitate in this case as it is so very small. When the supernatant liquid

was clear, a little (about 30 cc.) was filtered off through a fine paper (Green's 602). The concentration of sugars was then found by the polarimeter the whole being calculated to dextrin.

The results obtained by calculating the rotation all to dextrin are slightly low, about 0.3 % where a 3 % dextrin content is given. This is due to the maltose present having a slightly less specific rotation than dextrin.

The formation of an osazone derivative was tested for. Maltose was proved by the formation of maltosazone crystals, but no glucosazone was found.

Levulose was tested for, but negative results were obtained (Selivanov's resorcinol test). This sugar is very easily fermented by yeast whereas dextrin is not acted on, hence the latter is, on an average, a little above 66 % of the total carbohydrate present, the remainder being maltose [Chapman 1912].

RESULTS.

The following tables show the results of analyses of some typical beers, ales and stouts. The volume of each sample is noted.

TABLE V.

Beer (light).

	Tennent's Lager Beer		Draught Beer	
Total nitrogen	—	0.040 %	—	0.57 %
Protein nitrogen	0.006 %	—	0.021 %	—
= Total protein	—	0.038	—	0.133
Amino-acid nitrogen	0.017	—	0.014	—
Purine nitrogen	0.016	—	0.019	—
Undetermined nitrogen	0.001	—	0.003	—
Total solid matter	4.89 %		3.47 %	
Carbohydrate (dextrin)	2.46		1.30	
Specific gravity	1006.8		1006.1	
Bottle contained	250 cc.		550 cc.	

TABLE VI.

Beer (strong).

	Bass & Co.'s Pale Ale (A)		Bass & Co.'s Pale Ale (B)		
Total nitrogen	—	0.084 %	—	0.082 %	
Protein nitrogen	0.011	—	0.016	—	
= Total protein	—	0.074	—	0.102	
Amino-acid nitrogen	0.036	—	0.040	—	
Purine nitrogen	0.039	—	} not done	—	
Undetermined nitrogen	0.004	—			
<hr/>					
Total solid matter	6.37 %		6.20 %		
Carbohydrate (dextrin)	3.06		3.00		
Specific gravity	1009.4		1009.0		
Bottle contained	330 cc.		265 cc.		

TABLE VII.

Ales.

	Younger's Edinburgh Pale Ale		Worthington's Indian Pale Ale	
Total nitrogen	—	0.039 %	—	0.063 %
Protein nitrogeen	0.010 %	—	0.010 %	—
= Total protein	—	0.063	—	0.063
Amino-acid nitrogen	0.018	—	0.016	—
Purine Nitrogen	0.010	—	0.033	—
Undetermined nitrogen	0.001	—	0.004	—
Total solid matter	4.89 %		4.92 %	
Carbohydrate (dextrin)	2.46		2.24	
Specific gravity	1008.3		1007.0	
Bottle contained	535 cc.		330 cc.	

TABLE VIII.

Porter.

	Allsopp's Special Stout		Guinness's Extra Stout	
Total nitrogen	—	0.056 %	—	0.113 %
Protein nitrogen	0.012 %	—	0.029 %	—
= Total Protein	—	0.080	—	0.185
Amino-acid nitrogen	0.019	—	} not done	
Purine nitrogen	0.020	—		
Undetermined nitrogen	0.005	—		
Total solid matter	5.90 %		—	
Carbohydrate (dextrin)	2.05		—	
Specific gravity	1009.9		1010.0	
Bottle contained	305 cc.		310 cc.	

(All the above analyses were made in duplicate.)

Table IX shows the percentage on the total nitrogen of the nitrogen combined in various forms.

TABLE IX.

	Tennent's	Draught	Bass's	Younger's	Worthing- ton's	Allsopp's	Guinness's
Protein N in total N	15.1	37.0	13.2	25.8	15.9	21.4	25.0
Amino N ,,	42.5	24.5	35.7	46.0	25.4	33.8	—
Purine N ,,	40.0	33.2	46.4	25.8	52.4	35.7	—
Undetermined N in total N	2.4	5.3	4.7	2.4	6.3	9.1	—
	100	100	100	100	100	100	—

Consideration of results.

(1) From a study of Tables V to VIII it will be seen how close the figures for the total nitrogen are, to those given in Table III from which previous workers have calculated the total protein. According to the present investigation only about 22 % of the total nitrogen exists as protein.

(2) Horace Brown [1907] in his researches on malt extract obtained a mixture of crystalline substances. These proved to be asparagine (amino-succinamide), tyrosine (*p*-hydroxyphenyl- α -aminopropionic acid) and leucine (isobutyl- α -aminoacetic acid).

These amino-acids will ultimately find their way into the beer and it is probable that they may be the source of the amino-nitrogen.

(3) As regards the purines there is some evidence to show that these bases may be derived directly from the malt, hops, and yeast. A. C. Chapman has isolated adenine, hypoxanthine and xanthine from these but he does not give data as to the percentage of the bases.

With regard to the malt and yeast the presence of purine bases in these seems to be fairly well established. As far as the present investigation is concerned xanthine (dioxypurine) and uric acid (trioxypurine) were present in comparatively large quantities. Further, since these have been found in the present work and allantoin has been found in wort by previous workers [Brown 1907] it is possible that an addition to the purine nitrogen may result from the hydrolysis and oxidation of nucleic acid [Mathew, 1915, p. 729], partly from the grain and partly from the yeast.

EXAMINATION OF THE UNDETERMINED NITROGEN.

(1) *Alkaloids.*

On extracting a quantity of the beer made alkaline by sodium hydroxide with chloroform and evaporating off the solvent, a clear oil was obtained which turned brown on exposure to air, and assumed an odour very like that of tobacco juice from a pipe. It seemed also to be very volatile as it had to be dried in a desiccator for fear of loss which rapidly took place on drying at 100°.

A. Colour tests on the isolated alkaloid.

(i) A pale red colour was immediately produced by a drop of concentrated sulphuric acid in the cold. This colour was discharged on addition of a trace of nitric acid.

(ii) Sulphomolybdic acid gave a blood red colour in the cold which became darker on heating.

(iii) On evaporation with nitric acid a yellow colour was produced which turned rose pink on addition of alcoholic potash.

(iv) A deep pink colour was produced on heating with concentrated hydrochloric acid. This colour is discharged on addition of alkali.

B. The alkaloid dissolved in hydrochloric acid and in nitric acid quite freely and was liberated on addition of excess of caustic soda giving off the same odour as it had when extracted. It was proved by experiment that this alkaloid was not precipitated from its solutions by the tannic acid mixture used for the protein extraction, it is therefore another source of nitrogen.

The alkaloid seems to belong to the pyridine group. Many of the reactions are typical of "coniine" which has been found in hops by Chapman [1914]. The following table shows the percentage of alkaloid in a few samples of beer etc. examined.

TABLE X.

	Alkaloid
Draught Beer, Paterson & Co. ...	0.007 %
Younger's Edinburgh Ale ...	0.004
Allsopp's Stout ...	0.010
Guinness's Stout ...	0.012

(2) *Bases.*

These were taken out with a solution of phosphotungstic acid. The precipitate was filtered off and suspended in water, decomposed with barium hydroxide and filtered. The barium was removed by a current of CO_2 gas and the precipitate filtered off. Purines were then precipitated by ammoniacal silver nitrate solution, allowed to settle and filtered. The excess of silver was removed from the filtrate by hydrochloric acid and the solution concentrated to a syrup. Alcoholic mercuric chloride solution was added which precipitated the bases. This precipitate was decomposed with H_2S gas and the filtrate again concentrated. Picric acid solution was added and the whole set aside to crystallise. Bunches of yellow needles were formed which were fairly insoluble in water and had a melting point a little over 181° . This corresponds to the commonly occurring plant base, betaine, which has been found by previous workers in wort, see page 102. There was however an extremely small percentage of this in beer, only 0.005 % of the picrate, on an average, being isolated.

CONCLUSIONS.

1. The protein nitrogen of the beers analysed, varies from 13.2 to 37.0 % of the total nitrogen, *i.e.* the protein content of beer varies from 0.038 to 0.185 %.
2. The nitrogen as amino-acid varies from 25 to 46 % of the total nitrogen, *i.e.* the amino-acid nitrogen of beer varies from 0.014 to 0.040 %.

3. There is a varying proportion of nitrogen combined as purine compounds which in some cases exceeds that combined as amino-acid nitrogen. It varies from 25.8 to 52.4 % of the total nitrogen or from 0.010 to 0.039 % of the beer itself.

4. The undetermined nitrogen of these beers varies from 2.4 to 6.3 % of the total nitrogen, *i.e.* the percentage on the beer itself varies from 0.001 to 0.005.

5. The percentage of undetermined nitrogen is not quite made up by the addition of the nitrogen combined as alkaloid and base.

In the case of Allsopp's stout there is only 0.0011 % of nitrogen combined as the former and the stouts seem to contain the largest amount of alkaloid.

Of basic nitrogen there is only 0.0002 %, which is negligible.

I am indebted to Professor D. Noël Paton for his constant encouragement and advice during this work.

The expenses connected with the same were defrayed by the Medical Research Committee, to which body I tender my thanks.

REFERENCES.

- Allen (1914). *Commercial Organic Analysis*, 8.
Barger (1914). *The Simpler Natural Bases* (Longmans, London).
Brown, Horace (1903). *Trans. Guinness Research Laboratory*, i. i. 18.
Brown, Horace (1907). *J. Inst. Brewing*, New Series, 4, 394.
Chapman, A. C. (1912). *Brewing* (Cambridge, University Press).
Chapman, A. C. (1914). *J. Chem. Soc.* 105, 1895.
Folin and Denis (1913). *J. Biol. Chem.* 14, 95.
Mathew (1915). *Physiological Chemistry*.
Noorden, C. van (1907). *The pathology of metabolism*, 3, 1297.
Salkowski (1895). *J. Chem. Soc.* 2, 538.
Sutton (1904). *Volumetric Analysis*, 9th Ed.
Slyke, D. D. van (1912). *J. Biol. Chem.* 12, 278.
Völtz, Förster and Baudrexel (1910). *Pflüg. Arch.* 134, 133.