VI. ON THE COLOURS OF TWO SEA ANEMONES, *ACTINIA EQUINA* AND *ANEMONIA SULCATA*.

PART I. ENVIRONMENTAL.

By RICHARD ELMHIRST.

PART II. CHEMICAL.

By JOHN SMITH SHARPE.

*From the Marine Biological Station, Millport, and the Department of
Physiology, University of Glasgow.*

(Received December 24th, 1919.)

(With Plates III, IV.)

MANY and varied are the theories that hold the field of colour phenomena in nature. This is to be expected, for although a large amount of work has been published on this subject, very little of it takes one further than the surface. There are the theories of Darwin, Poulton, Wallace, Eimer, Simroth, etc., which are all based on a hypothesis of structure in relation to natural selection or environment, the works being so well known that it is unnecessary for us to enter into any detail here. The aim of the present investigation is to endeavour to obtain an interpretation for certain colours.

Part I.

ENVIRONMENTAL.

The commonest British Actinian is *Actinia equina* L., occurring on rocks and stones all around our coasts. Several colour varieties have been described and recorded in varying abundance from many localities. The species is abundant in the Clyde area: the prevailing form being the normal liver-red type, with bright blue "marginal spherules" and an equally blue border round the base, var. *hepatica* [Gosse, 1860, Pl. 6, fig. 2].

Ideal Habitat.

The ideal habitat for this species, as shown by the accompanying table, is about half-tide mark among the dense growths of *Ascophyllum* and *Fucus serratus*. The luxuriant growth of these weeds provides shelter and moisture during ebb and shade and protection from the force of the waves during flood. The presence of these weeds is, however, not essential because *Actinia* may occur abundantly on shelving rocks where they are replaced by *Chondrus* and *Gigartina*, short crisp weeds some three to four inches high.

In rock pools *Actinia* may occur at any tidal level up to and rarely above high water mark.

Vertical distribution of *Actinia*.

Sea-weed zone	Exposed upper surface of stones	Sheltered
<i>Pelvetia canaliculata</i>	—	Few
<i>Fucus platycarpus</i>	Rare	Few
<i>Fucus vesiculosus</i>	Few	Frequent
<i>Ascophyllum nodosum</i>	Common	Common
<i>Fucus serratus</i>	Common	Common
<i>Laminaria digitata</i>	Few	

Colour.

Within the limits of var. *hepatica* there are two shades of colour. (1) A deep red, occasionally tending almost to blackness, which occurs in exposed situations and (2) a paler bright red which occurs in shelter under sea-weed, below stones and in crevices. This agrees with Walton's observations on *Actinia equina* at Aberystwyth and in South Devon [1911, pp. 229 and 239]. Farquhar's on *Actinia tenebrosa* in New Zealand [1898] and Saville-Kent's on the coral *Euphyllia* [1893]. Of the other colour varieties described by Gosse, the brown var. *umbrina* forms less than 1 per cent. of all examined except in a few favoured localities where a pale var. *umbrina* seems to have become established. These varieties (*hepatica* and *umbrina*) merge through red browns so that it is often a matter of considerable difficulty to decide how to describe a particular specimen. Red browns and browns both vary in depth of colour according to the exposure of their surroundings. Two other varieties, the sea-weed coloured *olivacea* and red with green marks *fragacea*, are rare—less than 1 in 5000. Clyde var. *fragacea* are always small, 1 to 1.5 cm., apparently never attaining the inches of the Devon and Cornish specimens.

Relationship of Colours.

It is well known that the distinct colour varieties are connected by intermediate shades, which suggests that the pigments are chemically related to each other and probably derivatives from the same source. This is also borne out by several experimental observations of individuals changing colour, notably the fading of brilliant markings. Similar changes have been recorded for *Madrepora prostrata* [Saville-Kent, 1893].

In 1910 a normal *A. equina* was placed in a well-lit glass-sided aquarium. By the next year this specimen had passed from red through red-brown, brown-green to a deep olivaceous-green, i.e., had assumed the coloration of var. *olivacea* and attained a diameter of from 4 to 6 cm. This assumed coloration has persisted through a number of generations from 1911 to 1919. The young are usually pale greyish green about the shade of Gosse's var. *glauca*. A few, however, show a faint reddish tinge but in all cases the final colour has been that of var. *olivacea*.

Experiments are now in progress (1) to repeat the assumption of the green colour and (2) to reverse the process and get the var. *olivacea* to reassume the red of var. *hepatica*. With the former aim, three normal red var. *hepatica* were placed on May 10th under surroundings precisely similar to those in which the change *hepatica* to *olivacea* took place in 1910-11. Now, after five months, they have lost much of their original red and the tentacles are completely brown which, so far, is a repetition, though slower, of the 1910-11 process. With the latter aim a number of var. *olivacea* are being subjected to alternate dryness and submersion, while a second set are being kept in an artificial rock pool cut in the local sandstone. At present, after three months, no very definite results are noticeable but a suggestion of redness is appearing above the blue basal border in two cases.

Inheritance of Colour.

It is interesting to note that this assumed coloration has so far bred true. Gosse mentions [1860, pp. 178-9] that vars. *cerasum* and *chiococca* breed true. The same is suggested by the frequent occurrence on the shore of the young of any particular variety near an adult of that variety. At times a variety may establish itself in a suitable locality.

There is a rock-sheltered corner on the east shore of Cumbrae at the *Fucus serratus* level where *Metridium dianthus* was always found, both the red and white varieties (1906-15). On visiting this place in 1919 *Metridium* was found to be absent and in its place were a large number of pale brown *A. equina*, in colour between Gosse's vars. *umbrina* and *ochracea*. There were some 250 specimens of this variety which had established itself for a number of yards to the almost total exclusion of the normal red.

Colour Significance.

The significance of these colour forms has been ably discussed by Walton [1911] who classifies the coloration of actinians as

- (1) Warning;
- (2) Aggressive;
- (3) Protective;
- (4) Colours with some special physiological significance.

In *Actinia equina* the brilliant blue of the "marginal spherules" has been generally accepted as warning coloration. This may well be so, but unfortunately we have no observations bearing on the matter, although the fact that the normal colour often contrasts strongly with the environment may bear this interpretation. Because of this environmental colour contrast it is improbable that the aggressive coloration suggestion holds true although crustacea in aquaria have occasionally been seen to stumble onto an *A. equina*. That the normal colour is protective to any extent will be disproved by a single walk along the shore, despite the redness of the sandstone in several

localities. If the coloration were protective varieties should occur in definite association with environment, e.g. rock pools filled with *Cladophora*, backgrounds of sandstone, basalt or limestone, yet the normal red prevails in all these surroundings. In the *Cladophora* pools, particularly, dark reds which presented a marked contrast to their surroundings have been found occurring abundantly, 800 specimens without a single variety present. This brings us to colours with some special physiological significance, which interpretation is strongly suggested by (1) the variation in colour correlated with exposure already mentioned, and (2) the well-known fact that brightly coloured specimens when kept in aquaria and therefore continually submerged tend to lose their distinctive coloration. The obvious inference seems to be that the pigment is a light screen which has some physiological significance, possibly respiratory, since change of colour often takes place when specimens are kept continually submerged and not exposed to the alternate ebb and flow of the tide.

Anemonia sulcata (Pennant)=*Anthea cereus* (Ellis and Sol), occurs abundantly in suitable localities all round our southern and western coasts. In the Clyde, the normal "umber brown" is frequent on *Laminaria*, where it closely resembles the background of sea-weed: a case of aggressive coloration which obtains "much food in the form of deluded crustacea."

The ideal habitat of this species is a bed of *Laminaria digitata* sheltered from heavy waves and having a good exposure to sunlight. The first condition is favourable to the species because it is incapable of strong attachment and the second is essential for the life of the symbiotic algae inhabiting the tentacles. Excellent examples of this ideal habitat are seen in Castle Bay, Little Cumbrae, and between the Eilans in Millport Bay; the species being abundant in both places.

The need of direct strong sunlight is very well illustrated on the walls of the cambers in Gibraltar harbour, where on the walls running east and west, the north sides are practically devoid of *Anemonia* whereas the south sides are covered with thousands of fine specimens of the beautiful rose-tinted variety *smaragdina*.

SUMMARY.

1. The ideal habitat is at half-tide level.
2. Intensity of colour varies with exposure to light.
3. Individuals may show (environmental) colour change.
4. Individuals breed true.
5. A given variety may establish itself in suitable environment. In the present absence of evidence of the actual difference in environments it seems that one shade of colour may be as effective for the success of the species as another.
6. The coloration seems to have a special physiological function as a light screen.

Part II.**CHEMICAL.***Apparatus used.*

A Browning microspectroscope, with an inch objective and condenser, was used. The solutions of the colours were placed in small flat-bottomed glass tubes of about 10 cc. capacity. A compressorium was used for the examination of the fresh tissue which was cleared in glycerol. The positions of the various absorption bands and screens were determined by the Fraunhofer lines on a comparative solar spectrum.

Actinia equina.

1. *The ether-alcohol-soluble portion.* In the first part of the chemical work the interest was centred round substances of the nature of lipochromes. The fat solvents were all tried with varying success although the extracts were weak. In all cases the best extract of the ectodermic lining where the colouring matter was contained, was obtained by treatment with a mixture of three parts ether to one of alcohol. This gave in every case strong extractions of the red, brown-red or brown-yellow colours. On evaporation, oily drops were left corresponding in colour with the original coloured solution. These did not mix with water but lay on the surface as an oily layer. No haemoglobin derivatives could be demonstrated in this colouring matter; it is therefore not respiratory in function, at least as an oxygen carrier. Acids produce no change in the colour but on the addition of alkalies, NaOH for example, a brown colour resulted and the spectrum was changed. (See Plate III, A, spectrum 6.) This is important as it involves the possibility of acid or alkali playing a part in the colours of these animals, but whether this is produced by food or light is a question to be considered. There is one point that is significant in the absorption spectra of Plate III, A, that is the screening over the green and red portions of the spectrum.

According to Grotthus' law (1819) photo-chemical action cannot take place unless light is absorbed. Now Engelman has shown that red sea-weeds show the greatest carbon assimilation in the green and by the spectrum they show the greatest absorption. The chlorophyll in these red weeds is modified and acts as an optical sensitiser. This same effect is produced by light of various wave lengths, provided they are absorbed.

Again photo-chemical action seems not to depend on molecular construction, for with colouring matters ranging from red, yellow to blue, the molecules of which are not arranged in the same way, it has been shown that in every case where light is absorbed iodine can be set free by the released oxygen from potassium iodide, independent of absorption wave-length or position on the spectrum [Wager, 1914].

It will readily be seen that there is in these anemones a substance absorbing light in the green and red parts of the spectrum and generally resembling

chlorophyll in its action on light. This substance although not showing the distinctive bands of chlorophyll gives absorption wave-lengths in almost the same position as the maximum absorptions of chlorophyll.

Furthermore it was proved experimentally by Wager's method [1914] that these red and brown substances produce photo-chemical action with the release of active oxygen.

By the use of the fluoroscope a faint fluorescence is seen in the red almost over the lines *B*, *C*. This is in the position of maximal energy of the chlorophyll system as found by Engelmann and Timiriazev. According to Kerner red fluorescent substances have the power to transform the blue rays to red and ultra-red rays or in other words, rays of light into chemical rays. It is highly probable therefore, that such an action takes place in these anemones and as such may be concerned chiefly in their constructive metabolism.

This has also a bearing on the respiratory pigment which in these animals is very small in amount and must act more as a fixer than a carrier of oxygen.

Thus it will be seen that adequate conservation of the light is all important and that the colouring matter by virtue of the heavy screen towards the blue end affords protection against the harmful ultra-violet light.

In brown specimens a powerful screening of the blue and violet end is again secured by a brown substance soluble in ether-alcohol of the nature of a lipochrome. In spectrum 4, Plate III, A, it will be seen that even with a thin layer the screening towards the violet is very marked. The other parts of the screen behave in very much the same way as those of the red type.

In the lighter sheltered red and brown anemones there is only a variation in the intensity of colour but the photo-chemical nature remains the same.

2. *The respiratory pigment.* Portions of the ectoderm of a red and a brown anemone were cleared in glycerol and examined spectroscopically in a compress. Two bands were visible as indicated in Plate III, B, spectra 1 and 3. It will be noticed that in spectrum 3 the bands are a little further towards the blue end. MacMunn [1914] found a somewhat similar displacement in his brown specimens but he noted a band to the left of the *D* line. This could not be found with sufficient certainty to map. The presence of this band may vary with the state of oxidation of the respiratory pigment, but all our specimens were taken from fresh sea-water and used without delay. The demonstration of reduced alkaline haematin and also of haematoporphyrin is positive, Plate III, B, spectra 2, 4, 5, and in this we confirm MacMunn's findings.

The tests have to be carried out with great care as there is present an extremely small quantity of respiratory pigment. This was further proved by testing the oxygen content of the tissues, which gave spectroscopic evidence of haemoglobin derivatives, by the Barcroft [1914] method. Only 0.25 per cent. of oxygen by volume was given off by the ferricyanide from 5 g. fresh tissue as compared with mammalian blood which gives an average of from 17 to 18 per cent. evolved oxygen.

Experiment with a young red Anemone. A young red anemone was placed in fresh sea-water and screened from white light by various coloured glasses. When screened by either red or blue it moved into the yellow or any white light portion within a few hours. This experiment was repeated many times, the animal never failing to seek yellow or white light as rapidly as movement in these animals is possible. When the anemone was left altogether in white light it would remain in the same position for days.

The following are the absorptions of light by the coloured glasses used in this experiment:

Red. Nothing in the red, a strong dark band in the yellow and a heavy screen from the green to the end of the visible spectrum.

Yellow. A faint shading in the blue end.

Blue. A strong screen over the red and orange, a fainter screen from the mid-yellow to the end of the green and nothing in the blue.

Thus with these red and blue glasses the very light that the coloured ectoderm absorbs has been withheld from the anemone, making it necessary for the animal to change its position in order to obtain the rays of light most useful to it.

Anemonia sulcata.

1. *Ether-alcohol-soluble portion of ectoderm* (free from algae). A faint yellow colour was obtained giving no marked absorption spectrum, only a faint shading in the violet region.

2. *The respiratory pigment.* Alcoholic extracts of the tentacles and mesenteries of this anemone gave a greenish-yellow liquid which showed very characteristic absorption bands. The ether extract gave almost identical bands, the only difference being a change in the intensity of the bands on either side of the *C* line. In the alcohol solution the band to the left of the *C* line, as shown in Plate IV, C, spectrum 1, is very intense. Spectra 2 and 4 (Plate IV, C), the layers of which are only $\frac{1}{20}$ the thickness of those of 1 and 3, still show this band very markedly. The positions of the bands in ether solution show a very close resemblance to those of β -chlorophyll of the green leaf. The bands on either side of the *C* line occupy the position of the maximum intensity of the chlorophyll system, while in the thin layer the band between *F* and *G* is in the position of the more refrangible band of carotin or xanthophyll. We confirm MacMunn in these findings and also that this chlorophyll-like substance comes from the symbiotic algae contained in the tentacles and mesenteries of this animal. (See Microphotograph, Plate IV, D.)

Further no haemoglobin derivatives could be demonstrated by us in the tissues and it seems conclusive that this chlorophyll is here as an optical sensitiser to supply active oxygen for the use of the anemone. An experiment which may support this is as follows. We screened an anemone from light in a glass vessel containing sea-water. It climbed up the side until it could obtain the maximum daylight available. Whether this is due to a vital

necessity of the anemone or to a stimulus produced by the heliotropic action of the algae on their own behalf are questions which would be difficult to answer. Again Geddes [1882] has shown that *Anemonia sulcata* when placed in bright sunlight evolved oxygen in quantity sufficient for determination by gas analysis apparatus.

PREVIOUS WORK.

Moseley [1887] during the "Challenger" expedition made a large series of spectroscopic examinations of the various invertebrate colours. From our present standpoint his work upon the pigments of the Coelenterata whether respiratory or otherwise is of more direct importance. He isolated from certain species of Actiniae a madder red colour. In fresh condition this colour yielded a well defined absorption spectrum. Three bands were visible, one in the green and the others toward the less refrangible end. For this pigment the name polyperthyrin was proposed. MacMunn [1914] found that this polyperthyrin was to all appearances identical with haematoporphyrin although he indicates there is only spectroscopic evidence for this similarity. Moseley's polyperthyrin is very stable even on exposure to light. It is soluble in moderately strong hydrochloric and sulphuric acids. It is insoluble in water, glycerol, alcohol and ether. The colour in strong concentration of the above acids gave a screen from the *D* line extending all the way towards the violet end. When this solution was further weakened two distinct bands appeared close to and on opposite sides of the *D* line. The shading towards the less refrangible end was also lessened. This colouring matter was precipitated on the addition of alkalis as a burnt sienna flocculent deposit which produced on redissolving in acids the three-banded spectrum of the fresh substance.

According to Moseley [1887] polyperthyrin has been found in eleven different species comprising seven genera of Coelenterata. McKendrick [1881] examined the pigments of *Cyanea* from aqueous infusions of the animal. Two bands were observed, one in the orange and the other in the red. This pigment has been termed cyanein.

In Griffiths' *Physiology of the Invertebrata* it is stated: "There is no doubt that the chief mode of respiration of Coelenterates is by means of the ectodermic lining."

MacMunn [1885] devoted much time to the examination of respiratory pigments in Actiniae. The solid portions of *A. equina* were first examined in a compressorium by the microspectroscope. A band was seen which was not unlike that of reduced haemoglobin, and two other bands were noted nearer the violet end. In brown specimens he noted that the prominent band was nearer the violet and also that in some there was a band close to the left side of the *D* line. This pigment has been named by MacMunn actiniohaematin. Actiniohaematin is insoluble in the fat solvents but dissolves in glycerol. It is extracted by treatment with alcoholic potash solution with a change in condition. The spectrum thus seen was very like that of alkaline haematin

which gives a band at *D*. When reduced with ammonium sulphide two bands were observed, being to all appearances the spectrum of haemochromogen or reduced alkaline haematin.

Again the spectrum of haematoporphyrin was obtained by digesting the ectoderm of an *Actinia* in sulphuric acid and filtering through asbestos. This was further proved by the addition of ammonia in excess forming the four-banded spectrum characteristic of alkaline haematoporphyrin.

Moseley [1873] isolated from a *Tealia crassicornis* a substance which he called actiniochrome. MacMunn has proved that Moseley's actiniochrome is not the same as actiniohaematin for the products of decomposition are entirely different; actiniochrome could not be converted to reduced alkaline haematin whereas actiniohaematin very easily could. MacMunn concludes therefore that actiniohaematin is respiratory in function and actiniochrome is ornamental.

Again MacMunn has found certain "yellow cells" (symbiotic algae) in the tentacles and mesenteries of *Bunodes ballii* which give the spectrum of a chlorophyll-like substance.

Actiniochrome was found in this species but haemochromogen could not be demonstrated. He remarks that: "The replacement of actiniohaematin by the colouring matter of these 'yellow cells' is of great interest for it appears that they replace the red pigment of other species."

In the genus *Sagartia* MacMunn found actiniohaematin in many species, also the chlorophyll-like substance (chlorofucin) in *Sagartia bellis*. The "yellow cells" which lie in the endodermal lining of the tentacles were responsible for this spectrum. Further he finds that these "yellow cells" suppressed the respiratory pigment. In *Anemonia* "yellow cells" were also found which yielded the chlorophyll-like substance.

Geddes [1882] proposes the generic name of "philozoon" for the symbiotic algae or "yellow cells." In the same paper he gives a detailed account of chlorophyll-containing animals chiefly Coelenterates. He shows that there is a less amount of oxygen evolved by imprisoned algae, i.e. the symbiotic species, than by free living individuals. He concludes that the avidity for oxygen of the animal protoplasm accounts for this difference.

CONCLUSIONS.

1. *Actinia equina*. In view of the small quantity of respiratory pigment along with the alcohol-ether-soluble substances, it is concluded that the red and brown colours of these anemones are not ornamental as MacMunn and others hold, but that they act as optical sensitisers producing active oxygen for the use of the animal through the medium of the respiratory pigment. By virtue of the red screen in the position of the maximum absorption of chlorophyll they act as chemical sensitisers. Further there is reason to believe that these animals can make use of the blue and probably the violet rays.

2. *Anemonia sulcata*. In *Anemonia* there is no haemoglobin derivative present, but the animal contains minute algae lining the inside of the tentacles

A. *Actinia equina*.

Absorption spectra of ether-alcohol soluble extract of ectoderm.

B. *Actinia equina*.

Spectra of respiratory pigment.

C. *Anemonia sulcata*.
Absorption spectra of solutions.

D. Symbiotic algae from the tentacles and mesenteries of *Anemonia sulcata*. The rod-shaped objects are diatoms. $\times 150$ diam.

and mesenteries, which contain a chlorophyll-like substance. By its close similarity to the chlorophyll of the green leaf, the action of light on this body seems to be much the same, namely of optical and chemical sensitisation. Active oxygen therefore would be released by the contained algae and used by the animal if need be.

We have to thank Professor D. Noël Paton for help and advice. The expenses were defrayed by the Medical Research Committee.

REFERENCES.

- Barcroft (1914). *The Respiratory Function of the Blood*.
Farquhar (1898). *J. Linn. Soc.*, 26.
Geddes (1882). *Proc. Roy. Soc. Edin.*, 377.
Gosse (1860). *British Sea Anemones*.
McKendrick (1881). *J. Anat. Physiol.*, 15, 261.
MacMunn (1885). *Phil. Trans.*, II. 641.
— (1914). *Spectrum Analysis applied to Biology and Medicine*, 100.
Moseley (1873). *Quart. J. Microscop. Soc.*, N.S. 13, 143.
— (1887). *Quart. J. Microscop. Soc.*, N.S. 17, 1.
Saville-Kent (1893). *The Great Barrier Reef*.
Wager (1914). *Proc. Roy. Soc.*, B, 87, 386.
Walton (1911). *J. of Marine Biol. Assoc.*, N.S. 9, 228.