LIX. INVESTIGATIONS ON THE NITROGENOUS METABOLISM OF THE HIGHER PLANTS.

PART III. THE EFFECT OF LOW-TEMPERATURE DRYING ON THE DISTRIBUTION OF NITROGEN IN THE LEAVES OF THE RUNNER BEAN.

By ALBERT CHARLES CHIBNALL.

*From the Biochemical Department, Imperial College
of Science and Technology.*

(Received June 27th, 1922.)

IN any research into the metabolic processes taking place in plants, where fresh material is used, the period during which the research can be carried out must necessarily be restricted to the appropriate season during which the plants can be grown. To overcome the difficulty, thus entailed, some workers have used leaves that had been previously dried, either at low temperatures or from 50–70°.

As far as is known, however—and this statement applies especially to those who have investigated the N and protein metabolism in plants—no one has first investigated the possible changes taking place during drying. Both protein synthesis and degradation are continually going on in the leaf cell; consequently, proteolytic enzymes must be present, and it is strange that so many workers have assumed that these would remain inactive during the process of drying, often at temperatures of 30–50°, the most favourable for enzyme action.

The present research shows that in as far as the leaves of the runner bean are concerned, considerable proteolysis takes place, with consequent increase in the simpler water-soluble nitrogenous products, chiefly ammonia (in the form of ammonium salts or as the amide N of asparagine) and monoamino acids, the former being increased from 1 % to 5 % or 6 % of the total leaf N, the amount of increase depending on the conditions of drying.

On reviewing, therefore, the work done in the past to throw light on the N metabolism in the leaf, it would appear that the predominance given to asparagine as a final product of protein metabolism is based, to a certain extent, on an erroneous conception as to the amount of it actually present in the living leaf.

Boussingault [1860] was one of the first systematically to study the asparagine content of leaves, and he emphasised that it could only be isolated

when the plants were young. His results were confirmed by Meunier [1880] and Müller [1887], who extended their researches over a wide range of plants. Emmerling [1887], who studied the N content of the fresh leaves of *Vicia Faba maj.* throughout the life history of the plant, never found that either ammonia N or asparagine N exceeded 1.25 % of the total leaf N, a result the author of the present paper confirmed in an unpublished series of experiments in 1921. The latter also [Chibnall, 1922] found less than 1 % for each of these substances during the life of *Phaseolus vulgaris* v. *multifloris*. Kosutany [1897] found a similar result for *Vitis vinifera*. Yet side by side with these results others were being published which indicated that the asparagine content of leaves could be quite high. On investigation it would appear that in all these cases the leaves had been dried before analysis. Thus Suzuki [1897] gives 5 % of asparagine in leaves of *Phaseolus vulgaris* dried in an air oven at an unstated temperature. Wassilieff [1901] found 21.46 % of the total N as asparagine in young leaves of *Lupinus alba* dried at 70°. Jodidi, Kellogg and True [1920] found in spinach leaves (age unstated) dried at atmospheric temperature for 7 days, 4.8 % of the total N as ammonia and about 12 % as "amide N of asparagine." Yet the present author found in the fresh leaves from plants three weeks old only 0.6 % as ammonia and 2.05 % as "amide N of asparagine."

From the results given later, combined with those stated above, it seems fairly certain that these high values must be attributed, at any rate in part, to proteolysis during the drying. The same remark must apply to the results obtained by Prianischnikoff [1899], who investigated the seedlings of several legumes, though he analysed the plants as a whole, and not the leaves alone.

Whilst proteolysis thus affects the water-soluble products of the leaf, it is found that the general character of the leaf proteins, as shown by Hausmann's method of analysis, undergoes but little change, if any, during the drying.

EXPERIMENTAL DETAILS.

The plant used was *Phaseolus vulgaris* v. *multifloris* (Scarlet Champion). The samples of leaves investigated were from series 8, 9, and 10, of a former research [Chibnall, 1922]. In these the total weight of leaves picked was—in series 8, 2064 g. from 12 plants; in series 9, 2424 g. from 12 plants; in series 10, 1860 g. from 8 plants. For the distribution of N in the fresh leaves only about 500 g. were required, the remainder, in each case, being air-dried in a closed oven through which a slow stream of air was drawn. Series 8 was dried at 50° in approximately 100 hours, series 9 at 40° in approximately 120 hours, and series 10 at 40° in approximately 50 hours. The dried leaves were afterwards ground to a moderately fine powder and stored in a stoppered bottle. About six months elapsed before the present research was commenced.

Table I gives further details of the air-dried samples. As the drying took place in the dark, the leaves of series 8, which had been picked one hour before dawn, received no opportunity for photosynthetic action before death.

Table I. *Showing details of the air-dried samples used.*

Series number	Approximate age of plant above ground in weeks	Time of day when picked (G. M. T.)	Fresh weight of sample in g.	Air-dried weight of sample in g.	% of moisture in air-dried sample
8	11	2 a.m.	1365	213.0	7.65
9	11	3 p.m.	1925	335.1	10.92
10	14	9 a.m.	921	161.8	7.17

General methods of manipulation (separation of protein, soluble products and cellular matter). The method of grinding in an end-runner mill, as used with the fresh leaves [Chibnall, 1922], was found to be applicable in the case of the dried leaves. The ground up particles of cellular matter imbibe the water and swell up, so that only the very fine ones pass through the pores of the lawn when gentle pressure is applied. By using lawn of very close texture, as in the case of series 8, this can be reduced to a minimum, the percentage of N per dry-weight of the colloidal protein being but very little lower than that from the fresh leaves. Even when moderately coarse lawn is used, so as to allow the separation to be conducted in the minimum of time, the amount of carbohydrate mixed with the protein is not sufficient to interfere with the chemical analysis of the latter (*vide* series 9 in Table IV).

In some preliminary experiments the ground leaves were first shaken up with a small quantity of water and allowed to stand for some hours, with the object of facilitating the extraction of the less soluble products, but subsequent analysis of the extracts, as will be shown later, indicates the presence of proteolytic enzymes, which are activated by the addition of water. It is therefore necessary to carry out the extraction in the minimum of time.

In the preparations from 50 g. of the dried leaves of series 8 and 9, upon which the subsequent analyses given herein are based, 6 extractions by grinding, followed by six extractions with boiling water were given. Extraction with boiling water without the preliminary grinding in the cold was also tried. The protein- and proteose-free water-soluble N agreed with that given by the other method, but only a part of the proteoses passed into solution and, of course, no colloidal protein was obtained. On account of lack of material, the analysis given for series 10 in Table IV was of an extract prepared by this method. Table II illustrates the difference between the extracts prepared by the two methods.

Table II. *Showing the water-soluble N extracted from the dried leaves by cold and hot extraction.*

(In percentages of total leaf N.)

Series number	Number of extractions		Total water-sol. N	Precipitated by ammonium sulphate		Protein- and proteose-free water-soluble N
	By cold grinding	With hot water		Not resolvable in water (protein?)	Proteose N	
{ 8	6	6	30.78	0.51	3.62	26.65
{ 8	0	10	27.65	0.48	1.45	25.71
{ 9	6	6	33.68	0.39	3.74	29.55
{ 9	0	10	30.94	0.28	0.69	29.97

Analytical methods. The details given in the former paper were followed, with the following exceptions:

Proteose N. This was estimated by ammonium sulphate instead of zinc sulphate in acid solution [Baumann and Bömer, 1898]. No difficulty had been experienced with the latter method when using the extracts from the fresh leaves, but in those from the dried leaves the zinc sulphate separates a quantity of non-nitrogenous material of a slimy nature, which renders the subsequent filtering slow, and the washing of the separated proteoses almost impossible. No difficulty was experienced with the ammonium sulphate, the proteoses filtering readily at the pump. They were then washed with a saturated solution of ammonium sulphate until the washings were colourless and redissolved in water. The operation was repeated, and the N of the aqueous solution of the proteoses, after expulsion of the free ammonia by distillation with magnesia *in vacuo*, was determined by Kjeldahl's method. The N of the small insoluble residue on the filter paper, probably coagulated protein, was also determined, after thorough washing, by Kjeldahl's method.

Residue after extraction of the colloidal protein and water-soluble products. As this still contained about half the protein N of the leaf, three extractions with 700 cc. of 1 % HCl were given, the extracts being united and analysed as before.

RESULTS.

Effect of drying on the proteins. Table III gives a detailed account of the proteins extracted from the dried leaves of series 9, with the corresponding data for the fresh leaves. The amount of protein passing into colloidal solution is proportionately much smaller than with the fresh leaves, but could no doubt be increased by further grindings, since the 6th extract appeared as heavily charged with colloidal matter as the earlier ones. The bulk of the protein not passing into colloidal solution, however, is extracted by the HCl. Assuming, as in the case of the fresh leaves, that the small amount of N in the final residue of cellular matter is due to unextracted protein, it will be seen that the total protein N has fallen from 74.83 % to 61.15 % of the total leaf N.

Table III. *Comparison of the proteins and water-soluble N in the fresh and dried leaves of series 9.*

(In percentages of total leaf N.)

	Fresh leaves	Dried leaves
Colloidal protein extracted	59.70	29.44
Protein extracted by 1 % HCl	10.24	25.00
Remaining in residue	4.13	5.90
Coagulated by boiling	1.76	0.42
Coagulated by saturation with ammonium sulphate (?)	0.00	0.39
Total protein N	74.83	61.15
Proteose N	4.35	3.74
Protein- and proteose-free water-soluble N	17.22	29.55
Extracted by alcohol and ether	3.60	2.96

Table IV gives the distribution of N in the colloidal proteins from the fresh and dried leaves of series 8 and 9. It will be observed that no apparent change has taken place in the protein molecule.

Table IV. *Comparison of the distribution of N in the colloidal proteins from the fresh and dried leaves.*

Series number	% of N per dry weight	(In percentages of total protein N.)				
		Amide N	Humin N	Basic N	Monoamino N	Other N
8 fresh	11.75	6.14	3.55	19.33	59.25	11.73
8 dry	10.11	6.26	4.63	21.20	57.93	9.98
9 fresh	11.71	6.34	3.64	22.00	58.88	9.14
9 dry	8.64	6.73	4.71	21.67	56.30	10.51

Table V gives the distribution of N in the proteins extracted by the HCl from the fresh and dried leaves of series 9. In comparing them it must be remembered that they represent very different proportions of the total leaf protein. The HCl extract from the fresh residues represents only about $\frac{1}{6}$ of the total protein, that from the dried leaves nearly $\frac{1}{2}$. Since the leaf cell undoubtedly contains more than one protein, it follows that if the dried leaves, like the fresh, contain a small amount of protein rich in amide and basic N, these characteristics, on analysis, may be masked by the predominance of unextracted protein, of the type represented by the colloidal protein. If allowance be made for soluble carbohydrates due to the mild hydrolytic action of the HCl on the pentosans etc. of the cell wall material, which will increase the humin N at the expense of the monoamino N, the distribution of N in the HCl extract from the dried leaves does not differ from that of the corresponding colloidal protein, indicating that the major part of it is of this type. Interpreting the various distributions of N in terms of the amounts of the total leaf protein that they represent, it would appear that the leaf proteins do not undergo any appreciable change in character during the process of drying.

Table V. *Comparison of the distribution of N in the proteins extracted by 1 % HCl.*

Series number	(In percentages of total N in the extract.)				
	Amide N	Humin N	Basic N	Monoamino N	Other N
9 fresh	8.19	5.82	24.66	48.78	12.64
9 dry	6.96	6.16	21.57	54.01	11.30

Effect of drying on the proteoses. Table VI shows that in the cold extracts prepared from the dried leaves the proteoses have diminished. Since Table II shows that they are but slowly soluble in hot water, it may well be that after drying they are not so readily soluble in cold water, in which case the apparent loss may be due to incomplete extraction.

Table VI. *Comparison of the distribution of N in the aqueous extracts from the fresh and dried leaves.*

(In percentages of total leaf N.)

Series number	Total water-soluble N	Protein- and protease-free water-soluble N	Ammonia N	Amide N of Asparagine	Nitric N	Humin N	Basic N	Mono-amino N	Other N
8 fresh	21.07	*	0.36	0.0	3.56	1.00	*	7.06	2.34
8 dry	30.78	26.65	6.02	0.0	3.57	1.15	1.81	10.55	3.55
Diff.	9.71	—	5.66	—	—	—	—	3.49	1.21
9 fresh	21.57	17.22	0.44	0.53	2.73	0.99	2.00	7.76	2.77
9 dry	33.68	29.55	5.32	1.51	1.62	0.82	2.84	13.22	4.22
Diff.	12.11	12.33	4.88	0.98	1.11	—	0.84	5.46	1.45
10 fresh	22.75	18.16	0.37	0.47	3.53	1.06	2.30	6.81	3.62
10 dry	(1)	27.07	2.68	2.26	3.01	0.89	3.51	10.60	4.12
Diff.	—	8.91	2.31	1.79	0.52	—	1.21	3.79	0.50

* Proteoses not determined.

(1) Determined from an extract made with hot water only, which gives a low value for the proteoses.

Effect of drying on the protein- and protease-free water-soluble products. Table VI shows that in all three cases this has increased, the amount being roughly proportional to the time of drying. The products of the protein autolysis fall chiefly into ammonia (as ammonium salts or "amide N of asparagine") and monoamino acids. This is illustrated better by Table VII. Leaving for a moment the relation between the ammonium salts and asparagine, which is discussed later under the heading of enzymes, it will be seen that in the two samples picked after some hours' daylight, series 9 and 10, the amount of ammonia N is nearly equal to the monoamino N, whereas in series 8, picked after 6 hours' dark, the ammonia is in considerable excess. In the latter case this may possibly be due to the higher temperature of drying (50° instead of 40°) or, more probably, to the fact that the leaves, before drying, were deprived of most of their reserve carbohydrate by translocation away from the leaf at night.

Table VII. *Showing the ammonia and amino acids formed by proteolysis during drying.*

(In percentages of total autolysed protein N.)

Series number	Ammonia N + amide N of asparagine	Monoamino N	Total:	{ Ammonia N Amide N of asparagine Monoamino N
8	58.3	35.9		94.2
9	47.5	44.3		91.8
10	46.0	42.5		88.5

In series 9 and 10 again, there is a loss of nitric N, probably due to the necessary carbohydrate for the synthesis of organic nitrogenous products being present, since in series 8 there is no loss. The higher temperature

of drying may have accounted for this, however, since Couperot [1909] has shown that leaves dried at room temperature lost 25-50 % of their nitric N as hydrocyanic acid whereas those dried rapidly at 60° suffered no such loss.

Table VIII. *Showing the enzyme action produced by allowing the dried leaves to stand with cold water.*

(In percentages of total leaf N.)

Series No.	Hours standing with water	Total water-soluble N	Protein and protease N removed by ammonium sulphate	Protein- and protease-free water-soluble N	Ammonia N	Amide N of asparagine	Nitric N	Humin N	Basic N	Monoamino N	Other N
8	0	30.78	4.13	26.65	6.02	0.0	3.57	1.15	1.81	10.55	3.55
8	16	35.90	3.16	31.74	2.30	2.96	3.59	0.89	3.76	13.56	5.68
9	0	33.68	4.13	29.55	5.32	1.51	1.62	0.82	2.84	13.22	4.22
9	16	39.35	3.39	35.94	1.69	4.58	1.67	0.90	5.63	15.80	5.67
9	116	39.82	2.86	36.96	1.20	4.10	1.56	1.04	5.67	17.48	5.91
10	0	*	*	27.07	2.68	2.26	3.01	0.89	3.51	10.60	4.12
10	16	31.60	3.21	28.38	1.19	3.57	3.04	1.02	3.33	11.14	5.09

* Determined in a hot water extract only, which gives a low value for the proteoses.

Enzymes present in the dried leaves. As was stated earlier, if the ground leaves are allowed to remain suspended in water for some hours (under a layer of toluene), proteolytic enzymes become active, with consequent increase of the simpler nitrogenous products in solution. Table VIII shows the analyses that have been made to illustrate this. A study of the figures shows that the action is complex, but the following points stand out fairly clearly:

(1) There is at first a rapid increase in the water-soluble N tending to reach a limit with lapse of time.

(2) Monoamino N shows a continuous increase.

(3) Basic N shows an increase that is not maintained with lapse of time.

(4) Ammonia N shows at first a rapid decrease, tending to reach a limit. At the same time there is a nearly corresponding increase in the "amide N of asparagine."

(5) The sum ammonia N + "amide N of asparagine" shows a slow decrease, indicating re-synthesis of more complex bodies.

From (2) and (3) it is clear that an enzyme of the nature of a pepsin is present (the extract is slightly acid). But the chief point of interest is the rapid conversion of the ammonia N into the amide N of asparagine. In the leaf the origin and relationship between these two substances have not yet been definitely established. Under normal conditions the evidence is that the concentration of both of them is low. Whether the ammonia is due only to direct translocation from the root, or to protein degradation in the leaf itself or to both of these is not yet clear. Undoubtedly an enzyme capable of breaking down the protein into ammonia and amino acids exists in the dried leaves, but this may not be so in the normal fresh leaf. In a previous paper [Chibnall, 1922] there was described an experiment in which the living leaf

was starved under conditions that precluded the translocation of the products of metabolism, and in spite of considerable protein degradation, no appreciable increase in either ammonia or "amide N of asparagine" was noted¹. Suzuki [1897], by starving moist leaves for 48 hours, claims to have increased the "asparagine N" from 5 % to 13 % of the total N, but his results must be queried since he dried the leaves before analysis.

Asparagine itself cannot be a primary product of protein degradation, but must be a product of re-synthesis from the ammonia and amino acids. In very young plants direct translocation of it from the cotyledon reserve is possible, since Prianischnikoff [1896] showed in germinating seeds of *Vicia sativa* a steady increase of asparagine at the expense of the protein. Butkevitch [1909], who experimented with seedlings of lupins and peas, considers, as does Suzuki [1894], that the formation of asparagine is to remove an injurious excess of free ammonia. This view, if modified to include an excess of monoamino-dicarboxylic acids also, is very likely, and the reaction "ammonia + aspartic acid $\rightleftharpoons$ asparagine" is probably due to an enzyme governed by the H ion concentration of the medium in which it is working. As far as is known, an enzyme of this nature, an asparaginase, has not yet been indicated, and it is therefore of great interest to observe that one appears to be present in the dried leaves used in this research. Furthermore, under the conditions of the present experiment, it appears to have marked synthetic activity.

It is hoped at an appropriate season to repeat these enzyme experiments with fresh leaves, and if an asparaginase should be indicated, to study the effect of change in the H ion concentration upon its action.

In conclusion the author would like to thank Prof. S. B. Schryver for his advice and encouragement throughout the research, which was made possible by a grant from the Department of Scientific and Industrial Research.

SUMMARY.

1. The effect of low-temperature drying on the nitrogenous bodies in the leaves of the runner bean has been investigated.
2. Protein autolysis takes place, with increase in the simpler water-soluble N products.
3. These products are chiefly ammonium salts, asparagine and amino acids.
4. The leaf proteins, whilst they are diminished in amount, are not appreciably changed in character.
5. Proteolytic enzymes are present in the dried leaves, and are activated by the addition of water.

¹ The season 1921 was abnormal, the bean plants bearing but few pods. At the time this experiment was carried out, 23 August 1921, no pods were in process of formation. Further starvation experiments are being carried out this season, and the results so far to hand from leaves picked in August 1922, when pod formation was active, indicate that considerable quantities of asparagine are formed by degradation of the leaf protein. Ammonia N, as before, is unchanged.

6. The presence of an asparaginase, activated by the addition of water, is indicated. Under the conditions of the present research it possesses marked synthetic activity.

7. The position of ammonia and asparagine in the N metabolism of the leaf is discussed.

REFERENCES.

- Baumann and Bömer (1898). *Zeitsch. Untersuch. Nahr. u. Genussm.* **1**, 106.
 Boussingault (1860). *Agronomie, chimie agricole et physiologie*, 2nd ed.
 Butkewitsch (1909). *Biochem. Zeitsch.* **16**, 411.
 Chibnall (1922). *Biochem. J.* **16**, 344.
 Couperot (1909). *J. Pharm. Chim.* **29**, 100.
 Emmerling (1887). *Land. Vers. Stat.* **34**, 1.
 Jodidi, Kellogg and True (1920). *J. Amer. Chem. Soc.* **32**, 1061.
 Kosutany (1897). *Land. Vers. Stat.* **48**, 13.
 Meunier (1880). *Ann. Agron.* **6**, 275.
 Müller (1887). *Land. Vers. Stat.* **33**, 326.
 Prianischnikoff (1896). *Land. Vers. Stat.* **46**, 459.
 — (1899). *Land. Vers. Stat.* **52**, 137, 347.
 Suzuki (1894). *Bull. Coll. Agr. Tokyo*, **2**, 409.
 — (1897). *Bull. Coll. Agr. Tokyo*, **3**, 245.
 Wassilieff (1901). *Land. Vers. Stat.* **55**, 45.