VII. THE PRESENCE OF VITAMIN A IN YEAST FAT.

BY ETHEL MARJORIE LUCE AND IDA SMEDLEY MACLEAN.

*From the Departments of Experimental Pathology
and Biochemistry, Lister Institute.*

(Received December 12th, 1924.)

It is well known that yeast is comparatively rich in vitamin B and is frequently used as a source of this substance. The evidence as to the existence of the fat-soluble A vitamin in yeast is however not so clear and the presence of this accessory factor in yeast does not seem to be generally recognised.

Hopkins [1912], in his classic paper on the importance of accessory factors in normal dietaries, described experiments which certainly indicate the presence of the A vitamin in yeast. In these experiments rats were fed on a basal diet consisting of casein, starch, sugar, lard and salts; the addition of 4 % of milk to this dietary induced normal and continued growth. Protein-free alcoholic extracts of milk-solids and of yeast in very small amounts produced a similar acceleration of growth; it is also stated in the same paper that a minute amount of the ether-soluble fraction of the total alcohol extract of yeast induced great acceleration of growth. The significance of this observation seems to have escaped general notice. The diet used was deficient in the three accessory factors A, B and C, and the addition of yeast to the diet would certainly furnish an adequate supply of the B vitamin. It is improbable however that the ether-soluble fraction of the alcohol extract of yeast contained any significant amount of the B vitamin since as a general rule the B vitamin is not extracted by ether [Medical Research Committee, 1924]. It seems almost certain therefore that the acceleration of growth obtained in these experiments was caused by the addition of the ether-soluble A vitamin. Macallum [1919] also found that the addition of 2 or 6 % of dried brewer's yeast to a diet deficient in vitamins A and B induced satisfactory growth in rats for short periods of time.

It seemed to us desirable to investigate directly whether acceleration of growth in rats is produced by the addition of yeast fat (or more accurately, of the ether-soluble constituents of yeast) to a diet which already contained a supply of the B vitamin. In this case if the yeast fat was effective in inducing growth, the presence of the A vitamin in yeast would be satisfactorily confirmed.

A study of yeast fat by one of us [Smedley MacLean, 1922] has shown that the percentage of fat in yeast may be considerably increased by incubating the yeast in a solution of sugar, such as fructose, glucose or sucrose. It is

difficult however to obtain the increased amount of fat by directly extracting the dried yeast with ether; on the other hand, it is quite easily extracted if the yeast is first boiled for two hours with a normal solution of hydrochloric acid. The carbohydrate of the yeast is hydrolysed and both carbohydrate and protein pass into the acid solution, the fat being readily extracted from the dried residue. A comparison of the A vitamin content of the yeast fat obtained by direct extraction with that of the fat obtained from the yeast residue after hydrolysis of the carbohydrate was undertaken. It seemed of interest to determine whether the conditions which led to an increase of fat in the yeast cell also led to an increase of the A vitamin.

Almost all the additional fat formed when the yeast is incubated in a sugar solution is contained in the fat extracted after treatment of the yeast with the normal acid solution; it was therefore interesting to determine whether such a sample of fat contained the A vitamin or whether this vitamin was contained almost entirely in the first direct ether extract.

The experiments were carried out according to the method recommended by Zilva and Miura [1921] and the basal diet was that used by these observers containing the B and C vitamins but free from A vitamin. It consisted of a mixture of casein 20 %, starch 50 %, salts 5 %, hardened cottonseed oil 15 %, marmite 5 %, and decitrated lemon juice 5 %. After the usual preliminary period on the basal purified diet, indicated by dotted lines in the growth curves (Figs. 1, 2 and 3), the various substances to be tested were given to the animals daily for 28 days. The calcium content of the bones of both hind legs (femur, tibia and fibula) was estimated in each case by Aron and Sebaauer's method [1910].

The results of these experiments are shown in Figs. 1, 2 and 3. The rats in each group are of the same sex and taken from the same family.

Fig. 1. A comparison of growth obtained with ether extracted fat, dried yeast, incubated yeast, and maximal doses of cod-liver oil.

% Ca in dry weight of bones	
R 707	16.2
R 705	16.1
R 706	15.8
R 710	19.4
R 709	19.1
R 708	18.1

In Fig. 1 comparison is shown of the growth obtained with daily doses (a) of fat obtained by directly extracting dried yeast with ether (0.14 g.); (b) of dried yeast (0.5 g.); (c) of yeast incubated in carbohydrate solution to

increase the fat content and then dried (0.5 g.); and (d) of cod-liver oil (5 drops daily, about 0.14 g.). The results show absence of growth obtained with both specimens of dried yeast, positive growth with the yeast fat and with the cod-liver oil.

In Fig. 2 the results are given of the tests with the fat obtained by direct extraction of the dried yeast with ether and with the fat obtained from the residue by boiling it with acid and then drying it and extracting it with ether. In the first case there was growth, in the latter not.

Fig. 2. A comparison of growth obtained with ether-extracted and hydrolysed fat from the same yeast.

% Ca in dry weight of bones

R 784	12.6	R 793	14.7
R 789	14.2	R 788	16.3

Fig. 3. Test for the destruction of the growth factor in cod-liver oil by boiling with normal acid (minimal dose).

R 991	17.8	R 994	18.7
R 993	16.1	R 995	21.9

As far as these experiments go they indicate that the A vitamin is readily extracted by ether with the more easily obtained fat and there is no evidence that the residual fat is associated with the A vitamin.

Tests were then made to determine how far the process of hydrolysis used had a destructive effect on the A vitamin. A sample of cod-liver oil was submitted to the same process of boiling with normal acid and the effect of the oil thus treated compared with that of the original sample. If a dose of 5 drops of each of the two samples was given daily no difference could be detected in their results (Fig. 1), but with a minimal dose the effect becomes apparent. Fig. 3 shows that with a minimal dose of $\frac{1}{10}$ of a drop of oil there was some destruction of the growth factor in the sample that had been heated with the acid. The growth of rat 994 ran parallel with that of 995 until the dose was reduced from $\frac{1}{5}$ to $\frac{1}{10}$ of a drop.

DISCUSSION OF RESULTS.

The negative results obtained when dried yeast was added to the basal diet (Fig. 1) are probably explained by the dosage. Dried brewer's yeast used in these experiments contains about 3 % fat and 0.5 g. dried yeast contains therefore only about 0.015 g. of fat, a quantity insufficient to maintain growth; the dried yeast after incubation contains three or four times as much fat as the original sample but we have been unable to obtain evidence that there has been any corresponding increase in the growth factor. It was difficult to feed large doses of the dried yeast and the dosage is more satisfactorily tested by giving the ether-soluble constituents; the results obtained when a dose of 0.15 g. of the yeast fat was administered daily (Fig. 1) show that the growth factor is present in yeast; an increased calcium content is also found on analysis of the bones.

The striking difference in the results obtained when (a) the fat directly extracted by ether and (b) the fat dissolved by ether after the residue has been treated with acid shows that the A vitamin probably occurs in the cell in an easily extractable form and is not associated with the residual fat which is more firmly retained in the cell; we did not obtain any indication that increase of vitamin accompanied the increase of fat which occurs when the yeast is incubated in a sugar solution.

The yeast used in these experiments was ordinary brewery yeast grown in the absence of direct sunlight, the surface of the wort only being exposed to diffuse light. The question arises as to whether the yeast was able to synthesise the fat-soluble vitamin in the absence of any direct solar radiation or whether this accessory factor was picked up by the yeast from the medium and concentrated in the yeast.

Since the wort in which the yeast is grown consists essentially of an aqueous extract of malted barley with or without the addition of carbohydrate, if the fat-soluble vitamin has not been synthesised by the yeast it must have been preformed in the malted barley.

Steenbock, Kent and Gross [1918] have shown that whereas barley is rich in the water-soluble B vitamin it is deficient in the fat-soluble factor. They

found that in feeding experiments where all other requirements had been provided for, the diet of rats must contain 60 % of barley in order to get improved growth with a diet otherwise deficient in the fat-soluble factor.

Southgate [1924], who examined barley, kilned malt, unfinned and finned beers for the A vitamin was unable to detect it in any of the materials examined.

From what is known of the solubility of the fat-soluble vitamins it is improbable that the process of macerating the malted grain with water would result in the extraction of any appreciable amount of them. After boiling the wort the almost clear liquid is drawn off and as large a surface as possible of the hot liquid exposed to the air so that efficient aeration may take place, the aeration being continued during the process of cooling. Great importance is attached by the brewer to the thorough aeration of the wort and there is no doubt that this process would be highly detrimental to any A vitamin present in the original liquid.

It seems more probable therefore that the A vitamin is synthesised directly by the yeast cell and that this synthesis is brought about in the absence of any direct solar radiation. This result, if definitely established, would be of interest since, in so many instances, solar energy has been shown to play an important part in the production of substances rich in the fat-soluble vitamin.

Ehrlich [1907] showed that the higher alcohols, which are derived from the amino acids of the nutrient protein, are only formed when carbohydrate fermentation is taking place; it is possible that the energy set free by the fermentation of the carbohydrate enables the yeast cell to carry out various metabolic processes, not the least important of which may be the synthesis of the fat-soluble vitamin.

We desire to express our indebtedness to the Food Investigation Board for a grant which has enabled this investigation to be carried out; to thank Miss D. G. Hoffert for assisting us in the work and to acknowledge the courtesy of Messrs Watney Coombe and Reid in supplying us with material.

REFERENCES.

- Aron and Sebauer (1910). *Abderhalden's Handb. der biochem. Arbeitsmethoden*.
Ehrlich (1907). *Ber. deutsch. chem. Ges.* **40**, 1027.
Hopkins (1912). *J. Physiol.* **44**, 425.
Macallum (1919). *Trans. Roy. Canad. Inst.* **12**, 175.
Medical Research Council (1924). *Spec. Rep. Series* No. 38 (revised).
Smedley MacLean (1922). *Biochem. J.* **16**, 370.
Southgate (1924). *Biochem. J.* **18**, 769.
Steenbock, Kent and Gross (1918). *J. Biol. Chem.* **35**, 61.
Zilva and Miura (1921). *Biochem. J.* **15**, 654.