VI. INORGANIC CONSTITUENTS OF CEREBROSPINAL FLUID.

II. THE ULTRAFILTRATION OF CALCIUM AND MAGNESIUM FROM HUMAN SERA.

BY ELSIE WATCHORN AND ROBERT ALEXANDER McCANCE.

*From the Biochemical Laboratory, Cambridge, and the Biochemical
Department, King's College Hospital.*

(Received December 21st, 1931.)

In a recent paper McCance and Watchorn [1930-31] reported a preliminary investigation of the ultrafiltration of magnesium from serum. A further sixty cases have now been examined, including those with normal, subnormal and increased levels of serum-magnesium. No attempt has been made to study specifically any particular disease, as the main object has been to investigate the ultrafiltration of magnesium over a wide range of naturally occurring serum values. It is therefore impossible to deal statistically with separate diseases although some reference will be made to one or two, particularly nephritis. Normal pregnancy also will be discussed. As in the preliminary paper, calcium has been determined simultaneously in most cases for comparative purposes, although ultrafiltration of calcium from serum has already been extensively studied [for references see McCance and Watchorn, 1930-31].

Sources of material and technique.

The "normal" subjects were students, doctors or nurses, and were all in good health, leading active lives at the time the samples of blood were taken. All the other specimens of blood were taken from in- or out-patients of King's College Hospital. One of us [Watchorn, 1929] has shown that there is no appreciable variation in calcium and magnesium throughout the day, provided that violent exercise is not taken; consequently it is not necessary to ensure that the subject is fasting when blood is taken.

Experimental technique has been practically the same as that previously described [McCance and Watchorn, 1930-31]. To show that by this technique the ultrafiltrable calcium and magnesium can be accurately and consistently measured, a series of eight consecutive determinations is given in Table I. These were made on pooled samples of sera and each ultrafiltration was made with a separate membrane. It will be seen that the average deviations of the serum determinations are less than 1 %, and of the ultrafiltrates less than 2 %.

With regard to the ratio ultrafiltrable/total Mg or Ca it will be evident that the maximum possible variation for the series will extend from the lowest ultrafiltrable/highest total serum value to the highest ultrafiltrable/lowest total serum value. For magnesium this range is 0.732 to 0.779, and for calcium 0.533 to 0.561, indicating a possible maximum error of 6 % for the former and 5 % for the latter. The probable error is obviously less.

Table I. *Results on pooled sera.*

Magnesium (mg./100 cc.)		Calcium (mg./100 cc.)	
Serum	Ultrafiltrate	Serum	Ultrafiltrate
2.65	2.01	10.29	5.68
2.65	1.99	10.49	5.78
2.62	2.04	10.29	5.59
2.62	2.01	10.29	5.78
2.65	2.07	10.29	5.78
2.69	2.04	10.29	5.68
2.65	1.97	10.39	5.78
2.66	1.97	10.29	5.59
Mean	2.65 $\pm$ 0.014	10.33 $\pm$ 0.06	5.71 $\pm$ 0.07

It is evident that the individual variations recorded have a real meaning and are not merely technical in origin. The results therefore can be dealt with collectively and, when occasion demands, individually.

EXPERIMENTAL RESULTS.

Table II contains the results obtained from eight normal persons; it will be seen that their calcium, magnesium and protein values conform closely to

Table II. *Normal subjects.*

		Magnesium mg./100 cc.			Calcium mg./100 cc.			Protein %	
Case No.	Sex	Serum (S.)	Ultrafiltrate (U.)	U./S. ratio	Serum (S.)	Ultrafiltrate (U.)	U./S. ratio	Total	Albumin
1	F.	2.45	1.73	0.706	—	—	—	7.50	—
2	F.	2.47	1.83	0.741	10.58	4.80	0.454	6.00	4.80
3	F.	2.49	1.88	0.755	10.29	5.10	0.496	7.70	5.10
4	F.	2.50	1.60	0.640	—	—	—	7.90	5.35
5	M.	2.42	1.85	0.764	10.88	5.40	0.496	7.35	4.70
6	M.	2.48	1.90	0.766	10.39	4.85	0.467	7.70	5.40
7	M.	2.48	1.85	0.746	9.99	5.40	0.541	6.65	4.95
8	M.	2.59	1.78	0.687	9.80	5.84	0.596	6.70	—

the generally accepted normal values. Table III contains all the other results arranged in ascending order of the serum-magnesium, grouped as follows.

(a) Values below 2.3 mg./100 cc., *i.e.* definitely subnormal.

(b) Values over 2.3 and below 2.66 mg./100 cc., *i.e.* covering the widest possible limits of normality.

Table III. *Pathological cases.*

Case No.	Diagnosis	Magnesium			Calcium			Protein %	
		mg./100 cc.		U./S. ratio	mg./100 cc.		U./S. ratio	Total	Albumin
		Serum	Ultra-filtrate		Serum	Ultra-filtrate			
(a) Serum values under 2.30 mg. Mg/100 cc.									
9	Uraemia	1.57	1.29	0.822	6.86	4.51	0.657	7.70	—
10	Pregnancy	1.97	1.57	0.797	9.11	4.80	0.527	6.80	4.95
11	Enlarged prostate	2.01	1.82	0.905	—	—	—	8.50	4.45
12	Pregnancy	2.09	1.68	0.804	9.31	4.80	0.516	5.25	3.65
13	"	2.10	1.70	0.809	9.19	6.62	0.720	7.35	4.70
14	"	2.17	1.82	0.839	9.02	5.59	0.620	5.60	3.74
15	"	2.19	1.61	0.735	8.59	6.12	0.712	7.00	3.40
16	Before overbreathing	2.20	1.81	0.823	10.29	4.90	0.476	6.31	4.45
17	Hyperthyroidism	2.21	1.45	0.656	9.90	5.54	0.560	7.00	3.80
18	Enlarged prostate	2.24	1.87	0.835	12.08	5.54	0.459	6.30	—
19	Pregnancy	2.25	1.66	0.740	7.74	5.19	0.671	6.30	3.95
20	"	2.26	1.68	0.743	8.50	5.33	0.627	6.30	—
21	After overbreathing	2.28	1.44	0.632	11.46	6.62	0.578	7.55	5.20
(b) Serum values 2.30–2.65 mg. Mg/100 cc.									
22	Before overbreathing	2.34	1.57	0.671	10.96	5.64	0.514	—	—
23	Chronic parenchymatous nephritis	2.36	1.77	0.750	8.61	5.64	0.655	4.20	2.80
24	Pregnancy	2.47	2.11	0.854	8.92	6.17	0.692	6.15	4.20
25	"	2.49	2.00	0.803	9.58	5.93	0.618	8.50	4.90
26	Lymphadenoma	2.51	1.91	0.761	9.26	6.06	0.654	4.20	3.50
27	After parathyroid therapy	2.52	1.81	0.718	10.59	4.85	0.458	—	—
28	Pregnancy	2.54	1.93	0.760	8.00	5.83	0.729	6.30	—
29	Diabetes	2.54	2.00	0.787	12.60	8.13	0.645	7.00	4.00
30	?	2.63	1.85	0.720	9.90	5.30	0.535	7.00	—
31	Pulmonary T.B.	2.64	1.79	0.678	10.64	5.00	0.470	5.97	4.65
(c) Serum values 2.66–3.00 mg. Mg/100 cc.									
32	Acute anterior poliomyelitis	2.66	1.84	0.692	9.89	5.64	0.570	7.00	4.45
33	Pregnancy	2.70	2.04	0.756	9.58	6.81	0.711	8.05	5.80
34	After parathyroid therapy	2.70	1.87	0.693	9.40	5.00	0.532	7.50	—
35	Chronic parenchymatous nephritis	2.71	2.48	0.915	9.70	4.21	0.434	4.00	2.50
36	Enlarged prostate	2.72	2.04	0.750	12.45	7.55	0.606	7.15	4.45
37	Eclampsia	2.73	1.98	0.725	9.09	5.33	0.485	6.30	3.40
38	Nephritis	2.73	2.02	0.740	10.99	4.26	0.469	7.00	—
39	Uraemia	2.77	1.92	0.693	7.12	5.33	0.748	6.50	4.80
40	After overbreathing	2.77	1.74	0.628	10.49	5.19	0.495	—	—
41	Asthma	2.79	2.24	0.803	11.17	6.17	0.553	—	—
42	Lead neuritis	2.81	1.99	0.708	9.70	5.90	0.608	6.90	—
43	Uraemia	2.86	1.88	0.657	12.24	5.74	0.469	7.45	5.10
44	Chronic nephritis	2.91	2.51	0.862	9.58	5.83	0.609	7.30	4.30
45	Pregnancy and nephritis	2.96	2.70	0.912	9.70	5.70	0.588	5.70	—
46	Asthma	3.00	1.94	0.647	9.31	5.78	0.621	—	—
(d) Serum values over 3.00 mg. Mg/100 cc.									
47	Carcinoma of oesophagus	3.07	1.95	0.635	11.16	6.62	0.593	6.30	4.00
48	Asthma	3.14	2.38	0.758	11.56	5.98	0.517	—	—
49	Pulmonary T.B.	3.17	2.36	0.745	7.11	4.74	0.667	5.80	—
50	Streptococcal infection of throat	3.18	2.14	0.673	12.67	6.28	0.496	7.90	5.35
51	Uraemia	3.26	2.62	0.804	6.42	4.84	0.754	8.50	3.50
52	After intramuscular injection of magnesium	3.33	2.14	0.643	—	—	—	—	—
53	Nephritis	3.43	2.50	0.729	10.57	7.01	0.663	7.15	4.80
54	After intramuscular injection of magnesium	3.53	3.00	0.853	10.98	6.70	0.610	—	—
55	Diabetic coma	3.70	2.70	0.730	13.00	8.13	0.625	8.10	4.00
56	T.B. meningitis	3.81	3.38	0.887	12.03	6.32	0.525	—	—
57	Uraemia	3.92	3.28	0.837	3.38	2.06	0.609	6.20	3.80
58	"	3.95	2.88	0.729	8.81	5.01	0.569	6.30	4.65
59	Uraemia caused by chronic lead poisoning	5.77	4.27	0.740	13.62	7.98	0.586	6.95	5.35
60	Uraemia	10.34	7.86	0.760	3.82	2.35	0.615	6.30	4.30

(c) Values over 2.65 and below 3.00 mg./100 cc., *i.e.* definitely above normal. This class includes the great majority of hospital patients [McCance and Watchorn, 1930-31].

(d) Values over 3.00 mg./100 cc., *i.e.* wide departure from normal.

Table IV contains the averages of the normals and the four pathological groups, and also the averages within each group of the ratio of the ultrafiltrable magnesium to the total serum-magnesium (U./S. ratio). In this Table the results are also expressed as millimols per litre, correction having been made in the case of serum for the space occupied by proteins, according to the formula:

$$\frac{M \times 100}{(100 - P)}$$

where M represents millimols per litre of Mg or Ca, and P the percentage of total serum-proteins.

Table IV. *Average values.*

(I) Magnesium.

Range (mg./100 cc.)	Serum (S.)	Ultra- filtrate (U.)	U./S. ratio	Serum liquid (S.L.)	Ultra- filtrate	U./S.L. ratio	Serum- protein %
	mg./100 cc.			millimols per litre			
Normals	2.48	1.80	0.726	1.10	0.74	0.673	7.19
Group <i>a.</i> 1.57-2.28	2.11	1.65	0.782	0.93	0.68	0.731	6.77
„ <i>b.</i> 2.34-2.64	2.50	1.87	0.748	1.10	0.79	0.718	6.18
„ <i>c.</i> 2.66-3.00	2.79	2.08	0.745	1.22	0.86	0.705	6.48
„ <i>d.</i> over 3.00	3.63	2.74	0.751	1.64	1.13	0.689	7.02
Average for whole series	2.74	2.03	0.750	1.17	0.83	0.709	6.72

(II) Calcium.

Normals	10.32	5.23	0.508	2.77	1.28	0.462
Group <i>a</i>	9.34	5.46	0.584	2.50	1.36	0.544
„ <i>b</i>	9.91	5.85	0.590	2.64	1.46	0.553
„ <i>c</i>	10.03	5.63	0.561	2.68	1.41	0.526
„ <i>d</i>	9.62	5.69	0.592	2.55	1.42	0.557
Average for whole series	9.79	5.61	0.573	2.62	1.40	0.534

DISCUSSION.

The relationship of the ultrafiltrable to the total serum-magnesium and calcium.

Table IV shows that the average ratio of the ultrafiltrable/total serum-magnesium for the whole series is 0.750, and that the average ratios of the normal group and of Groups *a*, *b*, *c* and *d*, are all very close to this figure. The lowest individual figure is 0.628 and the highest 0.915, while the average deviation for the whole series is 0.058. It is evident therefore that however much magnesium there is in the serum in general the same fraction of it is ultrafiltrable, and that even individual variations are not very great. When corrections are made for the space occupied by the serum-proteins the ratio for the whole series (ultrafiltrable Mg/Mg in serum liquid) is 0.709.

The average ultrafiltrable calcium for the whole series is 5.61 mg./100 cc., or, expressed as a ratio of the total serum-calcium, 0.573. This is higher than Greenberg and Gunther's average [1930], and higher than the average ratio of our normal subjects, which is 0.508. As with magnesium the ultrafiltrable calcium tends to vary directly with the total serum-calcium. Thus, for all the serum-calcium values under 10 mg./100 cc., the average ultrafiltrable calcium was 4.97 mg./100 cc., and for serum-calcium values over 10 mg./100 cc. the ultrafiltrable value was 6.03 mg./100 cc. The corresponding ratios are 0.617 and 0.532 respectively. Case No. 60 is a good example of the constancy of the ultrafiltrable/total serum ratio (U./S. ratio) both for magnesium and calcium, even when the serum values are grossly abnormal. Thus, with a total serum-Mg of 10.34, the U./S. ratio is 0.760 and with a total serum-Ca of 3.82 the U./S. ratio is 0.615.

There is no relation between the level of calcium and of magnesium in the serum, both average values (Table IV) and individual cases show that these two blood constituents vary independently of each other.

It appears from these figures that on an average about 25 % of the total magnesium of serum and 41 % of the total calcium are non-diffusible. It is generally presumed that this non-diffusible calcium or magnesium is bound to the serum-proteins [Peters and Van Slyke, 1931, p. 811]. If this presumption is correct one would expect that the ratio of the ultrafiltrable/total serum-Mg or Ca would vary inversely with the total serum-proteins, for the smaller the protein content the more magnesium or calcium will be free to pass into the ultrafiltrate. When all the figures under review are considered together it is impossible to demonstrate this point. Taking all the available data from Tables II and III they can be arranged so that all cases with serum-proteins below 7 % form Group I and all with proteins 7 % or above Group II. The average U./S. ratios then obtained are:

Group I (serum-proteins below 7 %)	0.588 Ca U./S.	0.767 Mg U./S.
Group II (serum-proteins 7 % or above)	0.590 ,,	0.745 ,,

From these figures it appears that the amount of magnesium or calcium found in an ultrafiltrate is practically unaffected by the level of proteins in the serum. This is shown also in Table IV where considerable variations in serum-proteins are not accompanied by corresponding changes in the ultrafiltrable fraction of the serum-magnesium or -calcium. Many of these cases, however, though not showing marked abnormalities in the total serum-protein level have greatly disturbed relative amounts of albumin and globulin. When such cases are eliminated from the series (that is to say, those cases having albumin/globulin ratios outside the widest limits of normal, 1.6-4.0), then the relation between the serum-protein level and the ultrafiltrable Ca or Mg is:

Group I (serum-protein below 7 %)	0.581 Ca U./S.	0.767 Mg U./S.
Group II (serum-protein 7 % or above)	0.570 ,,	0.719 ,,

Again the lack of relation between the calcium and the serum-protein level is evident, but the results seem to indicate that in the majority of cases there is

a tendency for the ultrafiltrable magnesium to be inversely proportional to the total serum-protein. A further analysis of the figures confirms this:

Proteins below 6 %	magnesium U./S. ratio	0.797.
Proteins 6-7 %	„ „	0.752.
Proteins 7 % or over	„ „	0.718.

The fact that this relation is not universal over the whole range of cases under consideration suggests that some other factor besides the amount of protein present in the serum affects the ultrafiltrable magnesium, and may in many instances be the chief agent affecting the non-diffusible fraction. If the serum-magnesium and -calcium in colloidal form are not entirely in direct combination with the proteins they may be in part in combination with the organic lipid phosphoric acids of the serum. The salts of these acids would resemble alkali protein salts in behaviour. These acids have the requisite properties therefore and are present in sufficient quantity to bind considerable amounts of Mg and Ca in colloidal form. Presumably also these elements can vary independently of the proteins [Peters and Van Slyke, 1931, p. 1102]. Both Brull [1928] and Scholtz [1931, 2] believe that calcium and magnesium form colloidal phosphate complexes. This point is being further studied.

Though most writers have suggested that the serum-calcium level is influenced by the percentage of protein present, Stearns and Knowlton [1931] failed to find any relation between these two constituents in the sera of normal children.

Pregnancy.

Twelve cases were examined between the 32nd and 38th week of gestation. Their Case Numbers are 10, 12, 13, 14, 15, 19, 20, 24, 25, 28, 33, 45. Four points require discussion. (a) The total proteins in the serum average 6.60 % which is below the generally accepted normal value of 7 % [Salvesen, 1926] and below the normal average of 7.19 % found in the present study. This reduction of serum-protein during pregnancy is a recognised phenomenon. According to Plass and Bogert, Coetzee, and Oard and Peters [quoted from Peters and Van Slyke, 1931, p. 663] the serum-proteins are at their lowest level (average 6.2 %) at the sixth month of pregnancy and thereafter rise to nearly normal at term. The average of 6.6 % therefore at the 7-8th month agrees well with these observers. Kydd [1931] found that in pregnancy the serum-proteins were reduced at the expense of the albumin fraction. The present results tend to confirm this also. The albumin/globulin ratio of the small normal series is 2.3 while that of the pregnant series is 1.93. (b) The low serum-calcium found by many other workers [*e.g.* Aburel and Ornstein, 1930; Bokelmann and Bock, 1928; Plass and Bogert, 1923; Spiegler and Schol, 1929; Widdows, 1923] is here shown again. All the values are below 10 mg./100 cc. and 5 are below 9 mg./100 cc. (c) In 7 cases the serum-magnesium is also definitely low, in 3 it is normal and in 2 raised. One of the last was complicated by nephritis and therefore should probably be excluded from consideration here. As it is

unusual to find a serum-magnesium below normal, most known pathological processes tending to give a high figure, it is evident that pregnancy tends to lower the serum-magnesium as well as the serum-calcium. This generalisation cannot be too strictly applied to the individual cases where, for example, a low magnesium may be associated with a normal calcium and *vice versa*. (d) The ultrafiltrable fraction of the total serum-magnesium was increased. The average ultrafiltrable/total Mg (U./S.) ratio is 0.796, compared with a ratio of 0.726 for the normal cases, and 0.750 for the whole series. The ultrafiltrable fraction of the calcium is also increased, the average ratio being 0.643, against a ratio of 0.508 for the small normal series, and 0.573 for the whole series. Although the number of pregnant cases is rather small (12), and the results must be confirmed on a larger series suitably controlled, they suggest that during pregnancy a larger amount of the magnesium and calcium is ultrafiltrable than in normal conditions. This is not satisfactorily explained by the diminished quantities of protein in the serum during pregnancy, because (i) the massed results of the present study have failed to demonstrate any such dependence (*v. supra*); (ii) individual cases, for instance Nos. 13 and 25, may show a high proportion of ultrafiltrable Mg or Ca associated with a high percentage of serum-protein. Factors of which we have no certain knowledge are evidently at work in these cases. We can only say that some change in the normal physico-chemical equilibria must have taken place. Whatever the underlying mechanism, the increasing readiness with which calcium and magnesium can be ultrafiltered from the serum during pregnancy may be of considerable biological significance, because everything that passes through the placenta must presumably be in ultrafiltrable form. The physico-chemical changes that take place in the blood during pregnancy therefore favour the passage of these two essential elements from mother to foetus.

Uraemia and nephritis.

Eight cases are available for discussion, Nos. 9, 39, 43, 51, 57, 58, 59, 60. No. 59 was the result of chronic lead poisoning. These patients all showed a high degree of N-retention. Most had blood-ureas of 200–300 mg./100 cc. and two of over 800 mg./100 cc. (Nos. 51 and 57). Remarkable variations of both serum-calcium and -magnesium have been found in this group, magnesium values ranging from 1.57 to 10.34 mg./100 cc. and calcium values from 3.38 to 12.24 mg./100 cc. These highly abnormal fluctuations have not been accompanied by corresponding protein changes and even the extreme cases have shown normal U./S. ratios both for calcium and magnesium. Whatever their meaning these results do seem consistent and definite. Salvesen and Linder [1923, 2] and Pincus, Peterson and Kramer [1926] have claimed that the low blood-calcium associated with a normal amount of filtrable calcium found by them in non-uraemic nephritis is the result of a decrease in serum-proteins, and that the absence of tetany in these cases can be thus explained. It is evident that in uraemia the low serum-calcium is not the result of lowered serum-

proteins or associated with raised U./S. ratios. Tetany has never been observed in any case in this series even when the serum-calcium was as low as 3.38 mg./100 cc.

Five cases of nephritis without uraemia have been studied (Nos. 23, 35, 38, 44 and 45). Nos. 38, 44 and 45 were of the mixed type so often seen clinically with some N-retention and some oedema. These require no comment. Nos. 23 and 35 were diagnosed clinically as chronic parenchymatous nephritis and merit further consideration. Both had extensive oedema, complete absence of N-retention and very low serum-proteins. According to Salvesen and Linder [1923, 1] and Pincus, Peterson and Kramer [1926] one would have expected to find in both a low serum-calcium and a high ultrafiltrable/total Ca ratio. On general grounds also one would have expected the ultrafiltrable magnesium to have run parallel with the ultrafiltrable calcium. The results actually obtained were:

	Serum-Ca mg./100 cc.	U./S. ratio	Serum-Mg mg./100 cc.	U./S. ratio
No. 23	8.61	0.655	2.36	0.750
No. 35	9.70	0.434	2.71	0.915
Normal	9-11	0.508	2.3-2.65	0.726

It will be noted that (a) the predicted relationship holds for the Ca of No. 23, (b) it cannot be said to apply to the Mg of the same case, (c) the serum-Ca of No. 35 is strictly normal and the U./S. ratio low, (d) the serum-Mg is over normal and its U./S. ratio very high. These individual results, which cannot be explained away on technical grounds (see Table I), are quoted to show that low serum-proteins may be associated with abnormally high or low U./S. ratios, and that the U./S. ratio of Ca and Mg in the same serum may vary very widely from the normal in different directions. Many other examples of this may be found in the Tables. As one would expect any external change of conditions such as p_H to alter the ultrafiltrability of Ca and Mg in the same direction it is probable that the cause of these differences must lie in the serum itself.

Eclampsia.

The only case studied (No. 37) showed no abnormalities. There was no reduction of the ultrafiltrable calcium such as Rodecurt, Koenig and Regensburger [1928] found, and no reduction of the total calcium such as Halverson, Mohler and Bergeim [1917] reported.

Parathyroid administration.

Only one case has been available for study. This was a man aged 59 suffering from peripheral neuritis due to lead poisoning (case Nos. 42, 34 and 27). Collip's parathormone was given intramuscularly.

During treatment the total serum-Mg fell definitely but slightly and with it the ultrafiltrable Mg, so that the ratio U./S. remained constant. The variations in the latter are within the limits of experimental error. The total

serum-Ca did not rise beyond normal limits. Hunter and Aub [1926-27] have reported that in similar cases parathyroid sometimes failed to raise the serum-calcium, even when given in massive doses. In the case under discussion there was a progressive and definite fall in the ultrafiltrable fraction of calcium, which is in harmony with the findings of Cameron and Moorhouse [1925]; Cruickshank [1923, 1, 2] and Merritt and Bauer [1931], but contrary to those

	Magnesium			Calcium		
	Serum	Ultra-filtrate	U./S. ratio	Serum	Ultra-filtrate	U./S. ratio
Before treatment	2.81	1.99	0.708	9.70	5.90	0.608
After 6 days during which 180 units parathormone were given	2.70	1.87	0.693	9.40	5.00	0.532
After 6 more days during which 375 units parathormone were given, i.e. 555 units in 12 days	2.52	1.81	0.728	10.59	4.85	0.458

of Liu [1927], and of Snell and Wales [1930]. Moritz [1925] and Reed [1928], who also investigated the effect of parathormone on the ultrafiltrable or colloidal calcium in serum, have obtained inconclusive results. Scholtz [1931, 1] found that parathormone raised the serum-magnesium for a few hours after the injection, but his investigation is not comparable with ours, for which blood was never withdrawn within 6 hours of an injection of parathormone.

The partition of magnesium and calcium between the serum and cerebrospinal fluid.

Stary, Kral and Winternitz [1929] and McCance and Watchorn [1930-31] have each concluded that there is no relation between the diffusible serum-magnesium and the cerebrospinal fluid-magnesium. The present study confirms this opinion. In the whole range of 60 cases here presented the percentage of magnesium found in the ultrafiltrate only rises as high as the average normal c.s.f. percentage in Nos. 56, 59 and 60. Further the amount of magnesium found in an ultrafiltrate depends mainly upon the amount originally present in the serum, whereas the c.s.f.-magnesium tends to be constant in amount and independent of the serum value [McCance and Watchorn, 1930-31]. Case No. 60 illustrates this. The serum-magnesium was grossly abnormal (10.34 mg./100 cc.), the ultrafiltrate contained 7.86 mg./100 cc. and the ratio of the two was 0.760, which is normal. Yet the c.s.f. of this case contained only 4.69 mg./100 cc., which although definitely above normal was not much more than half the concentration in the ultrafiltrate.

Opinion is still divided as to whether ultrafiltration will wholly explain the calcium found in the cerebrospinal fluid. There is no doubt that when a series is averaged the c.s.f.-calcium and the ultrafiltrable calcium lie very close

together, and that both tend to rise and fall with the total serum value. In individual cases however the c.s.f. and ultrafiltrate values do not agree and the differences cannot be dismissed as experimental errors [see McCance and Watchorn, 1930-31, Tables III and IV]. Morgulis and Perley [1930] state that the c.s.f.-calcium is relatively much more constant in amount than the serum-calcium, and only follows to a slight degree experimental variations in the latter. Brucke [1925] found the c.s.f.-calcium relatively independent of that of the serum. Moreover, although Critchley and O'Flynn [1924] found the c.s.f.-calcium very much reduced in tetany, Leicher [1922-23] has reported a case of tetany with a very low serum-calcium and a normal c.s.f.-calcium. Hertz [1930] also, working with dogs, came to the conclusion that in acute variations of the serum-calcium the ultrafiltrable calcium no longer corresponds with the c.s.f.-calcium. One case from the present series (No. 60) shows definitely that the c.s.f.-calcium may be entirely dissociated from the ultrafiltrable calcium of the corresponding serum. The serum-calcium of this case was only 3.8 mg./100 cc.; the ultrafiltrable calcium was 2.35 mg./100 c.c., so that the U./S. ratio (0.615) was nearly normal; yet the c.s.f.-calcium was 4.21 mg./100 cc., practically in the normal range and actually higher than the serum-calcium. It is possible therefore that agreement between the ultrafiltrable and c.s.f.-calcium over the normal range of serum values is merely fortuitous and that the c.s.f.-calcium is regulated at its level of 4-6 mg./100 cc. by some mechanism allied to that which maintains the c.s.f.-magnesium about 3.33 mg./100 cc. From this it is evident that the opinion of Merritt and Bauer [1931] that the cerebrospinal fluid-calcium content is "a more accurate physiological representation of the diffusible calcium of the serum than the values obtained by the various filtration methods" cannot be entirely accepted.

SUMMARY.

- (1) Eight normal and 52 pathological cases have been examined.
- (2) A very constant fraction (about 75 %) of the total serum-magnesium is ultrafiltrable. On an average 57 % of the serum-calcium is ultrafiltrable, so that a smaller proportion of the serum-magnesium than of the serum-calcium is present in colloidal form.
- (3) The proportion of the total serum-calcium that is ultrafiltrable is not affected by the proteins in the serum. The ultrafiltrable magnesium may be inversely proportional to the total serum-protein within certain limits. This has been discussed. In the same serum the ultrafiltrable/total serum-calcium ratio may be very high and the magnesium ratio very low.
- (4) In pregnancy the serum-magnesium as well as the serum-calcium tends to fall and a higher proportion of both tends to be ultrafiltrable.
- (5) Great variations in both serum-magnesium and calcium may be encountered in the last stages of chronic interstitial nephritis (uraemia), but the ultrafiltrable/total ratio remains within normal limits.

(6) After parathormone administration one case showed a progressive fall in the ultrafiltrable/total serum-calcium ratio.

(7) The concentration of magnesium in the cerebrospinal fluid cannot be explained by ultrafiltration. Ultrafiltration may explain the c.s.f.-calcium over normal ranges of serum-calcium values although such an explanation is not wholly satisfactory. Ultrafiltration cannot explain the c.s.f.-calcium in some cases with very low serum-calcium values.

E. W. is indebted to the Medical Research Council for a whole time personal grant. R. A. McC. has carried on this work during the tenure of a Pinsent Darwin Studentship which he acknowledges with thanks.

REFERENCES.

- Aburel and Ornstein (1930). *Compt. Rend. Soc. Biol.* **105**, 195.
 Bokelmann and Bock (1928). *Arch. Gynäk.* **133**, 308.
 Brucke (1925). *Deutsch. Arch. klin. Med.* **148**, 183.
 Brull (1928). *Arch. int. Phys.* **30**, 1.
 Cameron and Moorhouse (1925). *J. Biol. Chem.* **63**, 687.
 Critchley and O'Flynn (1924). *Brain*, **47**, 337.
 Cruickshank (1923, 1). *Biochem. J.* **17**, 13.
 — (1923, 2). *Brit. J. Exp. Path.* **4**, 213.
 Greenberg and Gunther (1930). *Arch. Int. Med.* **24**, 72.
 Halverson, Mohler and Bergeim (1917). *J. Biol. Chem.* **32**, 171.
 Hertz (1930). *Biochem. Z.* **217**, 337.
 Hunter and Aub (1926–27). *Quart. J. Med.* **20**, 123.
 Kydd (1931). *J. Biol. Chem.* **91**, 63.
 Leicher (1922–23). *Deutsch. Arch. klin. Med.* **141**, 196.
 Liu (1927). *Chinese J. Physiol.* **1**, 331.
 McCance and Watchorn (1930–31). *Quart. J. Med.* **24**, 371.
 Merritt and Bauer (1931). *J. Biol. Chem.* **90**, 233.
 Morgulis and Perley (1930). *J. Biol. Chem.* **88**, 169.
 Moritz (1925). *J. Biol. Chem.* **66**, 343.
 Peters and Van Slyke (1931). *Quantitative clinical chemistry*, 2. (Baillière, Tindall and Cox, London.)
 Pincus, Peterson and Kramer (1926). *J. Biol. Chem.* **68**, 601.
 Plass and Bogert (1923). *Amer. J. Obst. Gyn.* **6**, 427.
 Reed (1928). *J. Biol. Chem.* **77**, 547.
 Rodecurt, Koenig and Regensburger (1928). *Z. Geburt. Gynäk.* **93**, 410.
 Salvesen (1926). *Acta Med. Scand.* **65**, 147.
 — and Linder (1923, 1). *J. Biol. Chem.* **58**, 617.
 — — (1923, 2). *J. Biol. Chem.* **58**, 635.
 Scholtz (1931, 1). *Arch. exp. Path. Pharm.* **159**, 233.
 — (1931, 2). *Biochem. Z.* **231**, 135.
 Snell and Wales (1930). *Proc. Staff Mayo Clinic.* **5**, 17.
 Spiegler and Schol (1929). *Arch. Gynäk.* **135**, 461.
 Stary, Kral and Winternitz (1929). *Z. ges. exp. Med.* **66**, 671.
 Stearns and Knowlton (1931). *J. Biol. Chem.* **92**, 639.
 Watchorn (1929). Thesis (Cambridge).
 Widdows (1923). *Biochem. J.* **17**, 34.