LIII. STUDIES IN THE BIOCHEMISTRY OF MICRO-ORGANISMS.

XXIII. PUBERULIC ACID $C_8H_6O_6$ AND AN ACID $C_8H_4O_6$, NEW PRODUCTS OF THE METABOLISM OF GLUCOSE BY *PENICILLIUM PUBERULUM* BAINIER AND *PENICILLIUM AURANTIO-VIRENS* BIOURGE. WITH AN APPENDIX ON CERTAIN DIHYDROXY- BENZENEDICARBOXYLIC ACIDS.

BY JOHN HOWARD BIRKINSHAW AND HAROLD RAISTRICK.

*From the Division of Biochemistry, London School of Hygiene and
Tropical Medicine, University of London.*

(Received January 6th, 1932.)

IN a biochemical and toxicological study of *Penicillium puberulum* Bainier, one of the organisms responsible for the deterioration of maize, Alsberg and Black [1913] isolated from the products of metabolism of this mould when grown on Raulin's medium a definite crystalline substance, to which they gave the name of penicillic acid. They determined its empirical formula ($C_8H_{10}O_4$) and described its properties in detail, but did not determine its constitution. Since the authentic culture used by Alsberg and Black was available to us through the courtesy of Dr Charles Thom, of the United States Bureau of Agriculture, it was decided to investigate further this interesting organism, the morphological characteristics of which are described on p. 272 of Thom's *Penicillia* [1930].

It soon became evident that *P. puberulum* produces, in addition to penicillic acid, small quantities of a variety of other products, some of which are new, and it is our purpose here to describe and characterise one of these products, an acid having the empirical formula $C_8H_6O_6$, and yielding a deep reddish-brown colour with ferric chloride. We propose to give to this acid the name "puberulic acid."

30 litres of Czapek-Dox medium of the following composition were made up. Glucose 50.0 g.; $NaNO_3$ 2.0 g.; KH_2PO_4 1.0 g.; KCl 0.5 g.; $MgSO_4 \cdot 7H_2O$ 0.5 g.; $FeSO_4 \cdot 7H_2O$ 0.01 g.; distilled water 1000 cc.

5 litres of this medium were distributed in each of 6 silica trays contained in the steriliser-incubator described by Birkinshaw, Charles, Lilly and Raistrick [1931]. After sterilisation, the medium was sown with a spore suspension of *P. puberulum* and incubated at 24° .

Sterile air was passed continuously over the surface of the mycelium at the rate of 4.5–6 cu. ft. per hr. After 12 days' incubation, the metabolism solution, which gave a brownish-red colour with ferric chloride, was removed and filtered.

The total solution, containing 0.58 % of residual glucose, and having a volume of 23.65 litres, was evaporated *in vacuo* and filtered from a yellowish-brown substance, which separated on evaporation. This substance proved to be the magnesium salt of an acid having the empirical formula $C_8H_4O_6$, which will form the subject of a future communication. The filtrate was acidified with a known amount of sulphuric acid and extracted with ether. The ether extract deposited on cooling a certain amount of yellow material which was separated, and weighed 1.5 g. This product, which forms the subject of the present communication, will be called puberulic acid.

The ethereal mother-liquor was evaporated, the residue taken up in water, filtered from a small amount of insoluble matter, and again extracted with ether, from which crystals finally separated. This product was found to agree with that described by Alsberg and Black under the name of penicillic acid.

During the progress of this work it was observed that a brownish-red colour reaction was also obtained by the addition of ferric chloride to the metabolism solution from another mould species, *P. aurantio-virens* Biourge, when grown on Czapek-Dox glucose solution. This culture was kindly supplied by Prof. Philip Biourge of the University of Louvain. It is Biourge's culture number B. 77, and its morphological characteristics are described by him [1923, p. 119] and by Thom [1930, p. 316]. Since this organism appears to give larger yields of both puberulic acid and the acid $C_8H_4O_6$ already referred to than does *P. puberulum*, it was used for the preparation of these two substances.

As the result of a large number of experiments with Czapek-Dox media, containing varying amounts of glucose and mineral salts, the following medium, for which we are indebted to our colleague Mr George Smith, was finally chosen as giving the best yields of puberulic acid and the acid $C_8H_4O_6$.

Glucose 70.0 g.; $NaNO_3$ 15.0 g.; KH_2PO_4 1.0 g.; KCl 0.5 g.; $MgSO_4 \cdot 7H_2O$ 0.5 g.; $FeSO_4 \cdot 7H_2O$ 0.005 g.; distilled water 1000 cc.

This was distributed in 350 cc. quantities in 1 litre conical flasks, sterilised, sown with a spore suspension of *P. aurantio-virens* and incubated at 24° for 42 days.

It was necessary, however, to modify the method employed for the isolation of the acids owing to the presence in the metabolism solution of a large amount of succinic acid, which was extracted by ether from the evaporated solution along with the other acids. The method adopted depends upon the observation that both puberulic acid and the acid $C_8H_4O_6$ are precipitated as deep red insoluble nickel salts on the addition of nickel sulphate solution to the neutralised metabolism solution, whereas succinic acid remains in solution.

The filtered metabolism solution was therefore first neutralised to phenolphthalein by the addition of 2N NaOH, the amount required being determined

by titration. Puberulic acid and the acid $C_8H_4O_6$ were then precipitated by the addition of a slight excess of *M* nickel sulphate solution, the amount added being usually about equal in volume to the NaOH added. After the bright red precipitate had settled, most of the liquid was siphoned off, and the precipitate filtered and washed. It was then ground up with an excess of 2*N* hydrochloric acid. The colour now changed from a bright red to a dirty yellow—the latter representing the colour of the crude acids. The mixture was then extracted with ether, and the evaporated ether extract taken up in a little methyl alcohol and allowed to crystallise. By this means two or three crops of a bright yellow colour were obtained of a mixture of puberulic acid and the acid $C_8H_4O_6$.

These two acids are difficult to separate by fractional crystallisation, but may be satisfactorily separated by taking advantage of the fact that, on treating the mixture of crude acids with acetic anhydride and sodium acetate, the acid $C_8H_4O_6$, which does not form an acetyl compound, separates from the diluted acetylation mixture as the acid sodium salt, whereas puberulic acid forms a diacetyl compound which remains in solution and may be precipitated by acidification with a mineral acid.

In a typical experiment the total crude mixed acids from 143 flasks (50 litres of medium), weighing 16.4 g., were heated with 33 g. of anhydrous sodium acetate and 65 cc. of acetic anhydride for 20 minutes, in an oil-bath, the temperature of which was maintained at 140–150°. The still warm mixture was poured into about 300 cc. of water and stirred. The whole dissolved, giving a deep yellow-brown solution, from which the acid sodium salt of the acid $C_8H_4O_6$ rapidly began to crystallise. After standing overnight, this was filtered off and dried, and weighed 8.5 g. The filtrate was acidified with 42 cc. of 10*N* sulphuric acid. The crude diacetylpuberulic acid separated as a somewhat dark crystalline solid which weighed 7.0 g. Most of the colour disappeared on washing with a little hot absolute alcohol. The product was recrystallised twice from the same solvent with the addition of a little animal charcoal. The final product, consisting of colourless needles, weighed 3.6 g. and melted at 212°.

This product, when mixed with a sample of diacetylpuberulic acid obtained from *P. puberulum*, showed no depression in melting-point, thus indicating the identity of the two products.

Puberulic acid, $C_8H_6O_6$, was obtained by gentle hydrolysis of the pure diacetyl compound (2.97 g.) by boiling with 150 cc. of *N* NaOH in an atmosphere of nitrogen for 20 minutes. The hydrolysis mixture was cooled and acidified with 15.5 cc. of 10*N* H_2SO_4 , when the free acid separated as a cream-coloured microcrystalline powder, which could not be satisfactorily recrystallised from any solvent (wt. 1.98 g.). It was, therefore, purified for analysis by sublimation in a high vacuum at 220°. Samples of puberulic acid prepared by hydrolysis of the diacetyl compound, or by direct sublimation of the ether extract from *P. puberulum*, were identical, since they were cream-coloured, melted at 316° with decomposition and gave rise to the same diacetyl compound. Puberulic acid is only slightly soluble in water. On the addition of a slight

excess of FeCl_3 solution to a dilute alcoholic solution of the acid a deep reddish-brown colour is obtained. When the test is made by adding very dilute FeCl_3 solution to an alcoholic solution of the acid a yellow-brown colour is first obtained, followed by a distinct amethyst shade as further additions are made, and finally by the deep reddish-brown colour referred to above. Puberulic acid contains no nitrogen and no methoxyl groups. Micro-analysis (Schoeller): C, 48.60 %, 48.48 %; H, 3.16 %, 3.09 %. ($\text{C}_8\text{H}_6\text{O}_6$ requires C, 48.48 %; H, 3.05 %.)

Equivalent by titration, 95.4; for a dibasic acid, $\text{C}_8\text{H}_6\text{O}_6$, the calculated equivalent is 99.

The acid $\text{C}_8\text{H}_4\text{O}_6$ was prepared from the crude sodium salt separating from the diluted acetylation mixture of the mixed acids (p. 443). The crude sodium salt was dissolved in methyl alcohol, two volumes of ether were added to precipitate a dark-coloured amorphous material, and the product, after removal of solvent, was recrystallised from ethyl alcohol. It was finally obtained in beautiful orange needles containing 10.12 % Na. ($\text{C}_8\text{H}_3\text{O}_6\cdot\text{Na}$ requires Na 10.55 %.)

The pure sodium salt was dissolved in hot water and acidified with HCl . On cooling, the free acid separated in shining yellow plates, m.p. 296° (decomp.). An alcoholic solution of the acid gives with FeCl_3 a final colour indistinguishable from that given by puberulic acid. Micro-analysis (Schoeller) (two different samples of the free acid): C, 48.75 %, 48.92 %; 48.58 %, 48.52 %; H, 2.30 %, 2.39 %; 2.09 %, 2.08 %; mol. wt., 217, 187. ($\text{C}_8\text{H}_4\text{O}_6$ requires C, 48.97 %; H, 2.06 %; mol. wt., 196.)

Although it seems probable that puberulic acid and the acid $\text{C}_8\text{H}_4\text{O}_6$ are intimately related, the former being probably a reduced form of the latter, we have not yet succeeded in transforming either acid into the other.

Diacetylpuberulic acid, $\text{C}_8\text{H}_4\text{O}_4(\text{O}\cdot\text{CO}\cdot\text{CH}_3)_2$, was prepared from samples of puberulic acid obtained from both *P. puberulum* and *P. aurantio-virens*, by acetylation with sodium acetate and acetic anhydride as previously described. In each case the product was obtained as well-defined colourless prisms, melting at 212° and having the same mixed melting-point. The acetyl compound is readily soluble in aqueous sodium acetate solution and in alcoholic solution gives no colour with FeCl_3 .

Micro-analysis (Schoeller): C, 51.08 %, 51.24 %; H, 3.77 %, 3.72 %. Mol. wt. by Rast method, 216, 232. ($\text{C}_{12}\text{H}_{10}\text{O}_8$ requires C, 51.05 %; H, 3.57; mol. wt., 282.)

0.0812 g. dissolved in 10 cc. alcohol, required 5.72 cc. $N/10$ NaOH for neutralisation to phenolphthalein, corresponding to an equivalent of 142. (Theoretical for $\text{C}_{12}\text{H}_{10}\text{O}_8$ titrating as a dibasic acid, 141.) 0.4030 g. when hydrolysed in an atmosphere of nitrogen with $N/5$ NaOH yielded acetic acid equivalent to 13.91 cc. of $N/5$ NaOH (calc. for 2 acetyl groups, 14.29 cc.).

Dimethylpuberulic acid, $\text{C}_8\text{H}_4\text{O}_4(\text{OCH}_3)_2$. 1 g. of puberulic acid and 10 cc. (a large excess) of dimethyl sulphate were shaken vigorously and 10 %

aqueous NaOH were added continuously to maintain an alkaline reaction. When most of the dimethyl sulphate had disappeared an excess of NaOH was added, the mixture was boiled for some time, cooled, acidified, extracted with ether, and the ethereal solution evaporated to dryness. The residual solid, the yield of which was small, was crystallised from a mixture of methyl alcohol and water, and finally purified for analysis by sublimation in a high vacuum. The sublimate, which was colourless, melted at 271–272°, but gave a weak reddish-brown colour in alcoholic solution with FeCl_3 .

Micro-analysis (Schoeller): C, 53.10 %; H, 4.68 %; OCH_3 , 25.8 %. ($\text{C}_8\text{H}_4\text{O}_4(\text{OCH}_3)_2$ requires C, 53.08 %; H, 4.45 %; OCH_3 , 27.4 %.)

Equivalent by titration, 116. (Theoretical for $\text{C}_8\text{H}_4\text{O}_4(\text{OCH}_3)_2$, titrating as a dibasic acid, 113.)

A sample of dimethylpuberulic acid, made by hydrolysing tetramethylpuberulic acid (p. 446) with boiling aqueous *N* sodium hydroxide, gave no colour with FeCl_3 in alcoholic solution. This sample contained 26.67 % OCH_3 . It thus appears that the colour given with FeCl_3 by the sample prepared by methylation with methyl sulphate and sodium hydroxide is due to the presence of traces of partially methylated material and indicates that at least one hydroxyl group in puberulic acid is difficult to methylate under these conditions.

Dimethyldiacetylpuberulic acid, $\text{C}_8\text{H}_2\text{O}_2(\text{OCH}_3)_2(\text{O}.\text{CO}.\text{CH}_3)_2$. When diacetylpuberulic acid was treated with an excess of diazomethane in ether, a syrup was obtained which could not be crystallised. When, however, diacetylpuberulic acid was treated with just sufficient diazomethane in ether to dissolve it, at which point the initial violent evolution of nitrogen ceased, a crystalline product was obtained. 0.7 g. of diacetylpuberulic acid, treated in this way, gave, after evaporation in air of most of the ether, a mass of needles which separated from the ether mother-liquors. The product on crystallisation from methyl alcohol gave 0.45 g. of colourless needles, melting sharply at 102° and quickly turning red on exposure to light. The crystals were neutral in reaction and gave no colour with FeCl_3 in alcoholic solution.

Micro-analysis (Schoeller): C, 54.28 %; H, 4.51 %; OCH_3 , 19.94 %; mol. wt., 286. ($\text{C}_8\text{H}_2\text{O}_2(\text{OCH}_3)_2(\text{O}.\text{CO}.\text{CH}_3)_2$ requires C, 54.18 %; H, 4.55 %; OCH_3 , 20.01 %; mol. wt., 310.)

0.4261 g. when hydrolysed in nitrogen with *N* NaOH yielded acetic acid equivalent to 27.54 cc. of *N*/10 NaOH, indicating an acetyl content of 27.8 % (calc. 27.75 %).

0.1047 g. of the substance was hydrolysed by boiling for 2½ hours in an atmosphere of nitrogen with a measured excess (25 cc.) of *N*/5 NaOH. The mixture was cooled and the residual NaOH estimated. It was thus shown that an increase in acidity corresponding to 13.44 cc. *N*/10 acid had been produced, compared with the theoretical figure of 13.51 cc. required for the liberation of 4 acidic groups. It is thus obvious that, since two molecules of acetic acid are produced on hydrolysis, two other acid reacting groups have been unmasked during hydrolysis.

Monomethylpuberulic acid, $C_8H_5O_5(OCH_3)$. The alkaline hydrolysis mixture from dimethyldiacetylpuberulic acid yielded, when acidified, a product crystallising readily from water in the form of glistening plates, M.P. 272–276°, which on further heating decomposed slowly at 320–330°. It gave a brown colour in alcoholic solution with ferric chloride.

Micro-analysis (Schoeller): C, 51.23, 51.19 %; H, 3.81, 3.82 %; OCH_3 , 14.59, 14.43 %. ($C_8H_5O_5(OCH_3)$ requires C, 50.93 %; H, 3.80 %; OCH_3 , 14.63 %.)

Equivalent by titration, 105.1. (Theoretical for $C_8H_5O_5(OCH_3)$ titrating as a dibasic acid, 106.)

Tetramethyl derivative, $C_8H_2O_2(OCH_3)_4$. 1.5 g. of puberulic acid were suspended in dry ether and treated with a large excess of an ethereal solution of diazomethane. A vigorous effervescence took place. When this had ceased the solution was filtered from a small amount of insoluble material (0.08 g.) and evaporated to about 50 cc. On chilling the solution the product separated in clusters of pale yellow needles, weight 0.67 g.; M.P. 112–113°. The residue consisted of a yellow oil.

The tetramethyl derivative, surprisingly enough, is readily soluble in water, and in most other solvents, is neutral in reaction, gives no colour in alcoholic solution with $FeCl_3$, and contains no nitrogen.

Micro-analysis (Schoeller): C, 56.64, 56.64 %; H, 5.59, 5.62 %; OCH_3 , 48.18, 48.20 %. ($C_8H_2O_2(OCH_3)_4$ requires C, 56.67 %; H, 5.55 %; OCH_3 , 48.83 %.)

Molecular weight estimations (calc. 254) by the Rast method for which we are indebted to Prof. C. R. Harington gave abnormal results. They varied from an initial value of 474 to a final and fairly steady value of 275 after repeated remelting and cooling, in the course of which there was no obvious gross decomposition.

The results obtained may now be reviewed in the light of their bearing on the constitution of puberulic acid.

(1) Puberulic acid, which is cream-coloured, has the empirical formula $C_8H_6O_6$, titrates sharply to phenolphthalein as a dibasic acid, and gives a purple-brown colour with $FeCl_3$.

(2) Puberulic acid gives with diazomethane a tetramethyl derivative, $C_8H_2O_2(OCH_3)_4$, which is pale yellow-coloured, readily soluble in cold water, neutral in reaction, and gives no colour with $FeCl_3$. Puberulic acid therefore contains four OH groups.

(3) Puberulic acid gives a colourless diacetyl compound, $C_8H_4O_4(O.CO.CH_3)_2$, with acetic anhydride and sodium acetate, and a colourless dimethyl derivative, $C_8H_4O_4(OCH_3)_2$, with dimethyl sulphate and sodium hydroxide. Both the diacetyl and dimethyl derivatives titrate sharply to phenolphthalein as dibasic acids. Neither diacetyl- nor dimethyl-puberulic acid gives a colour with $FeCl_3$.

(4) Diacetylpuberulic acid, $C_8H_4O_4(O.CO.CH_3)_2$, gives with diazomethane dimethyldiacetylpuberulic acid in colourless crystals turning pink on exposure

to light. This compound, $C_8H_2O_2(OCH_3)_2(O.CO.CH_3)_2$, is neutral in reaction, and gives no colour with $FeCl_3$. On gentle hydrolysis it loses both acetyl groups and one OCH_3 group to give monomethylpuberulic acid, $C_8H_5O_5(OCH_3)$, a substance titrating sharply to phenolphthalein as a dibasic acid, and giving a brown colour with $FeCl_3$.

It is thus abundantly clear that puberulic acid contains four OH groups, two of which are probably phenolic in nature, since puberulic acid gives a strong colour with $FeCl_3$, and the diacetyl and dimethyl derivatives give no colour with $FeCl_3$. The nature of the two remaining OH groups, which confer strong acidic properties on puberulic acid, but give no colour with $FeCl_3$, is not quite certain. The most obvious explanation is that they are present in carboxyl groups in which case the formula for puberulic acid becomes $C_6H_2(OH)_2(COOH)_2$ and the acid would probably be a dihydroxybenzenedicarboxylic acid. This explanation agrees with most of the facts presented. Thus the diacetyl and dimethyl compounds would be diacetyl- and dimethoxybenzenedicarboxylic acids, while the diacetyldimethyl- and tetramethylcompounds would be the dimethyl esters of diacetyl- and dimethoxybenzenedicarboxylic acids respectively. The most serious objection to this hypothesis arises when puberulic acid and its derivatives are compared with the known dihydroxybenzenedicarboxylic acids, eleven of which are theoretically possible.

These may be grouped as derivatives of phthalic, isophthalic and terephthalic acids.

A. Dihydroxyphthalic acids, $COOH : COOH : 1 : 2$; (i) 3 : 4-dihydroxy; (ii) 3 : 5-dihydroxy; (iii) 3 : 6-dihydroxy; (iv) 4 : 5-dihydroxy. Of these (i), (iii) and (iv) cannot be identical with puberulic acid, since they all readily form anhydrides, the colours they give with $FeCl_3$ are deep blue, blue-violet and emerald green respectively, and they melt at $210-212^\circ$, $219-220^\circ$ and $183-185^\circ$ respectively. Acid (ii) has not been synthesised. β -Resodicarboxylic acid was formerly thought to have this constitution, but is now known to have formula (x) [Brunner, Madersbacher and Goritschan, 1928]. Since, however, acid (ii) would almost certainly readily form an anhydride, it is improbable that puberulic acid has this constitution.

B. Dihydroxyisophthalic acids, $COOH : COOH : 1 : 3$; (v) 2 : 4-dihydroxy; (vi) 2 : 5-dihydroxy; (vii) 4 : 5-dihydroxy; (viii) 4 : 6-dihydroxy.

Acid (v), α -resodicarboxylic acid, has a recorded melting-point (305° , 312°) sufficiently near to that of puberulic acid (316°) to suggest a possible identity. A quantity of acid (v) was therefore synthesised from β -resorcylic acid by the method of Errera [1899]. It melted at 309° , a sublimed sample of puberulic acid melted at 316° , and a mixture of the two at 292° under identical conditions of heating. This fact, together with a totally different colour reaction with $FeCl_3$ —wine-red against the purple-brown given by puberulic acid—definitely exclude (v) as the constitution of puberulic acid. Further, acid (v), in contrast with puberulic acid, does not give an acetyl derivative on heating with acetic anhydride and sodium acetate.

Acid (vi) has not been previously synthesised. Its synthesis by two different methods is described in the Appendix. The acid prepared by either method gave a blue-violet colour with FeCl_3 and melted at $248\text{--}250^\circ$, the dimethyl ether melted at 172° . These melting-points are very different from those of puberulic acid and its dimethyl ether, 316° , and 272° . Acid (vi) is thus definitely not identical with puberulic acid.

The free acid (vii) has not been described, but it can scarcely be identical with puberulic acid since its dimethyl derivative melts at $245\text{--}246^\circ$ [Tiemann and Mendelsohn, 1877] while dimethylpuberulic acid melts at $271\text{--}272^\circ$, and, since acid (vii) is a catechol derivative, it would probably give a blue or green colour with FeCl_3 .

The free acid (viii) has not been described previously, although a dimethyl ether, m.p. 266° , prepared by Eykman [1905], was thought by its discoverer to be 4 : 6-dimethoxybenzene-1 : 3-dicarboxylic acid but is now shown (see Appendix) to be identical with α -resodicarboxylic acid. We prepared a quantity of this material by the method given by the above author and, by demethylation with concentrated HI, obtained the parent dihydroxy-acid. The latter melts at $311\text{--}312^\circ$ but is not identical with puberulic acid since the acids have a mixed m.p. of 288° , it gives a wine-red colour with FeCl_3 , and does not give an acetyl derivative on heating with acetic anhydride and sodium acetate.

C. Dihydroxyterephthalic acids, $\text{COOH}:\text{COOH}:1:4$; (ix) 2 : 3-dihydroxy; (x) 2 : 6-dihydroxy; (xi) 2 : 5-dihydroxy.

Acid (ix), described by Kawai [1926], cannot be identical with puberulic acid. It melts at 290° , gives a deep indigo-blue colour with FeCl_3 , and its dimethyl ether melts at $213\text{--}215^\circ$.

Acid (x), β -resodicarboxylic acid, was formerly thought to have formula (ii), but has now been shown [Brunner, Madersbacher and Goritschan, 1928] to have the formula indicated. A specimen of this acid was made and compared with puberulic acid. The two acids are definitely not identical. Acid (x) melted at 284° , and gave a blue-violet colour with FeCl_3 in alcohol. Further, its dimethyl ether was found to melt at 301° (Wedekind and Fleischer [1924] give 295°) and its diacetyl derivative is recorded as melting at 198° [Brunner, Madersbacher and Goritschan, 1928].

Acid (xi), which is yellow in colour, is well known. A specimen of the acid made by Herrmann's method [1882] melted at 346° , and gave a pure blue colour with FeCl_3 . Further, since its diacetyl derivative was found not to melt, even at 360° (Thiele and Gunther [1906] state that it does not melt up to 260°), there can be no doubt that puberulic acid has not the formula of acid (xi).

If the evidence afforded by the dihydroxybenzenedicarboxylic acids which we have prepared, and if the chemical constitutions and physical characteristics of the other dihydroxybenzenedicarboxylic acids described in the literature are accepted as correct, the possibility of puberulic acid being a dihydroxybenzenedicarboxylic acid is practically excluded, though there is, of course, the remote possibility that it may be either 2 : 4-dihydroxybenzene-1 : 3-di-

carboxylic acid or 4 : 6-dihydroxybenzene-1 : 3-dicarboxylic acid, depending on which of these formulae is finally assigned to α -resodicarboxylic acid. Even this possibility seems to be ruled out by the yellow colour of the tetramethyl derivative and its ready solubility in water.

In an attempt to demonstrate the presence of aldehydic or ketonic groups the following qualitative tests were carried out in each case on cold neutral aqueous solutions of the ammonium salts of the free acid, the monomethyl and diacetyl derivatives, and on cold aqueous solutions of the tetramethyl and the diacetyldimethyl derivatives.

- (i) Schiff's reagent gave a negative in all cases.
- (ii) 2 : 4-Dinitrophenylhydrazine gave no precipitate in any case.
- (iii) Cold ammoniacal silver nitrate gave an immediate strong reduction with puberulic acid. There was no immediate reduction with any of the derivatives, though the diacetyl derivative gave a partial reduction after standing for about an hour and the diacetyldimethyl and monomethyl derivatives after standing overnight. On the other hand, tetramethylpuberulic acid showed no sign of reduction, even after standing for 2 days.
- (iv) Cold neutral KMnO_4 . Immediate strong reduction was given by puberulic acid and monomethylpuberulic acid. A moderately quick though not immediate reduction was given by diacetylpuberulic acid, and a slow reduction on standing was given by the diacetyldimethyl and tetramethyl derivatives.

Tests (i) and (ii) give no support to the view that puberulic acid, or any of its derivatives, contains either an aldehydic or ketonic grouping. On the other hand, test (iii) might be taken to indicate the presence of an aldehydic grouping in puberulic acid. In this connection it is of interest to note that while the so-called 2 : 4-dihydroxybenzene-1 : 3-dicarboxylic acid (acid v) and 2 : 6-dihydroxybenzene-1 : 4-dicarboxylic acid (acid x) give no immediate reduction with either ammoniacal silver or cold neutral KMnO_4 solution, 2 : 5-dihydroxybenzene-1 : 3-dicarboxylic acid (acid vi) and 2 : 5-dihydroxybenzene-1 : 4-dicarboxylic acid (acid xi) give immediate strong reduction of both reagents.

We therefore propose to leave the question of the exact determination of the constitution of puberulic acid to a future communication, in the opinion that, in spite of resemblances to certain of the dihydroxybenzenedicarboxylic acids, it is improbable that puberulic acid is a compound of this type.

Since there are good *a priori* reasons for assuming that puberulic acid $\text{C}_8\text{H}_6\text{O}_6$ and the acid $\text{C}_8\text{H}_4\text{O}_6$ are closely related, *e.g.* their formulae, their colour reactions with FeCl_3 and the fact that both acids are present in the same metabolism solution, it is tempting to speculate on the possibility of these two acids forming an oxidation-reduction system for the utilisation of atmospheric oxygen by the moulds which produce them—*P. puberulum* and *P. aurantiovirens*. This aspect of the question we propose to develop also in a future communication.

SUMMARY.

The production is described of two new mould metabolic products, puberulic acid $C_8H_6O_6$ and an acid $C_8H_4O_6$, in synthetic solutions containing glucose as sole source of carbon. Each of these acids is produced by *P. puberulum* Bainier and *P. aurantio-virens* Biourge. Puberulic acid and its monomethyl, dimethyl and diacetyl derivatives are dibasic acids, whereas the tetramethyl and diacetyldimethyl compounds are neutral substances. The possibility of puberulic acid being a dihydroxybenzenedicarboxylic acid is considered, but it is thought that this is improbable. It is suggested that puberulic acid $C_8H_6O_6$ and the acid $C_8H_4O_6$ may form an oxidation-reduction system.

We desire to express our thanks to Prof. Robert Robinson for much kindly advice and helpful criticism.

REFERENCES.

- Alsberg and Black (1913). U.S. Dept. of Agriculture, Bureau of Plant Industry. Bulletin No. 270.
 Biourge (1923). *La Cellule*, **33**, 119.
 Birkinshaw, Charles, Lilly and Raistrick (1931). *Phil. Trans. Roy. Soc. Lond.* B **220**, 127.
 Brunner, Madersbacher and Goritschan (1928). *Monatsh. Chem.* **50**, 216.
 Errera (1899). *Ber. deutsch. chem. Ges.* **32**, 2797.
 Eykman (1905). *Chem. Weekblad*, **2**, 59-72, 79-93.
 Herrmann (1882). *Liebig's Ann.* **211**, 327.
 Kawai (1926). *Chem. Zentr.* i, 3144.
 Thiele and Gunther (1906). *Liebig's Ann.* **349**, 61.
 Thom (1930). *The Penicillia*. (Baillière, Tindall and Cox, London.)
 Tiemann and Mendelsohn (1877). *Ber. deutsch. chem. Ges.* **10**, 398.
 Wedekind and Fleischer (1924). *Ber. deutsch. chem. Ges.* **57**, 1121.

APPENDIX ON CERTAIN DIHYDROXYBENZENEDICARBOXYLIC ACIDS.

In the course of work described above it became necessary to prepare certain new dihydroxybenzenedicarboxylic acids, and also certain derivatives, not previously described, of dihydroxybenzenedicarboxylic acids already appearing in the literature. An account of this work follows.

Synthesis of 2:5-dihydroxybenzene-1:3-dicarboxylic acid.

This acid, which has not previously been described, was synthesised from 2:5-dihydroxy-1:3-dimethylbenzene [cf. Bamberger and Rising, 1901] by conversion (a) into the potassium sulphuric ester, or (b) into the dimethyl ether followed by oxidation and hydrolysis.

(a) *2:5-Dihydroxy-1:3-dimethylphenyl dipotassium disulphate*. 2 g. of 2:5-dihydroxy-1:3-dimethylbenzene were mixed with 4.6 g. of *N*-pyridinium sulphonic acid and heated in an oil-bath at 100°. The mixture slowly began to melt. A few drops of pyridine were then added. After a few minutes the mixture was dissolved in water and extracted with ether. 2.5 g. of potassium hydroxide (i.e. a slight excess) in water were added to the ether-extracted solution which was evaporated *in vacuo* to crystallisation. Yield of crystalline product, 3.2 g.

2:5-Dihydroxybenzene-1:3-dicarboxylic acid. 3.2 g. of the above compound were dissolved in about 300 cc. of water and oxidised with 5.5 g. of potassium permanganate at 100°. The mixture was then rendered weakly acid with sulphuric acid and hydrolysed for half an hour at 100°. On keeping, a brownish crystalline deposit was obtained, which was extracted with ether, filtered

from the insoluble colouring matter, and the fluorescent ether solution was evaporated, giving a mass of crystalline needles. The acid was further purified by neutralisation with sodium hydroxide, filtration from a small amount of insoluble matter, and reprecipitation by addition of a slight excess of sulphuric acid. It crystallised in beautiful long colourless needles with $1\text{H}_2\text{O}$, m.p. 248–250°, giving a blue-violet colour with FeCl_3 . 0.0905 g. dried at 100° to constant weight (2 hours) lost 0.0068 g., i.e. 7.52 %. Theoretical for $\text{C}_8\text{H}_6\text{O}_6 \cdot \text{H}_2\text{O}$, 8.33 %. 0.0837 g. (anhydrous) required 8.54 cc. $N/10$ NaOH giving an equivalent of 98 (calc. 99).

Micro-analysis (Schoeller) on the material dried at 100° gave the following results: C, 48.61 %; H, 3.10 %. (Theoretical for $\text{C}_8\text{H}_6\text{O}_6$, C, 48.48 %; H, 3.05 %.)

(b) 2 : 5-Dimethoxy-1 : 3-dimethylbenzene. 13.8 g. of crude 2 : 5-dihydroxy-1 : 3-dimethylbenzene were shaken with 40 cc. of 20 % sodium hydroxide and 26 g. of dimethyl sulphate. After some time a further addition of alkali and dimethyl sulphate was made, and the mixture was heated for a few minutes under reflux, the reaction being maintained permanently alkaline. The resulting product was distilled in steam, extracted from the distillate with ether, dried with calcium chloride and distilled *in vacuo*. Practically the whole of it boiled at 117°/25 mm. Yield 10.6 g.

2 : 5-Dimethoxybenzene-1 : 3-dicarboxylic acid. To 38 g. of KMnO_4 in 750 cc. of water were added 10 g. of the dimethoxy compound, and the mixture was heated in a water-bath under reflux for several hours. The hydrated manganese dioxide was then removed by filtration and the filtrate acidified with sulphuric acid and evaporated. The residue was extracted with ether and the product was recrystallised three times from water, from which it separated in silky needles melting at 172°. The yield was only small as the loss on oxidation was considerable. 0.0809 g. required 7.17 cc. $N/10$ NaOH giving an equivalent of 112.9. (Theoretical for $\text{C}_8\text{H}_2(\text{COOH})_2(\text{OCH}_3)_2$, 113.) In a Zeisel estimation 0.0912 g. gave 0.1872 g. of silver iodide. OCH_3 , 27.1 %. (Theoretical for $\text{C}_8\text{H}_2(\text{COOH})_2(\text{OCH}_3)_2$, 27.4 %.) The acid gave no colour with ferric chloride.

2 : 5-Dihydroxybenzene-1 : 3-dicarboxylic acid. The residue from the Zeisel estimation gave well formed needles on cooling. These were separated by filtration from the hydriodic acid, washed and recrystallised from water. The product agreed in m.p., colour with ferric chloride, and mixed m.p. with the 2 : 5-dihydroxybenzene-1 : 3-dicarboxylic acid synthesised by method (a).

Preparation of 4 : 6-dihydroxybenzene-1 : 3-dicarboxylic acid.

The compound described by Eykman [1905] as 4 : 6-dimethoxybenzene-1 : 3-dicarboxylic acid was prepared in quantity by his method and had m.p. 276° (decomp.) (Eykman, 266°). It crystallised in colourless needles, gave no colour with FeCl_3 , and gave the following results on analysis (Schoeller), C, 52.89 %; H, 4.45 %; OCH_3 , 26.8 %. (Theoretical for $\text{C}_8\text{H}_2(\text{OCH}_3)_2(\text{COOH})_2$, C, 53.08 %; H, 4.45 %; OCH_3 , 27.4 %.)

0.0523 g. required 4.645 cc. $N/10$ NaOH for neutralisation to phenolphthalein corresponding to an equivalent of 112.6. (Theoretical for $\text{C}_8\text{H}_2(\text{OCH}_3)_2(\text{COOH})_2$ titrating as a dibasic acid, 113.) The dimethyl ester (OCH_3 , 48.7 %; calc., 48.8 %), prepared with diazomethane in ethereal solution, formed glistening plates, m.p. 151–153° after softening at 150°.

4 : 6-Dihydroxybenzene-1 : 3-dicarboxylic acid. 5 g. of the dimethoxy-acid were de-methylated with 50 cc. of HI (sp. gr. 1.7) in the Zeisel apparatus. The mixture was heated to 120–130° for 2½ hours, when no further reaction was obtained from the exit gas with alcoholic silver nitrate, cooled and allowed to stand. The solid material separating was filtered through a sintered glass funnel, washed with water and dried. It melted at 311–312°, with decomposition, and gave a wine-red colour in alcoholic solution with ferric chloride.

Micro-analysis (Schoeller) gave the following results: C, 48.53 %; H, 3.09 %. (Theoretical for $\text{C}_8\text{H}_6\text{O}_6$, C, 48.48 %; H, 3.05 %.)

0.1000 g. required 10.12 cc. of $N/10$ sodium hydroxide giving an equivalent of 98.8. (Theoretical for $\text{C}_8\text{H}_6\text{O}_6$ titrating as a dibasic acid, 99.) It was found impossible to acetylate this acid by heating with acetic anhydride and sodium acetate. The only product which could be recovered (0.62 g. from 1 g. of the dihydroxy-acid) melted at 302°, gave a deep red colour with FeCl_3 , and contained no free acidic groups.

α-Resodicarboxylic acid.

α-Resodicarboxylic acid, which is thought to have the constitution 2 : 4-dihydroxybenzene-1 : 3-dicarboxylic acid, was synthesised from *β*-resorecylic acid by the method of Errera [1899].

The colourless product, melting at 309° with decomposition, gave a wine-red colour with FeCl_3 , and did not give an acetyl compound on heating with acetic anhydride and sodium acetate.

Micro-analysis (Schoeller) gave the following results: C, 48.27 %; H, 3.09 %. (Theoretical for $\text{C}_8\text{H}_6\text{O}_6$, C, 48.48 %; H, 3.05 %.)

0.0472 g. required 4.72 cc. $N/10$ NaOH for neutralisation to phenolphthalein, corresponding to an equivalent of 100. (Theoretical for $\text{C}_8\text{H}_6\text{O}_6$ titrating as a dibasic acid, 99.)

Direct comparison of α -resodicarboxylic acid with the acid prepared by demethylation of Eykman's dimethoxybenzenedicarboxylic acid (see above) showed that they are probably the same substance. Their melting-points are 309°, and 311–312° respectively, while a mixture of the two melted at 309–310°. They each give the same wine-red colour with FeCl_3 , and neither of them gives an acetyl derivative on heating with acetic anhydride and sodium acetate.

Dimethyl- α -resodicarboxylic acid. α -Resodicarboxylic acid was methylated three times with methyl sulphate and excess of sodium hydroxide (20 %). The methylation mixture was allowed to warm up spontaneously and then boiled for a short time, cooled and finally acidified with H_2SO_4 . This protracted methylation is necessary since α -resodicarboxylic acid is difficult to methylate. The solid separating was washed and dried.

Micro-analysis (Schoeller) gave the following results: C, 53.16 %; H, 4.61 %; OCH_3 , 26.9 %. (Theoretical for $\text{C}_8\text{H}_2(\text{OCH}_3)_2(\text{COOH})_2$, C, 53.08 %; H, 4.45 %; OCH_3 , 27.4 %.)

0.0506 g. required 4.54 cc. $N/10$ NaOH for neutralisation to phenolphthalein, corresponding to an equivalent of 111.7. (Theoretical for $\text{C}_8\text{H}_2(\text{OCH}_3)_2(\text{COOH})_2$ titrating as a dibasic acid, 113.)

The product gave no colour with FeCl_3 and had a melting-point of 275° (decomp.). A mixed melting-point with Eykman's dimethoxybenzenedicarboxylic acid was also 275°.

Treatment with diazomethane in ether gave a dimethyl ester identical with that similarly prepared from Eykman's acid. The same dimethyl ester is given, though in smaller yield since methylation even with diazomethane is not complete, by methylating α -resodicarboxylic acid directly with diazomethane.

It is thus obvious that α -resodicarboxylic acid and the dihydroxybenzenedicarboxylic acid prepared from Eykman's 4:6-dimethoxybenzene-1:3-dicarboxylic acid are one and the same substance. Since α -resodicarboxylic acid is thought to have the constitution 2:4-dihydroxybenzene-1:3-dicarboxylic acid, and since Eykman's product is thought to be the dimethyl derivative of 4:6-dihydroxybenzene-1:3-dicarboxylic acid, it is obvious that the formula for one or other of these acids must be incorrect.

The formula for α -resodicarboxylic acid and the evidence supporting this formula were presented by Waitz [1911]. Waitz showed that when either β -resorecylic acid or γ -resorecylic acid is heated with ammonium carbonate and water at 130° for 20 hours, α -resodicarboxylic acid is produced in each case.

Waitz points out that this proof of constitution is open to the objection that one or other of the substituent groups may have "wandered," but supports his evidence by showing that when α -resodicarboxylic acid is heated in water for 48 hours at 100° in a sealed tube, a mixture of β - and γ -resorecylic acids is formed. Accepting this evidence therefore, and omitting any possibility of the "wandering" of substituent groups, there appears to be no alternative to the acceptance of the statement that α -resodicarboxylic acid is 2:4-dihydroxybenzene-1:3-dicarboxylic acid.

The evidence for the constitution of Eykman's dimethoxybenzenedicarboxylic acid, thought by him to be 4:6-dimethoxybenzene-1:3-dicarboxylic acid, obviously depends on the orientation of the groups in the parent dihydroxydiacetylbenzene or "resodiacetophenone" which was prepared by heating resorcinol diacetate with ZnCl_2 . Eykman prepared a large number of mixed ethers by the action of alkyl iodides or bromides on the potassium salt of his dihydroxydiacetylbenzene. Since mixed ethers of the type $\text{C}_6\text{H}_2(\text{COCH}_3)_2(\text{OR}_1)(\text{OR}_2)$ are identical, independently of the order of the introduction of the groups R_1 and R_2 , the parent acetyl compound must be

4:6-dihydroxy-1:3-diacetylbenzene and hence the acid resulting from the oxidation of the dimethyl ether must be 4:6-dimethoxybenzene-1:3-dicarboxylic acid.

Further work is contemplated to settle which of these constitutions is correct.

SUMMARY.

The synthesis by two different methods of 2:5-dihydroxybenzene-1:3-dicarboxylic acid is described. It is proved that α -resodicarboxylic acid, at present thought to be 2:4-dihydroxybenzene-1:3-dicarboxylic acid, is identical with the dihydroxybenzenedicarboxylic acid obtained by demethylation of the substance described by Eykman as 4:6-dimethoxybenzene-1:3-dicarboxylic acid. This follows since the dihydroxy-acids, the dimethyl ethers and the dimethyl ether dimethyl esters are identical in each case.

REFERENCES.

- Bamberger and Rising (1901). *Liebig's Ann.* **316**, 202.
Errera (1899). *Ber. deutsch. chem. Ges.* **32**, 2797.
Eykman (1905). *Chem. Weekblad*, **2**, 59-72, 79-93; *J. Chem. Soc. Abstr.* **88**, 1, 359.
Waitz (1911). *Monatsh. Chem.* **32**, 427.