CLXXVI. THE METABOLISM OF NORMAL AND TUMOUR TISSUE.

XVII. THE ACTION OF SOME DERIVATIVES OF PHENAZINE, QUINOLINE AND PYRIDINE ON THE PASTEUR REACTION.

By FRANK DICKENS.

From the Cancer Research Laboratory, North of England Council of the British Empire Cancer Campaign, Royal Victoria Infirmary, Newcastle-upon-Tyne.

(Received 27 May 1936.)

IN the course of work on the action of reversible oxidation-reduction dyes upon tissue metabolism it was found [Dickens, 1935, 1, 2] that the negative dyestuff phenosafranine (E_h at pH 7 = -0.252 v.) even in 10^{-5} *M* solution could inhibit completely the Pasteur mechanism in normal and tumour tissues. This action has been further studied in the present paper and has been found to extend to other substances showing a certain chemical relationship with phenosafranine.

Methods.

The experiments were made at 37.5° in the Warburg apparatus, using tissue slices suspended in bicarbonate- or phosphate-Ringer solution at pH 7.4 containing 0.2 % glucose. The duration of the experiments was usually 1 hour, unless otherwise stated. In these experiments it was desirable to follow the time course of the metabolism, and for this Warburg's 2-vessel method is most suitable. For measurement of R.Q. the method of Dickens & Simer was used.

Phenosafranine (B.D.H.) could be dissolved directly in the solution, but many of the substances added are sparingly soluble in the cold and were therefore dissolved with warming and grinding in the salt solution before adding bicarbonate or glucose; even in the high dilutions used, solution of some of the reagents was incomplete.

Phenosafranine.

Added to slices of normal, highly glycolysing tissues (brain cortex, extra-embryonic membranes) or tumour tissue (JRS, Walker 256 carcinoma of rat), phenosafranine in appropriate concentration has little effect on respiration, but causes an immediate increase of aerobic glycolysis (Table I). The optimum concentration is close to 10^{-5} *M* for brain and a little higher for the other tissues. Above the optimum concentration the respiration is rapidly poisoned, whilst at lower concentrations (10^{-6} *M*) there is no change from the normal metabolism. The R.Q. of brain is not appreciably depressed below the carbohydrate level by 10^{-5} *M* phenosafranine, and chemical estimation by Clausen's method showed that the acid formed is lactic acid (Exp. 3, Table I). In some experiments a slight increase of respiration occurred with phenosafranine; however, the respiration of brain fell rather rapidly with concentrations of 5×10^{-5} *M* or a little above, but not to the same extent with the other tissues in Table I. The aerobic glycolysis in presence of phenosafranine rises until it becomes nearly equal to the anaerobic glycolysis; this is best seen with the more strongly

Table I. *Action of phenosafranine (formula I) on tissue metabolism.*

Medium	Conc. <i>M</i>	Phenosafranine			Control		
Rat brain cortex		Q_{O_2}	$Q_G^{O_2}$	$Q_G^{N_2}$	Q_{O_2}	$Q_G^{O_2}$	$Q_G^{N_2}$
1. Phosph.	10^{-4}	- 8.3			} - 10.6	—	—
	3×10^{-5}	- 9.2	—	—			
	10^{-5}	- 11.3					
2. Bicarb.	10^{-5}	- 12.8	+ 20.5		} - 12.6	+ 1.4	—
	10^{-6}	- 10.0	+ 1.9	—			
3. Bicarb. 2 hours:	10^{-5}	{ - 13.2 dupl. { - 15.7	+ 9.7	R.Q. 0.95	- 13.2	+ 1.7	R.Q. 1.03
			+ 12.4	1.01	—	—	—
Chemical estimation of lactic acid:		0.71 mg.			0.07 mg.		
Manometric ditto.:		0.77 mg.			0.09 mg.		
4. Bicarb. in darkness:	10^{-5}	- 10.3	+ 10.5	—	- 11.3	+ 0.3	—
Mouse yolk-sac							
5. Bicarb.	5×10^{-5}	- 16.1	+ 29.8	+ 27.0	- 12.3	+ 6.8	+ 32.0
Jensen rat sarcoma							
6. Bicarb.	5×10^{-5}	90 min.					
		- 9.3	+ 35.5	—	- 11.3	+ 22.9	—
	3×10^{-4}	- 9.7	+ 44.9	—	- 10.8	+ 28.3	—
Walker 256 carcinoma							
7. Phosph.	3×10^{-4}	- 9.3			} - 9.0	—	—
	10^{-4}	- 7.7	—	—			
	3×10^{-5}	- 8.4					
Bicarb.	10^{-4}	—	+ 36.2	+ 45.8	—	+ 24.5	+ 42.0
8. JRS. Anaerobic glycolysis; phenosafranine tipped in after 40 min.							
		$Q_G^{N_2}$ (2nd. 40 min.)	conc. 10^{-3}		3×10^{-4}	10^{-4}	Control
		$Q_G^{N_2}$ (1st. 40 min.)	$\times 100$		61 %	81 %	86 %
							84 %

glycolysing embryonic and tumour tissues (Exps. 5 and 7). Anaerobic glycolysis of these tissues is little affected (Exps. 5, 7 and 8) by concentrations of $3 \times 10^{-4} M$ or less.

Ashford & Dixon [1935] have shown that high concentrations of KCl increase respiration and aerobic glycolysis, and depress anaerobic glycolysis, of brain. Dickens & Greville [1935] showed that this effect was restricted to brain tissue. The action of phenosafranine differs from the KCl effect in several important respects: (1) the anaerobic glycolysis is not depressed by phenosafranine; (2) the action of phenosafranine is not restricted to brain; (3) the respiration, though sometimes a little increased, is not affected so much as by KCl; (4) the active concentration of phenosafranine is so low and its action on metabolism so selective that it is difficult to conceive of its action otherwise than as a specific anticatalyst of certain cell enzymes; whereas the action of KCl, or of disturbance of salt balance [Dickens & Greville, 1935], is an action on permeability and dispersion of cell colloids [Dickens & Greville, 1935; Dixon, 1936; Dixon & Holmes, 1935].

Mechanism of phenosafranine action: the effect of related compounds.

In seeking an explanation of this action, it appeared possible that the negative potential of phenosafranine (E_h at pH 7 = 0.252 v.) might be responsible. Thus Lipmann [1934] considers that negative systems favour fermentative reactions

and Bumm & Appel [1932] and Quastel & Wheatley [1932] have described a similar property of cysteine or reduced glutathione.

Consequently other negative dyes were tested (Table II) for their effects on respiration and glycolysis of rat brain cortex, but benzylviologen ($E_h-0.359$ v.) and even the chemically closely related compound neutral red ($E_h-0.34$ v.) were without action. Hence factors other than the negative potential are needed for the phenosafranine action.

Table II. *Other negative dyes.*

Medium	Conc.	With dye		Control	
Neutral red		Q_{O_2}	$Q_G^{O_2}$	Q_{O_2}	$Q_G^{O_2}$
Bicarb.	ca. 10^{-4}	- 8.8	+ 1.7	- 10.0	+ 1.4
Benzylviologen					
Bicarb.	10^{-5}	- 13.0	+ 2.5	} - 11.1	+ 1.1
	10^{-4}	- 8.9	+ 1.0		

Phenosafranine belongs to a group of dyes which are in use in photography as photosensitizers and desensitizers, phenosafranine itself being a powerful desensitizer. The precise mechanism of this photochemical action is still obscure, but it was decided to study the effects of members of this group on tissue metabolism. The formulae and reference numbers of the substances tested are shown on p. 1236, and their actions on metabolism of rat brain cortex are given in Table III.

Table III. *Sensitizers and desensitizers: action on metabolism of rat brain cortex.*

Formula No.	Name of dye	Sens. or desens.	Conc. M	Metabolism	
				Q_{O_2}	$Q_G^{O_2}$
	None added	—	—	- 10.5	+ 1.1 (average of controls)
I	Phenosafranine	D	(For data see Table I)		
II	<i>p</i> -Dimethylaminophenyl-azomethine-2-quinoline ethiodide	D	10^{-5}	- 9.2	+ 17.5
III	Pinacryptol yellow	D	ca. 10^{-5}	- 9.0	+ 11.3
IV	Pinacyanol	S	ca. 10^{-5}	- 13.8	+ 16.8
V	<i>p</i> -Dimethylaminobenzylidene-quinaldine ethiodide	S	10^{-5}	- 8.5	+ 18.6
VI	<i>p</i> -Dimethylaminobenzylidene- α -picoline methiodide	S	10^{-4}	- 7.6	+ 32.0 (toxic)
VII	Ethyl red	S	10^{-5}	- 12.0	+ 26.0
VIII	<i>p</i> -Dimethylaminobenzylidene- β -napthoquinaldine ethiodide	S	10^{-5}	- 9.2	+ 17.1
	Derivatives of fluorescein:				
	Erythrosin B.	S	10^{-4}	- 8.2	+ 3.4
			10^{-5}	- 8.2	+ 0.6
	Eosin	S	10^{-4}	- 8.1	+ 0.8

It is evident from this table that representatives of both groups, sensitizers and desensitizers alike, may act in the same way as phenosafranine. Some of those tested quickly poisoned the respiration even in low concentration and

were therefore discarded; others had hardly any action on respiration and in a few cases a slight increase was observed.

With all the active dyes however the aerobic glycolysis rose almost immediately on adding the dyes to a value approximately equal to that found in nitrogen.

The concentrations needed are very low, usually $10^{-5} M$ as with phenosafranine. In higher concentration the dyes were found to be markedly poisonous, whilst in lower concentration ($10^{-6} M$) they were usually inactive.

Desensitizers.

Phenosafranine

I.

p-Dimethylaminophenylazomethine-2-quinoline ethiodide

II.

Pinacryptol yellow

III.

Sensitizers.

Pinacyanol

IV.

p-Dimethylaminobenzylidene-quinaldine ethiodide

V.

p-Dimethylaminobenzylidene- α -picoline methiodide

VI.

Ethyl red

VII.

p-Dimethylaminobenzylidene- β -naphthoquinaldine ethiodide

VIII.

Related compounds.

Quinanil (Cohen)

X.

Acriflavine

IX.

Benzylidenequinaldine ethiodide

XI.

Since both photosensitizers and desensitizers act similarly, it seems unlikely that the metabolic effect is connected with the photoactivity of the dyes, but more probably with a definite type of chemical constitution. This view is supported by the following facts: (a) the action of phenosafranine proceeds in as nearly complete darkness as is possible for manometric readings with the usual

apparatus (Table I, Exp. 4); (b) eosin and erythrosin (also photosensitizers, but of completely different chemical constitution) have little or no effect on metabolism (Table III, Exps. 9 & 10); (c) chemically related compounds, some of which are not known to be sensitizers or desensitizers, act similarly to those already described (see Table IV).

Table IV. *Various related compounds: action on rat brain cortex.*

Formula No.	Substance added	Medium	Conc. <i>M</i>	Metabolism		Control	
				<i>Q</i> _{O₂}	<i>Q</i> _G ^{O₂}	<i>Q</i> _{O₂}	<i>Q</i> _G ^{O₂}
Group I							
IX. Acriflavine	Phosph.	10 ⁻⁴	- 7.5	falling		- 12.3	
		10 ⁻⁵	- 11.0				
	Bicarb.	10 ⁻⁵	- 10.5	+ 16.5		- 9.1	+ 4.2
X. Quinanil (<i>p</i> -Dimethylamino-phenylazomethine-2-quinoline metho- chloride)	Bicarb.	10 ⁻⁵	- 11.2	+ 13.0	(1st hour)	- 9.8	+ 2.5
			- 8.9	+ 13.2	(2nd hour)	- 10.6	+ 0.8
	Phosph.	10 ⁻⁵	- 11.5	(1st hour)			
			- 9.5	(2nd hour)			
XI. Benzyldinequinaldine ethiodide	Bicarb.	10 ⁻⁵	- 6.6	+ 17.6	<i>Q</i> _G ^{N₂} + 19.5	- 12.0	+ 1.9 <i>Q</i> _G ^{N₂} + 19.2
		10 ⁻⁶	- 12.0	+ 3.8		- 13.0	+ 0.7
Group II.							
β-Diethylamino-ethyl-1-aminoacridine hydrochloride [Clemo & Hook, 1936]	Bicarb.	10 ⁻⁵	- 7.4	+ 2.7		- 7.8	+ 1.9
Quinoline methiodide	Bicarb.	10 ⁻⁴	- 10.5	+ 3.5		- 13.0	+ 2.0
Quinaldine ethiodide	Bicarb.	10 ⁻⁵	- 12.0	+ 1.7			

It is possible from the data presented in Tables I-IV to draw some provisional inferences as to the general chemical constitution needed in substances of this group in order that they may specifically inhibit the Pasteur mechanism. All such substances are derivatives of a N-containing ring which may be phenazine (as in phenosafranine), quinoline or pyridine (as in the substances in Tables III and IV), or acridine (as in acriflavine, Table IV). In all the active derivatives of these parent substances described, one cyclic nitrogen is quaternary (phenochloride in phenosafranine, alkohalide in the other compounds). This combination alone is insufficient however, for quinoline methiodide and quinaldine ethiodide have hardly any action (Table IV) whilst phenazine methiodide and *NN'*-dimethyldihydrophenazine [Dickens, 1936] have a powerful catalytic type of action, instead of behaving as inhibitors or anticatalysts of the Pasteur mechanism. The introduction of the basic amino groups as in phenosafranine causes a complete alteration of properties of the *N*-substituted phenazine; the converse effect is seen in the presence of the acidic hydroxyl group in pyocyanine, which like phenazine methiodide depresses glycolysis. Similarly in phenosafranine loss of amino-groups or replacement by hydroxyl results in loss of desensitizing properties. The presence of the amino- (or substituted amino-) group may not be indispensable, however, in all cases; for benzyldinequinaldine ethiodide (Table IV) has a somewhat similar action to the other inhibitors, but this effect is here complicated by a rather strong poisoning effect on respiration. All active compounds contain a fairly long conjugated chain which, except in benzyldinequinaldine ethiodide and in pinacryptol yellow (III) which has a

m-nitro-group, terminates in the amino-group. In this chain, as in the *p*-position to the cyclic N atom, =CH— may be replaced by a nitrogen atom (e.g. compounds II and V).

Reduction of phenazine methiodide and of a derivative of quinoline ethiodide.

Phenazine methiodide (8.01 mg.) was ground with 4.0 ml. *N*/20 HCl until it was mostly dissolved and 1.5 ml. of the greenish yellow suspension were then transferred to a conical Warburg manometer vessel containing in 2 side bulbs 8 mg. Pt black in 0.5 ml. water. The vessel was filled with hydrogen and shaken nearly to absorption equilibrium in a bath at 37.5°. On tipping in the Pt, H₂ was taken up and the absorption was: 60 min. 160 μl.; 120 min. 190 μl.; 180 min. 206 μl., when it became exceedingly slow. Calculated for 1 mol. H₂ per mol. phenazine methiodide 208.5 μl. The solution became a very pale greenish colour and on standing in the air turned orange (cf. the oxidation of *N*-methyl-dihydrophenazine [Dickens, 1936] to a red aposafranone).

p-Dimethylaminophenylazomethine-2-quinoline ethiodide (II), 2.53 mg., dissolved with warming in 1.0 ml. water, were placed in the side bulbs of a Warburg manometer filled with H₂. The main part contained 10 mg. Pt black in 2 ml. water. H₂ uptake: 30 min. 186 μl. (colourless or very pale yellow); 200 min. 261 μl.; 240 min. 262 μl. (both exps. were corrected for a slow residual H₂ absorption by Pt). Calc. for 2H₂ per mol. dye = 264 μl. H₂.

This exp. shows that the uptake of 1 mol. H₂ is accompanied by decoloration of the dye: this stage is reversible, shaking with air reoxidized the dye to give the original violet colour. With the entry of the second H₂ the colour is not restored by shaking with air.

The substances described in this paper therefore behave on mild reduction like the methiodide of nicotinic acid amide [Karrer & Warburg, 1936] and the coenzymes of dehydrogenation and fermentation.

Relationship of the inhibitors to cell catalysts.

As already pointed out, it is probable that these substances depend for their action on specific poisoning (or anticatalysis) of certain cell enzymes. Warburg [1926] has given a similar explanation of the inhibition of the Pasteur reaction by ethyl isocyanide, which has a special affinity to certain heavy metal catalysts in "model" experiments.

The substances described in the present paper resemble ethyl isocyanide in their effects on cell metabolism, but have an even higher affinity for the enzymes concerned in the Pasteur reaction, as is shown by their 100-fold lower active concentration. Whilst the nature of these enzymes remains unknown it is clear that the discussion of mechanism of their poisoning cannot be carried far: at the same time the use of specific inhibitors has often in the past provided a means of analysis of complicated enzyme actions. It is proposed to study the effects on various isolated enzyme systems of members of this group of substances with this end in view.

Until quite recently it was not known that substances of general chemical character resembling in any way that already described for the inhibitors were concerned in cell-catalysis. Warburg *et al.* [1936] have however shown that both the hydrogen-transporting coenzyme and the fermentation coenzyme contain as an integral part of the molecule nicotinic acid amide, and in a very recent paper [Warburg & Christian, 1936] it has been shown that the methiodide of the latter compound on reduction yields a substance of identical spectroscopic and reducing properties with the reduced active group of these coenzymes.

The chemical similarity of the substances described in this paper (e.g. the *N*-methylpyridine ring, with conjugated chain, of the substance VI) with Warburg's compound suggests that substances of this type may perhaps have a

special affinity for enzymes, possibly those participating in the Pasteur mechanism. In the case of Warburg's coenzymes, an activation of the enzymes results from this combination; may it not be that with these inhibitors the same affinities cause the displacement of the active coenzyme by an inert anticatalyst, with consequent inactivation of the enzyme? Possibly the substances acting in an opposite direction (pyocyanine, phenazine methiodide, dimethyldihydrophenazine) may form similar, but active, complexes with the enzyme. Whether this is in fact the actual mechanism or not remains to be proved, but some such type of action is suggested by the specificity and the low active concentration of these substances.

Weil-Malherbe [1935], working in this laboratory, has found that simple dibasic acids (glutamic and maleic acids) strongly influence the Pasteur mechanism in certain tissues. It is probable that this is also primarily an interaction with tissue enzymes; its mechanism is being further examined.

Action of a sarcoma-producing quinoline derivative on metabolism.

Browning *et al.* [1933; 1936] have described the production of sarcoma in mice following a single injection of an aqueous solution of 2 (*p*-aminostyryl)-6- (*p*-acetamidobenzamido)quinoline methoacetate; i.e. of a compound closely related in chemical constitution to the 2-styrylquinoline derivatives found to inhibit the Pasteur mechanism as described above.

This resemblance caused us to study the effect of Browning's compound ("styryl 430") on tissue metabolism: the results are shown in Table V. The substance is freely soluble in water, but addition of salts to form Ringer solution caused precipitation of all but a small fraction; the freshly prepared semi-colloidal suspension thus provided was used for manometric experiments; a slow absorption by the tissue occurred as shown by the colour.

The most marked effect was observed with rat brain cortex. Corresponding with the slow absorption, the effect on metabolism is a gradual one, except anaerobically, when the anaerobic glycolysis is increased from the start. It is known that permeability is in some cases increased by anaerobic conditions [see Dickens & Greville, 1935]. In this tissue the respiration rose slowly (Table V), while the aerobic glycolysis similarly increased until it became about equal to the enhanced anaerobic glycolysis caused by the dye. This effect was shown by 7×10^{-5} *M* suspension, but was insignificant with a further 10-fold dilution, at least in a 4-hour experiment.

Guinea-pig brain cortex showed only the increased respiration, and this soon fell off when the same concentration as is active in rat brain was used. The reason for this difference is unknown. With the highly glycolysing growing tissues mouse yolk-sac and Jensen sarcoma of rat, the anaerobic glycolysis was increased at first in the former but not in the latter; increase of respiration was not observed and of aerobic glycolysis was very slight compared with rat brain.

Although we are unable to account for these differences, the effect with rat brain shows that under favourable conditions styryl 430 has a powerful effect in increasing respiration and glycolysis which resembles that of the other quaternary salts of 2-quinoline derivatives already described, together with some additional activities peculiar to itself and perhaps associated with the physical properties of this compound of high molecular weight. Whilst the effect on respiration is similar to the lyotropic action of K, Cs and Rb [Dickens & Greville, 1935] except that it is more slowly produced, the anaerobic glycolysis is much increased with the styryl compound whereas with KCl it falls. Brilliant cresyl blue [Dickens, 1936] caused a similar increase of $Q_{\text{g}}^{\text{N}_2}$, which however occurred in both

Table V. *Action of "styryl 430" (Browning).*

Concentration 7×10^{-5} *M* as suspension, unless otherwise stated.
Bicarbonate-glucose-Ringer solution pH 7.4. 37.5°.

Tissue	Hours of Exp.	With dye			Control		
		Q_{O_2}	$Q_G^{O_2}$	$Q_G^{N_2}$	Q_{O_2}	$Q_G^{O_2}$	$Q_G^{N_2}$
Rat brain cortex	1	-12.1	+ 3.2	+31.2	-11.9	+ 1.6	+12.7
	2	-14.5	+ 6.1	+27.8	-12.3	+ 0.0	+13.2
	3	-20.4	+15.2	+24.6	-13.7	+ 0.5	+11.5
	4	-23.0	+24.7	+24.0	-13.7	+ 0.9	+12.5
(Four other exps. gave similar result with 7×10^{-5} <i>M</i> suspension.)							
(Same exp.: conc. 7×10^{-6} <i>M</i> : $\begin{cases} Q_{O_2} \\ Q_G^{N_2} \end{cases}$		(1) -13.6	(2) -13.5	(3) -14.7	(4) -14.2		
		+ 2.6	+ 1.6	+ 2.1	+ 2.6		
Guinea-pig brain cortex	1	-17.3	+ 2.7	+17.6	-10.4	+ 2.5	+17.0
	2	-11.0	+ 4.1	+14.9	- 9.1	+ 4.2	+13.1
	3	- 4.2	+ 2.5	+16.5	- 8.3	+ 3.6	+13.7
(Another exp. gave similar result, $Q_G^{O_2}$ increasing from +2.6 to +2.8 only, after 5 hours' exp.)							
Mouse yolk-sac	1	-16.2	+ 5.5	+33.0	-18.5	+ 6.6	+25.4
	2	-13.4	+ 7.7	+26.0	-16.5	+ 7.2	+23.4
	3	- 9.8	+10.3	+21.6	-14.9	+ 8.6	+24.0
(Another exp. gave similar result.)							
Jensen rat sarcoma	1	-10.5	+24.0	+30.4	-10.5	+22.4	+30.4
	2	-10.9	+23.2	+29.0	- 9.6	+20.0	+32.0
	3	-11.5	+25.5	+33.6	-10.8	+23.6	+34.0
	4	- 8.7	+26.5	+31.8	- 8.5	+22.0	+36.0

rat and guinea-pig brain: 2:4-dinitrophenol [Dodds & Greville, 1934] increases respiration and glycolysis of tumour tissues, but in this material styryl 430 has little effect. Although alterations of permeability might be advanced to explain these phenomena, such an explanation is too vague to explain the different behaviours of anaerobic glycolysis towards KCl and styryl 430, for example, as well as the species differences reported above: the lack of sufficient data on permeability enables almost any problem of cell metabolism to receive a theoretical interpretation based on assumed alterations of permeability. The ingenious theories of Dixon [1936] appear to us to suffer from such a disadvantage.

In styryl 430 sarcoma production is seen to be associated with a special action on metabolism, whereby in favourable cases the anaerobic glycolysis of a "resting" tissue may be made to rise to the level found in rapidly growing embryonic and tumour tissues. The aerobic glycolysis increases more slowly, so that at one stage (e.g. 3rd hour, 1st Exp., Table V) the metabolism passes through a stage where it appears the same as for many tumours.

Unlike other cancer-producing agents which have been shown to have a direct influence on metabolism (arsenic, cyanide) styryl 430 induces experimental cancer readily in a high percentage of the animals used [Browning *et al.*, 1933; 1936], and at the same time it has the advantage over the carcinogenic hydrocarbons of being sufficiently soluble for metabolic effects to be produced *in vitro*. The rapidity of action on metabolism (at least with brain tissue) suggests a primary direct influence of the substance itself or of its reduced form rather than of some product formed from it by a slow reaction within the tissues. It is therefore peculiarly fitted for the study of the metabolic changes associated with induction of neoplastic growth.

The precipitation of styryl 430 by tissue fluids has been shown by Browning *et al.* [1936] to lead to the formation of a depot containing the dye at the site of injection, which remains for several months; this may determine the property of tumour induction following a single injection, which is seen with this substance. The other substances described in this paper are more soluble under these conditions, and as yet have not led to the production of malignant growths following a single injection. These animal experiments are being continued and extended.

SUMMARY.

The negative dyestuff phenosafranine inhibits the Pasteur mechanism in animal tissues, the active concentration being in the region of 10^{-5} *M*. The negative potential is not alone sufficient to account for this action, since other negative dyes did not act similarly on metabolism. On the other hand a group of derivatives of pyridine, acridine and quinoline, containing pentavalent nitrogen and a conjugated chain terminating usually in a basic grouping, had essentially the same action on metabolism as phenosafranine, though some members were more poisonous.

The activity of this group is ascribed to an anticatalytic action on certain cell enzymes. Their general chemical similarity to the Warburg pyridine-body, the active group of the H-transporting and fermentation coenzymes, suggests that they may displace the coenzymes by preferential adsorption on the colloidal carrier, forming an inactive complex; but this remains to be proved.

The sarcoma-producing "styryl 430" of Browning, in concentration similar to phenosafranine increases respiration, aerobic and anaerobic glycolysis of rat brain very strongly; its effect on other tissues was less marked. These observations, and their possible relationship to experimental carcinogenesis, are briefly discussed.

To the following who have given valuable material and much helpful advice I would like to express my thanks: Prof. G. R. Clemo and Dr R. Raper (University of Durham), Dr Frances M. Hamer (Kodak Research Laboratories). I am specially indebted to Prof. C. H. Browning (Glasgow University) for kindly supplying a tested sample of purified styryl 430.

REFERENCES.

- Ashford & Dixon (1935). *Biochem. J.* **29**, 157.
 Browning, Cohen, Cooper, Ellingworth & Gulbransen (1933). *Proc. roy. Soc. B* **113**, 300.
 — Gulbransen & Niven (1936). *J. Path. Bact.* **42**, 155.
 Bumm & Appel (1932). *Hoppe-Seyl. Z.* **210**, 79.
 Clemo & Hook (1936). *J. chem. Soc.* 608.
 Dickens (1935, 1). *Nature, Lond.*, **135**, 762.
 — (1935, 2). *J. Soc. chem. Ind.* **54**, 1117.
 — (1936). *Biochem. J.* **30**, .
 — & Greville (1935). *Biochem. J.* **29**, 1468.
 Dixon (1936). *Nature, Lond.*, **137**, 742.
 — & Holmes (1935). *Nature, Lond.*, **135**, 995.
 Dodds & Greville (1934). *Lancet*, i, 398.
 Karrer & Warburg (1936). *Biochem. Z.* **285**, 297.
 Lipmann (1934). *Biochem. Z.* **274**, 329.
 Quastel & Wheatley (1932). *Biochem. J.* **26**, 2169.
 Warburg (1926). *Biochem. Z.* **172**, 432.
 — & Christian (1936). *Biochem. Z.* **285**, 156.
 — & Griesse (1936). *Biochem. Z.* **279**, 143.
 Weil-Malherbe (1935). *J. Soc. chem. Ind.* **54**, 1116.