CLXXXVII. STUDIES IN THE BIOCHEMISTRY OF MICRO-ORGANISMS.

L. RAVENELIN (3-METHYL-1:4:8-TRIHYDROXY- XANTHONE), A NEW METABOLIC PRODUCT OF *HELMINTHOSPORIUM RAVENELII* CURTIS AND OF *H. TURCICUM* PASSERINI.

BY HAROLD RAISTRICK, ROBERT ROBINSON AND
DOUGLAS ELWOOD WHITE.

*From the Division of Biochemistry, London School of Hygiene and Tropical
Medicine, University of London, and the Dyson Perrins Laboratory,
Oxford University.*

(Received 10 June 1936.)

IN a recent series of communications from these laboratories we have recorded the isolation from the dried mycelium of several different species of *Helminthosporium*, when grown on a synthetic medium containing glucose as the sole source of carbon, of a number of hitherto unrecorded polyhydroxyanthraquinones [Charles *et al.* 1933; Raistrick *et al.* 1933; 1934]. These mould metabolic products included helminthosporin (2-methyl-4:5:8-trihydroxyanthraquinone) from *H. gramineum* Rabenhorst and *H. catenarium* Drechsler; catenarin ($\alpha\beta$ -(hydroxymethyl)-1:5:8-trihydroxyanthraquinone) from *H. catenarium* Drechsler, *H. gramineum* Rabenhorst and *H. velutinum* Link; cynodontin (2-methyl-1:4:5:8-tetrahydroxyanthraquinone) from *H. cynodontis* Marignoni, *H. euchlaenae* Zimmermann, and *H. avenae* Eidam, and tritisporin (6(or 7)-(hydroxymethyl)-1:3:5:8-tetrahydroxyanthraquinone) from *H. tritici-vulgaris* Nisikado. In some cases these polyhydroxyanthraquinones are present in surprisingly large amounts, comprising in the case of *H. gramineum* about 30 % of the weight of the dried mycelium.

During the investigation of some 40 species of *Helminthosporium* it was found that, in the cases of *H. Ravenelii* Curtis and *H. turcicum* Passerini, although solvent extraction of their dried mycelia yielded considerable amounts of a yellow crystalline material, this was not a polyhydroxyanthraquinone. This substance, for which the name *ravenelin* is proposed, has now been purified and characterized as 3-methyl-1:4:8-trihydroxyxanthone, thus affording the first recorded instance of the production by any of the lower fungi of a polyhydroxyxanthone. Two naturally occurring hydroxyxanthones have been previously examined, namely (a) euxanthone (1:7-dihydroxyxanthone), which occurs as the glucoside mangiferin in the leaves of *Mangifera indica* L., the Indian mango tree, and, coupled with glucuronic acid, as euxanthic acid in the pigment "piuri", or Indian yellow, prepared from the urine of cows fed on mango leaves, and (b) gentisin (the 3-methyl ether of 1:3:7-trihydroxyxanthone) which is present probably as a glucoside in the rhizomes of *Gentiana lutea* L.

Nisikado [1929] divided the genus *Helminthosporium* into two subgenera on morphological grounds, i.e. the subgenus *Eu-Helminthosporium* and the subgenus *Cylindro-Helminthosporium*. It is perhaps significant that of the 40 species

examined biochemically by us which are common to the 24 species examined morphologically by Nisikado, all those species (with one possible exception) which we have found to produce polyhydroxyanthraquinones are included by Nisikado in the subgenus *Cylindro-Helminthosporium*, whilst those species which fail to produce polyhydroxyanthraquinones, including *H. Ravenelii* and *H. turcicum* which produce a polyhydroxyxanthone, are included in the subgenus *Eu-Helminthosporium*.

H. Ravenelii Curtis (syn. *H. hoffmanni* Berkeley and Curtis, *H. tonkinense* Karsten & Roumeguère, and *H. crustaceum* Hennings), although of very little economic importance, has become one of the best known members of the genus *Helminthosporium* because of its conspicuousness and wide occurrence in many of the warmer regions of the world. It was first described in 1848 by Curtis who noted its abundant distribution in North and South Carolina as a parasite on the grass *Sporobolus indicus* R. Since that time it has been reported by many different mycologists, on *Sporobolus indicus* growing in different parts of North and South America and China. A full account of it is given by Drechsler [1923].

H. turcicum Passerini (syn. *H. inconspicuum* Cooke & Ellis) is the cause of a disease of maize (*Zea Mays* L.) commonly known as "leaf blight" and less frequently as "white blast". It appears to have been first noticed in 1876 in Italy by Passerini. Leaf blight disease of maize is of almost universal occurrence, having been reported throughout the maize belt of the United States, in New South Wales, Japan, India, South Africa and Russia. *H. turcicum* has also been reported on cultivated sorghum by Butler [1918] in India, Egypt and China and by other authors in different parts of the world. A full account of it is given by Drechsler [1923].

Chemistry of ravenelin. The intensely yellow crystalline substance, M.P. 267–268° (corr.), was found to be homogeneous; it is optically inactive and has pseudo-acidic characters. The composition, $C_{14}H_{10}O_5$, was established by elementary analysis, mol. wt. determinations (cryoscopic in camphor) and by the preparation and analysis of derivatives such as the triacetyl-, tribenzoyl- and trianisoyl-ravenelins and ravenelin dimethyl and trimethyl ethers.

These experiments also demonstrated the absence of the carboxyl group confirmed by the failure to effect esterification with the help of the usual acid catalysts.

It is difficult to formulate a trihydroxy-compound, $C_{14}H_{10}O_5$, as a derivative of naphthalene, anthracene or phenanthrene, and yet a polycyclic system was clearly indicated. Hence, especially as the function of two oxygen atoms remained undecided and lactonic character was not evinced, we suspected a xanthone structure. A substance of the composition $C_{14}H_{10}O_5$ might be a trihydroxymethylxanthone. Direct evidence of the presence of a methyl group was obtained by means of the Kuhn-Roth micro-method.

According to the conditions of the experiment, ravenelin affords a dimethyl or a trimethyl ether when it is treated with diazomethane in alcoholic-ethereal solution; the dimethyl ether can be further methylated by means of methyl sulphate and sodium hydroxide in aqueous acetone solution.

The trimethyl ether is inert towards alkalis and acetic anhydride and its most interesting property is its basic character. It is a rather strong oxonium base and a *ferrichloride*, $C_{17}H_{17}O_5 \cdot FeCl_4$, was isolated and analysed.

This observation gave very strong support to the view that ravenelin is a trihydroxymethylxanthone. Confirmation was obtained by the isolation of the ethyl ether of a substituted phenylxanthidrol after reaction between *O*-trimethyl-ravenelin and phenyl magnesium bromide.

Degradation of ravenelin proved to be a very difficult operation, but resorcinol was obtained by fusion with potassium hydroxide. The structure could therefore be expressed as (I) or (II).

Eventually it was found that the pyrone ring could be opened as the result of submission of ravenelin trimethyl ether to the action of sodamide in boiling xylene solution. The effect was to remove a carbon and an oxygen atom and to introduce two hydrogen atoms. It was evident that the reaction had gone beyond the usual first stage of the Haller process (scission of —CO—R affording —CONH_2 and HR) and that, in all probability, the fission of a xanthone ring occurred in accordance with the scheme:

The presumed substituted diphenyl ether was a homogeneous oil, but fortunately it afforded a characteristic crystalline *tetrabromo-derivative*. It will be noticed that structures of the types I and II are derived from one and the same diphenyl ether and as the number of plausible formulae was limited we turned to synthesis in order to decide between them. In this we confined ourselves to substances in which the hydroxyl groups in the trihydroxytoluene nucleus were in *o*- or *p*-relation because ravenelin in alkaline solution is readily oxidized by atmospheric oxygen. The isomeric ethers III, IV and V were prepared by appropriate modifications of Ullmann's method from *m*-methoxyphenol and the requisite dimethoxybromotoluenes.

The isomerides III and V proved to be crystallizable but IV was liquid and identical with the substance obtained from ravenelin trimethyl ether by interaction with sodamide. This important conclusion followed a careful comparison of the tetrabromo-derivatives. It follows that ravenelin is VI or VII.

The analytical evidence at our disposal did not distinguish between these possibilities although the formation of a dimethyl ether might be held to support VI because VII contains two hydroxyls in the *o*-position to the carbonyl. However, it is probable that the inactivation of hydroxyl by neighbouring carbonyl is due to chelation and theoretically this implies that one carbonyl group can inhibit the methylation of only one hydroxyl group at a time.

Actually VII must be the correct formula because we have been able to synthesize the trimethyl ether of VI and this substance, although very similar to, is different from *O*-trimethylravenelin.

The condensation of 3-bromo-2:5-dimethoxytoluene with 2-hydroxy-4-methoxybenzoxynitrile in the presence of potassium hydroxide or ethoxide and copper bronze at 200° affords 2:5-dimethoxytoluene, 2:4-dimethoxybenzoxynitrile (!) and 2-cyano-3'-methyl-5:2':5'-trimethoxydiphenyl ether (VIII) which is hydrolysed by boiling aqueous barium hydroxide to the corresponding acid. Hot phosphoryl chloride then effects the ring-closure to 3-methyl-1:4:6-trimethoxyxanthone (IX).

It is of interest to compare the constitution of ravenelin (VII) with that of helminthosporin (X) because the methyl group and two of the hydroxyl groups are identically situated in these two metabolic products of species of *Helminthosporium*.

In cynodontin (XI), a congener of helminthosporin, the fourth hydroxyl corresponds to the third of ravenelin. It is not known whether cynodontin is an oxidation product of helminthosporin or whether the latter is derived from the former by reduction, but a Dakin-type oxidation of XI followed by reduction could yield VII. This particular hypothesis is naturally a pure speculation and is mentioned only as an example of the kind of relation that may exist between these natural products.

EXPERIMENTAL.

Cultures. The culture of *Helminthosporium Ravenelii* Curtis (L.S.H.T.M. Cat. No. Ag. 138) used throughout this work was purchased from the Centraalbureau voor Schimmelcultures, Baarn, Holland, in June 1933. It was received by them in November 1929 from Nisikado, who isolated it in Japan from *Sporobolus* sp. (strain No. 216). The culture of *H. turcicum* Passerini (L.S.H.T.M. Cat. No. Ag. 124) used was purchased from Baarn in May 1932. It was received by them in November 1929 from Nisikado, who isolated it in Japan from *Zea Mays* L. (strain No. 164).

A. *Cultural conditions and course of metabolism.*

35 litres of Czapek-Dox medium of the following composition were used: glucose, 50 g.; NaNO_3 , 2.0 g.; KH_2PO_4 , 1.0 g.; KCl , 0.5 g.; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.5 g.; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g.; distilled water, 1 litre. This was distributed in 350 ml. amounts in 100 one-litre conical flasks and sterilized. After sterilization a measured quantity, previously determined, of sterile N NaOH solution was added to each flask to bring the pH of the medium to 8.0. Each flask was then sown with the same volume of a suspension in sterile Czapek-Dox solution of a pure culture of *H. Ravenelii* grown on Czapek-Dox agar slopes for 2–3 weeks. The inoculated flasks were incubated at 24° and the mixed contents of 5 flasks were analysed after varying incubation periods. The methods used for treatment of the mycelium are described in section C. The results obtained are given in Table I in which the weights recorded are those obtained from 5 flasks in each case. After 69 days' incubation the remaining 65 flasks were taken off and the figures given in Table I for this incubation period are calculated from the results obtained.

Table I.

Incubation period in days	% glucose in metabolism solution by polarimeter	Wt. of mycelium g.	Wt. of crude ravenelin g.	M.P. of crude ravenelin $^\circ\text{C}$.	Wt. of crude fat g.
21	4.66	2.42	Trace	Very indefinite	0.21
28	4.09	5.16	0.15	253–6	0.33
35	2.82	11.77	0.24	258–263	1.10
42	2.48	12.41	0.30	255–8	1.40
49	2.10	12.25	0.32	252–5	1.18
61	1.50	10.67	0.37	254–6	1.15
66	0.61	16.58	0.38	255–7	2.33
69	0.64	15.23	0.34	259–261	2.91

B. *Effects of temperature and pH on yield of ravenelin.*

A batch of Czapek-Dox medium of the composition given above was made up, distributed in 350 ml. quantities and sterilized. After sterilization some of the flasks were inoculated with *H. Ravenelii* without further treatment (pH of medium = 4.2) while some were adjusted to pH 6.0 and the remainder to pH 8.0 before inoculation. Five flasks at each pH were then incubated at 24° , 5 at 27° and 5 at 30° , in all cases for 52 days. The results obtained are summarized in Table II and refer to yields from 5 flasks in all cases.

Table II.

Initial pH	Incubation temperature $^\circ\text{C}$.	Residual glucose by polarimeter %	Wt. of mycelium g.	Wt. of crude fat g.	Wt. of crude ravenelin g.	M.P. of crude ravenelin $^\circ\text{C}$.
4.2	24	2.21	12.1	1.51	1.72	255–8
6.0	24	1.43	14.1	1.84	1.08	260–1
8.0	24	0.90	15.2	2.50	0.50	261–3
4.2	27	0.90	13.8	1.71	1.71	258–61
6.0	27	1.25	13.4	1.12	1.20	246–50
8.0	27	1.33	14.1	0.90	0.86	246–51
4.2	30	2.47	10.1	0.40	1.28	249–58
6.0	30	2.56	9.0	0.29	0.83	249–56
8.0	30	1.76	11.2	0.25	0.73	252–56

It is clear from these figures that the optimum conditions for the production of ravenelin by *H. Ravenelii* on Czapek-Dox medium are an incubation tem-

perature of 24–27° and an initial *pH* of 4.2. However, since ravenelin, in quantities sufficient for its chemical investigation, had already been prepared before these findings were available, no use was made of them for the large-scale preparation of ravenelin except for the preparation of one batch (see Table III, batch 6).

C. Preparation of ravenelin.

Several batches of ravenelin have been prepared and in each case the medium (adjusted before inoculation to *pH* 8.0 except batch 6, Table III, in which the initial *pH* was 4.2), cultural conditions etc. employed were the same as are described in section A. 100 flasks were used in each batch. At the end of the incubation period the flasks were similar in appearance, the mould growth having a crumpled dark grey to buff surface, with a brown or red margin and a khaki-brown reverse, while the metabolism solution was red-brown in colour. The mycelium was filtered from the metabolism solution, squeezed, washed with cold water, pressed and dried in a vacuum oven at about 40°. It was ground to a fine powder in a coffee mill and extracted for 1 working day with light petroleum (B.P. 40–50°) to remove most of the fat and sterols. The light petroleum-extracted mycelium was dried, re-ground and exhaustively extracted under a reflux condenser with boiling chloroform. (Ethyl ether is almost as efficient.) The crude ravenelin separating from the chloroform (or ether) extracts was filtered, washed, dried and weighed. The solvent was removed from the chloroform (or ether) mother-liquors and the residue was extracted with a small amount of light petroleum (40–50°). The undissolved residue consisted, as did the material separating from the original light petroleum extract, mainly of ergosterol together with small amounts of ravenelin. The combined light petroleum extracts yielded, on evaporation, considerable amounts of crude fat.

Further extraction with acetone or methyl alcohol of the chloroform-extracted mycelium yielded small amounts of mannitol.

The experimental details of 6 batches are summarized in Table III in which all figures refer to 100 flasks.

Table III.

Batch	Incubation period in days	% glucose in metabolism solution by polarimeter	Wt. of mycelium g.	Wt. of crude ravenelin g.	Wt. of crude fat g.	Wt. of crude ergosterol g.
1	52	0.90	307	10.0	53.0	1.75
2	53	0.92	307	10.0	49.6	1.31
3	56	0.63	324	11.2	63.8	2.58
4	67	0.90	300	8.9	80.5	2.02
5	67	0.47	295	11.0	69.1	3.97
6	68	0.94	266	24.8	20.4	4.46

The M.P. of the crude ravenelin was in all cases about 260°.

Isolation of ravenelin from H. turcicum. 100 flasks (35 litres) of Czapek-Dox medium (see p. 1307), initial *pH* 4.2 unadjusted, were sown with *H. turcicum* and incubated at 24° for 60 days; residual glucose by polarimeter, 1.52%; weight of dried mycelium, 273 g. The ground mycelium was extracted as described in section C for *H. Ravenelii*, giving 10.6 g. of a mixture of crude fat and crude ergosterol and 0.75 g. crude ravenelin, M.P. 258–261°, alone or mixed with ravenelin from *H. Ravenelii*.

Chemistry of ravenelin.

Ravenelin (batch 1 *ex H. Ravenelii*) had M.P. 265–266° (corr.) raised to 267–268° (corr.) by crystallization from acetone-chloroform as microscopic needles and prismatic needles (found: C, 65.3, 65.3, 65.3, 64.8; H, 4.0, 3.8, 3.9, 4.0%;

mol. wt. (cryoscopic in camphor), 258, 255) or pyridine (found: C, 65.5; H, 4.0. $C_{14}H_{10}O_5$ requires C, 65.1; H, 3.9%; mol. wt., 258). Found on sample *ex H. turcicum*: C, 65.3; H, 4.2%. On oxidation with chromic acid (Kuhn-Roth method) evidence of the presence of one side-chain methyl group was obtained (found: 89.9, 77.3% of $1C_2H_4O_2$). The intensely yellow substance gives yellow solutions in organic solvents in most of which it is sparingly soluble. It is freely soluble in cold pyridine and hot acetic acid, moderately readily soluble in hot alcohol and acetone and sparingly soluble in hot benzene and light petroleum.

It is insoluble in boiling aqueous sodium acetate but dissolves in hot aqueous sodium carbonate to a brownish yellow solution. The yellow solution in oxygen-free aqueous NaOH quickly becomes deep brown on shaking with air. The ferric chloride reaction in alcoholic solution is an intense dull greenish brown. On heating the orange-yellow solution in sulphuric acid it becomes redder and deeper in tone and then orange-brown.

Ravenelin itself does not appear to form an oxonium salt in contact with concentrated hydrochloric acid.

O-Triacetylravenelin. A mixture of ravenelin (2 g.), acetic anhydride (20 ml.) and pyridine (0.3 ml.) was heated on the steam-bath for 4 hours. The derivative isolated in the usual manner crystallized from acetic acid in slender, colourless needles, m.p. 204–205° (corr.) (2 g.). (Found: C, 62.5; H, 4.2%. $C_{20}H_{16}O_8$ requires C, 62.4; H, 4.2%.) A determination of acetyl groups by the Kuhn-Roth oxidation process indicated the presence of three acetate residues and the side-chain methyl group. (Found: 90.2, 89.8% of $4C_2H_4O_2$.) The substance was optically inactive in chloroform solution. It is sparingly soluble in cold alcohol and benzene but readily soluble in the hot solvents. The material crystallized from acetic acid dissolved in cold acetone and quickly separated again in the form of hair-fine needles.

O-Tribenzoylravenelin. Ravenelin (0.5 g.) was benzoylated in pyridine solution by means of benzoyl chloride (2.5 ml.), at first in the cold and eventually by boiling the mixture for 2 min. The product crystallized on cooling and was collected, washed and crystallized from pyridine. It separated as colourless, elongated, microscopic, prismatic needles with pointed ends, m.p. 255° (corr.). (Found: C, 73.6; H, 3.9%. $C_{35}H_{22}O_8$ requires C, 73.6; H, 3.9%.) The derivative is sparingly soluble in boiling alcohol, moderately readily soluble in hot acetic acid and readily soluble in warm pyridine.

O-Trianisoylravenelin. This derivative was prepared like the benzoate. It crystallized from pyridine and then from acetic acid as colourless, flat, friable needles, m.p. 216–218° (corr.). (Found: C, 69.3; H, 4.5%. $C_{38}H_{28}O_{11}$ requires C, 69.1; H, 4.3%.) The substance was insoluble in aqueous alkalis, and ferric chloride-negative, so that apparently it contained no free hydroxyl group; it gave consistently low values in methoxyl estimations.

Action of nitric acid on ravenelin. Vigorous oxidation occurred under most conditions but the only products isolated were oxalic acid and a sparingly soluble nitro-derivative obtained in insufficient amount for investigation.

Tribromoravenelin. When ravenelin (0.2 g.) was brominated in hot acetic acid by means of bromine (1 g.) the derivative crystallized on cooling and was recrystallized from acetic acid in the form of long, thin, deep yellow, opaque splinters, not possessing a definite melting-point. (Found: Br, 48.0%. $C_{14}H_7O_5Br_3$ requires Br, 48.4%.)

Fusion of ravenelin with alkalis and other experiments. Only unchanged material was obtained as the result of attempted reduction of ravenelin by means of hydriodic acid and phosphorus at 180° or by distillation over zinc dust in a

stream of hydrogen. The attempted oxidation of the triacetate by means of chromic acid in acetic acid solution was also fruitless.

A mixture of ravenelin (3 g.), sodium hydroxide (10 g.), potassium hydroxide (10 g.) and water (20 ml.) was heated in a nickel crucible (aluminium block at 240–275°) for 2 hours and finally for 15 min. at a higher temperature (block at 340°, melt at 250°). The product was dissolved in water, acidified, saturated with ammonium sulphate and extracted with ether. The syrup isolated in this way, which gave a strong fluorescein reaction test for resorcinol, was treated with methyl sulphate and aqueous sodium hydroxide in excess, and the resulting oil isolated by means of ether. It had B.P. 210–211°/736 mm. (corr.) (yield, 0.3 g.). (Found: C, 69.9; H, 7.5%. Calc. for $C_8H_{10}O_2$: C, 70.0; H, 7.3%.) The substance was recognized as resorcinol dimethyl ether by its odour and B.P. (authentic specimen, B.P. 209–214°/736 mm.) and by the preparation of the dibromo-derivative, M.P. 142–143° (corr.) alone or mixed with an authentic specimen. The other products of the alkali fusion could not be isolated in a pure condition.

Methylation of ravenelin. Direct processes using methyl sulphate and alkalis gave unsatisfactory results.

(a) A suspension of ravenelin (3 g.) in absolute alcohol (200 ml.) was treated with diazomethane (from 10 g. of nitrosomethylurea, method of Arndt & Amende [1930]) for 48 hours at room temperature. The solid products were isolated and treated again in exactly the same way. The solvent was then removed by distillation and the residue extracted with hot alcohol (some insoluble residue). The cooled solution deposited long, yellow, hair-like needles, M.P. 285–287° (corr.) (0.15 g.). (Found: C, 66.7; H, 5.0; MeO, 21.8%. $C_{16}H_{14}O_5$ requires C, 67.1; H, 4.9; 2MeO, 21.7%.)

(b) The crude dimethyl ether (0.3 g.) was dissolved in a little acetone and methyl sulphate (5 ml.) added. The mixture was then treated alternately with aqueous NaOH and methyl sulphate until the aqueous alkaline solution was no longer intensely yellow in colour. The acetone was then evaporated and the solid collected (0.25 g.) and crystallized from alcohol. It separated as colourless prisms, M.P. 178–179° (corr.).

(c) The same substance was obtained by the action of diazomethane on ravenelin in alcohol containing a little water. Ravenelin (2 g.) in 99.5% alcohol (300 ml.) was treated with diazomethane (from 20 g. of nitrosomethylurea using undried ether). After keeping for 48 hours at the room temperature the excess of diazomethane was decomposed by acetic acid and the solvents removed. The solids obtained on addition of water were collected (1.95 g.) and dissolved in alcohol (0.2 g., residue). On cooling the solution deposited colourless prisms, M.P. 178–179° after recrystallization from alcohol. (Found: C, 68.4, 68.1; H, 5.7, 5.4; MeO, 31.9%. $C_{17}H_{16}O_5$ requires C, 68.0; H, 5.4; 3MeO, 31.0%.) The procedure described under (b) above was applied to the material in the mother-liquors and in this way 0.9 g. of the trimethyl ether was recovered.

The derivative dissolves in cold conc. HCl to an intense orange solution, the colour being discharged by the addition of water or alcohol.

The hydrobromide crystallized as orange-yellow needles when the substance was dissolved in acetic acid containing hydrogen bromide. This trimethyl-ravenelin was unchanged after treatment with hydroxylamine, semicarbazide, boiling aqueous sodium hydroxide or barium hydroxide or alcoholic potassium hydroxide and acetic anhydride under various conditions.

Ferrichloride. Trimethylravenelin (0.15 g.) was dissolved in conc. HCl and a solution of $FeCl_3$ (0.3 g.) in conc. HCl added. The derivative crystallized from

acetic acid containing HCl and FeCl_3 as red prisms, M.P. $174\text{--}175^\circ$ (corr.). (Found: C, 40.8; H, 3.8%. $\text{C}_{17}\text{H}_{17}\text{O}_5\text{FeCl}_4$ requires C, 40.8; H, 3.4%.)

Nitrodimethylravenelin. After the addition of HNO_3 (1.5 ml. sp.gr. 1.42) to a solution of trimethylravenelin (0.1 g.) in acetic acid (5 ml.), the nitro-derivative crystallized (0.1 g.) in a few minutes in the cold. It separated from acetic acid in flat, elongated, rather irregular prisms, M.P. $224\text{--}226^\circ$ (corr.). (Found: C, 58.4; H, 4.0; N, 4.4%. $\text{C}_{16}\text{H}_{13}\text{O}_7\text{N}$ requires C, 58.0; H, 3.9; N, 4.2%.) The derivative is orange in mass and pale yellow by transmitted light under the microscope. It is insoluble in aqueous NaOH but its alcoholic solution becomes intensely greenish brown on the addition of ferric chloride and this reaction confirms the loss of a methyl group indicated by the analytical results.

Action of sodamide on trimethylravenelin. Very little reaction occurred when the experiment was conducted in boiling benzene solution during $8\frac{1}{2}$ hours. A mixture of trimethylravenelin (2.5 g.), powdered sodamide (10 g.) and xylene (50 ml.) was refluxed (bath at $145\text{--}150^\circ$) for 7 hours. Ice was added to the cooled mixture and the xylene layer separated, dried and distilled. After the removal of the solvent there remained a nitrogen-free oil (1.5 g.) which distilled at 0.5 mm. (bath at 195°). (Found: C, 70.2; H, 6.6; MeO, 33.9%. $\text{C}_{16}\text{H}_{18}\text{O}_4$ requires C, 70.0; H, 6.6; 3MeO, 34.0%.) This *trimethoxymethyldiphenyl ether* could not be induced to crystallize, but it was characterized and identified by means of its tetrabromo-derivative, M.P. 152° , which is described below in connexion with the synthetic specimen of unambiguous constitution.

Ethyl ether of phenyldihydro-O-trimethylravenelin (ethyl ether of a phenyltrimethoxyxanthidrol). A solution of trimethylravenelin (0.3 g.) in benzene (10 ml.) was added to one of phenyl magnesium bromide (bromobenzene, 2.4 g.; magnesium, 0.4 g.; ether, 8 ml.) and the mixture was refluxed for 4 hours. The mass was decomposed with water, diluted with benzene and the benzene layer separated and extracted with successive small volumes of conc. HCl until these extracts were pale in colour. The orange-yellow aqueous acid solutions were neutralized with ammonia and the precipitated solid was collected (0.35 g.) and boiled with alcohol. This process ethylated the carbinol and the colourless ethyl ether crystallized in twinned, flat, rhombic plates, M.P. 166° (corr.). (Found: C, 73.6; H, 7.3; H, 6.4, 6.5%. $\text{C}_{25}\text{H}_{26}\text{O}_5$ requires C, 73.9; H, 6.5%.) On treatment with cold aqueous HCl the ether was hydrolysed and an orange-yellow solution of the corresponding xanthylum salt was obtained.

Synthetical experiments bearing on the constitution of ravenelin.

2:5-Dimethoxy-4-bromotoluene. 2:5-Dihydroxy-4-bromotoluene [Clark, 1892] was methylated by means of methyl sulphate and NaOH in aqueous acetone solution. The dimethyl ether crystallized from alcohol in polyhedra, M.P. $90\text{--}91^\circ$ (corr.). (Found: Br, 34.4%. $\text{C}_9\text{H}_{11}\text{O}_2\text{Br}$ requires Br, 34.7%.)

2:5:3'-Trimethoxy-4-methyldiphenyl ether (III). 2:5-Dimethoxy-4-bromotoluene (11.6 g.), *m*-methoxyphenol (6.2 g. purified as described by Perkin *et al.* [1926]) and powdered potassium hydroxide (2.8 g.) were dissolved in a little alcohol and the solution heated (bath at $200\text{--}220^\circ$) for 7 hours. At first alcohol distilled and later a little of the bromo-compound and water. The product was isolated by means of ether and distilled, B.P. $192\text{--}195^\circ/2.5$ mm. (4.8 g.). The distillate solidified and the substance crystallized from methyl alcohol as stout, colourless prisms, M.P. $72\text{--}74^\circ$ (corr.). (Found: C, 70.1; H, 6.6%. $\text{C}_{16}\text{H}_{18}\text{O}_4$ requires C, 70.0; H, 6.6%.)

The *tribromo-derivative* was prepared by the addition of bromine to a moderately concentrated solution of the ether in acetic acid. It crystallized from methyl

alcohol in colourless needles, m.p. 130°. (Found: C, 37.5; H, 3.2; Br, 46.7%. $C_{16}H_{15}O_4Br_3$ requires C, 37.6; H, 2.9; Br, 47.0%.)

2:5:3'-Trimethoxy-3-methyldiphenyl ether (IV). This substance was prepared in exactly the same way as the isomeride last described, but starting from 2:5-dimethoxy-3-bromotoluene [Kohn & Steiner, 1931] (11.6 g.). 5 g. of a colourless oil, b.p. 165–166°/0.5 mm. were obtained. (Found: C, 70.1; H, 6.5%.)

The *tribromo-derivative* was obtained under the following conditions. The trimethoxymethyldiphenyl ether (0.1 g.) was dissolved in acetic acid (1.5 ml.) and bromine (0.5 ml.) added gradually with stirring. After 2 min. the product (0.15 g.) was isolated after the addition of aqueous sodium bisulphite. The derivative crystallized from methyl alcohol in colourless needles, m.p. 133° (corr.). (Found: C, 37.4; H, 3.3; Br, 47.2%. $C_{16}H_{15}O_4Br_3$ requires C, 37.6; H, 2.9; Br, 47.0%.)

The characteristic *tetrabromo-derivative* is obtained when an excess of bromine is added to a mixture of the ether with about three times its volume of acetic acid and, if the scale of the operation is so small that the liquid does not become warm as the result of the reaction, the mixture is heated to 50° for 1 min. After allowing to cool, aqueous sodium bisulphite is added and the precipitate collected and crystallized from methyl alcohol, in which the substance is very sparingly soluble, and from ethyl alcohol. It crystallizes in well-shaped, colourless prisms, m.p. 152°. (Found: C, 32.7; H, 2.6; Br, 54.6%. $C_{16}H_{14}O_4Br_4$ requires C, 32.6; H, 2.4; Br, 54.3%.) This tetrabromo-compound was prepared from the product of decomposition of ravenelin trimethyl ether with sodamide and had m.p. 152°, alone or mixed with the synthetic specimen. The crystal forms and solubility relations of the two specimens were also identical. The colourless solution in concentrated sulphuric acid becomes successively pink, yellow, brown and reddish brown on heating.

4:5:3'-Trimethoxy-2-methyldiphenyl ether (V). This was prepared like the isomerides from 3:4-dimethoxy-6-bromotoluene [Heap *et al.* 1927; Erdtmann, 1933] and had b.p. 160–162°/0.2 mm. The distillate crystallized from alcohol, separating as colourless prisms, m.p. 68–69° (corr.). (Found: C, 70.6; H, 6.0%.)

Tetrabromo-derivative. This was prepared by the action of an excess of bromine on a cold solution of the ether in thrice its weight of acetic acid.

The product crystallized from methyl alcohol, in which it was more readily soluble than the isomeride described above, in colourless, elongated prisms, m.p. 115°. (Found: Br, 54.7%. $C_{16}H_{14}O_4Br_4$ requires Br, 54.3%.) It may be noted that considerable depressions of melting-point were observed on mixing the various tribromo- and tetrabromo-derivatives described in this memoir.

Intermediates for the synthesis of 1:4:6-trimethoxy-3-methylxanthone (VI). These intermediates have all been previously described, but the methods of preparation have been improved.

m-Bromotoluquinone [Claus & Jackson, 1888] (80 g., m.p. 93°) was obtained by oxidation of crude 4:6-dibromo-*o*-cresol [Kohn & Segel, 1925] (266 g.) by means of chromic acid (200 g.) in acetic acid (1 litre) and water (700 ml.) at 60° (maximum). The quinone (60 g.) was reduced by means of stannous chloride (100 g., hydrated) in conc. HCl (100 ml.) and water (200 ml.) at 80° (maximum). The quinol separated on cooling and was collected and directly methylated by means of methyl sulphate (100 ml.) and 25% aqueous NaOH in methyl alcohol (200 ml.). After isolation there were obtained 44 g. of 3-bromo-2:5-dimethoxytoluene, b.p. 133°/13 mm.

2-Hydroxy-4-methoxybenzaloxime [cf. Späth & Klager, 1934] (73 g., m.p. 140°) was obtained from 2-hydroxy-4-methoxybenzaldehyde [Tiemann & Parrisius, 1880; method of A. M. & R. Robinson, 1932] (66 g.) by oximation (hydroxylamine hydrochloride, 50 g.) in pyridine (200 ml.) for 1 hour on the steam-bath. When the oxime (69 g.) was refluxed with acetic anhydride (350 ml.) for 3 hours, an acetyl derivative was produced and this was isolated and hydrolysed by heating

with 10% aqueous NaOH (300 ml.) during 15 min. on the steam-bath. On acidification of the solution 2-hydroxy-4-methoxybenzonitrile (44 g., m.p. 179°) separated and was purified by crystallization from aqueous methyl alcohol.

2-(2':5'-Dimethoxy-m-tolyloxy)-4-methoxybenzoic acid (related to VIII). Potassium (3 g.) was dissolved in alcohol (30 ml.) and the solution mixed with 2-hydroxy-4-methoxybenzonitrile (11.4 g.), 3-bromo-2:5-dimethoxytoluene (18 g.) and copper bronze (0.3 g.); the whole was then heated at 200° for 8 hours. The neutral products were isolated by means of ether and distilled, giving:

(a) 2.5 g., B.P. 105–170°/12 mm. (nearly all at 105–110°), B.P. 218–220°/751 mm. This was identified as 2:5-dimethoxytoluene [cf. Erdtmann, 1933].

(b) 2.7 g., B.P. 120–170°/0.15 mm., crystallizing on cooling and separating from methyl alcohol as long, colourless needles, m.p. 96.5°. (Found: C, 66.4; H, 5.6; N, 9.1%. Calc. for $C_9H_9O_2N$: C, 66.3; H, 5.5; N, 8.6%.) This was identified as 2:4-dimethoxybenzonitrile and afforded 2:4-dimethoxybenzoic acid, m.p. 108°, on hydrolysis with baryta.

(c) 2.0 g., B.P. 178–188°/0.13 mm., solidifying on cooling and separating from alcohol, light petroleum (B.P. 80–100°) or aqueous acetic acid as short colourless prisms, m.p. 94–95°; analysis indicated that this consisted essentially of 2-cyano-5:2':5'-trimethoxy-3'-methyldiphenyl ether (VIII), but it was contaminated with a small proportion of a substance of lower carbon content.

(d) 0.3 g., B.P. 188–200°/0.13 mm., largely the same nitrile as (c).

The formation of 2:4-dimethoxybenzonitrile is the result of a curious reaction and the substance was also obtained for purposes of comparison by methylation of 2-hydroxy-4-methoxybenzonitrile by means of methyl sulphate and 10% aqueous NaOH in methyl alcoholic solution. The purified substance had m.p. 96.5°, alone or mixed with the reaction product mentioned above.

The fractions (c) and (d) afforded the desired acid on hydrolysis. The cyanotrimethoxymethyl-diphenyl ether (100 mg.) was refluxed with barium hydroxide (10 g.) and water (10 ml.) for 12 hours. The cooled solution was acidified with HCl, the precipitate collected and the acid redissolved in aqueous NaOH and reprecipitated. It was then crystallized from alcohol, separating in the form of microscopic, elongated, rhombic prisms, m.p. 208° (corr.) (20 mg.). (Found: C, 64.2; H, 5.9%. $C_{17}H_{18}O_6$ requires C, 64.1; H, 5.7%.)

1:4:6-Trimethoxy-3-methylxanthone (IX). The carboxytrimethoxymethyl-diphenyl ether (40 mg.) was boiled with phosphoryl chloride (7 ml.) for 3 min. The deep yellowish brown solution was mixed with water and made alkaline with NaOH, and the solid was collected (30 mg., m.p. 155°). The substance separated from alcohol as tiny rods, m.p. 157° (corr.). (Found: C, 68.0; H, 5.5%. $C_{17}H_{16}O_5$ requires C, 68.0; H, 5.3%.) The melting point was depressed on admixture with *O*-trimethylravenelin, but, although different from this substance, the synthetic product exhibits a general resemblance to it.

The deep yellow solution in conc. HCl gives a yellow precipitate on the addition of a concentrated solution of $FeCl_3$ in the same solvent.

SUMMARY.

The dried mycelia of the fungi *Helminthosporium Ravenelii* Curtis, a plant pathogen parasitizing the grass *Sporobolus indicus* R., and *Helminthosporium turcicum* Passerini, the causative agent of leaf blight disease of maize, *Zea Mays* L., have been shown to contain considerable amounts of a hitherto undescribed methyltrihydroxyxanthone, for which the name ravenelin is proposed. Ravenelin, $C_{14}H_{10}O_5$, has been shown to be 3-methyl-1:4:8-trihydroxyxanthone, and a number of derivatives and breakdown products are described. The optimum cultural conditions for the production of ravenelin by *H. Ravenelii* have been determined.

We desire to express our indebtedness to Mr W. K. Anslow for much technical assistance during the preparation of crude ravenelin.

REFERENCES.

- Arndt & Amende (1930). *Z. angew. Chem.* **43**, 444.
Butler (1918). *Fungi and Diseases in Plants*. Calcutta.
Charles, Raistrick, Robinson & Todd (1933). *Biochem. J.* **27**, 499.
Clark (1892). *J. Amer. chem. Soc.* **14**, 569.
Claus & Jackson (1888). *J. prakt. Chem.* **38**, 321.
Drechsler (1923). *J. agric. Res.* **24**, 688, 712.
Erdtmann (1933). *Proc. roy. Soc. A* **143**, 177.
Heap, Jones & Robinson (1927). *J. chem. Soc.* 2021.
Kohn & Segel (1925). *Mh. Chem.* **46**, 661.
—— & Steiner (1931). *Mh. Chem.* **58**, 92.
Nisikado (1929). *Ber. Ohara Inst.* **4**, 111.
Perkin, Ray & Robinson (1926). *J. chem. Soc.* 945.
Raistrick, Robinson & Todd (1933). *Biochem. J.* **27**, 1170.
—— ——— (1934). *Biochem. J.* **28**, 559.
Robinson, A. M. & Robinson, R. (1932). *J. chem. Soc.* 1439.
Späth & Klager (1934). *Ber. dtsh. chem. Ges.* **67**, 859.
Tiemann & Parrisius (1880). *Ber. dtsh. chem. Ges.* **13**, 2365.