113. COMPARISON OF THE GLOBOGLYCOID, CRYSTALBUMIN, SEROGLYCOID AND SEROMUCOID FRACTIONS OF NORMAL HORSE SERUM

BY C. RIMINGTON AND M. VAN DEN ENDE

From the National Institute for Medical Research, Hampstead, N.W. 3

(Received 29 April 1940)

LARGELY as a result of his studies of the albumin fraction of serum, Sørensen [1930] developed the view that the soluble proteins comprise reversibly dissociable component systems, from which by suitable adjustment of conditions a whole series of fractions may be obtained, varying progressively in such characteristics as solubility, carbohydrate content etc. Sørensen's investigations remain a monument of precision and detail and together with his deductions form a classic monograph in protein chemistry. It is of particular significance that though all his albumin fractions were crystalline, this afforded no guarantee of their chemical "purity", since, as he showed, their solubility was not independent of the amounts of solid phase introduced into the system. His most readily soluble albumin fraction contained as much as 50 times the quantity of carbohydrate present in the least soluble fractions, intermediate fractions possessing intermediate values. The tendency of different proteins to dissociate into their component systems was shown to vary. Serum albumin presents rather an extreme case, being very readily dissociable, whereas in egg albumin the tendency to dissociation is considerably less.

The work of Hewitt [1934; 1936; 1937, 1; 1938; 1939], in which attention was focused upon the carbohydrate contents of his serum protein fractions, led him to postulate the existence of three proteins in the albumin group, viz. globoglycoid, crystalbumin and seroglycoid. Crystalbumin contained no carbohydrate, whilst the other two, as their names imply, were relatively rich in bound carbohydrate. Hewitt's crystalbumin was obtained by several successive recrystallizations, at the isoelectric point and half-saturation with ammonium sulphate, of crude crystalline albumin from native serum. Globoglycoid is the name he gave to a protein fraction, described as a globulin, which separates when twice crystallized albumin from crude serum is brought to pH 7.0 and half-saturated with ammonium sulphate. It is recorded as containing about 7.5% of carbohydrate (calculated as glucosaminodihexose). Seroglycoid, the best preparation of which contained 10–11% of carbohydrate, was the protein left in the mother liquors from the albumin crystallizations and was apparently purified by removing the remainders of other proteins either by heat coagulation, or salt precipitation, or both.

If Hewitt's conclusions are correct, then the crystalline specimens of albumin which still contain carbohydrate must be regarded as admixed with either one or both of the other proteins. It appeared to us that the validity of this hypothesis might be conveniently tested by applying the anaphylactic technique in addition to chemical methods, and the investigation originally planned was broadened to include also the seromucoid fraction, a study of which, by one of us, has been presented in the preceding paper [Rimington, 1940].

Plan of the experiments

Our operations have been so planned that in one main experiment we pursued a scheme of fractionation leading to ever purer globoglycoid, with a main albumin fraction as a by-product. These operations were carried out upon the proteins from 5 l. of horse serum freed from lipins by the Hardy-Gardiner process, and have been repeated on smaller quantities of native serum. In the second main experiment untreated serum was used, and the fractionation scheme adopted achieved an ever increasing purification of the serum albumin, since, after every recrystallization, it was brought to the necessary salt and pH conditions to remove a further crop of "globoglycoid". It will be seen that we found the formation of globoglycoid to be an inexhaustible process. From the filtrates of the first albumin crystallization, in each case, seroglycoid fractions were prepared, as detailed below, and subsidiary experiments starting with fresh lots of native serum afforded us our seromucoid samples, the method of preparation being based upon the investigations reported in the accompanying communication.

Details concerning other methods employed may be summarized as follows. Protein concentrations: calculated from the N content (micro-Kjeldahl) and the figure of 15.6 % for the N content of serum albumin [Adair & Robinson, 1930]; checked by measurement of refractive index by Zeiss dipping refractometer. Carbohydrate (hexose) content of proteins: determined by the orcinol method [Rimington, 1940]. Optical rotatory power: measured in triple field instrument using Hg green ($\lambda = 5461 \text{ \AA}$). Dialysis of protein solutions: carried out against distilled water at 0° in "Visking" (cellulose) sausage skins. Precipitation of fractions, adjustments of pH etc.: in every instance reagents added dropwise to the mechanically stirred solutions, final pH being checked electrometrically.

Details of fractionation scheme no. 1

In this instance it was desired to obtain a globoglycoid preparation in as nearly pure form as possible. The crude precipitate obtained at pH 7.0 and half-saturation with $(\text{NH}_4)_2\text{SO}_4$ was therefore reprecipitated under the same conditions four successive times, attention being paid each time to the material left in the mother liquors. In order to obviate any difficulties in preparation that might be caused by the presence of lipins associated with the crude albumin, the serum proteins were freed from lipins by the Hardy-Gardiner process.

5 l. of filtered normal horse serum were chilled and added in the cold room with constant agitation to 15 l. of cold alcohol. Next day the precipitate was removed on large folded filter papers, washed with cold absolute alcohol, and then transferred to 6 l. of cold anhydrous ether. Exhaustion with dry ether followed in the usual manner. To reconstitute the protein solution, the resulting powder was dissolved in 5 l. of 0.85 % NaCl and filtered from a small sediment.

The globulin fraction was precipitated by adding saturated $(\text{NH}_4)_2\text{SO}_4$ until 52 % saturation was reached. After keeping overnight in the cold room, the solution was filtered on large Büchner funnels and the precipitate washed with 52 % saturated $(\text{NH}_4)_2\text{SO}_4$. One-half of the entire crude globulin was purified by successive reprecipitations at half-saturation with $(\text{NH}_4)_2\text{SO}_4$, followed by prolonged dialysis at 0° against 0.85 % of NaCl.

To the filtrate and washings from the globulin precipitate (8.4 l.) N acetic acid (109.2 ml.) was added dropwise, until a faint turbidity appeared. Crystallization occurred rapidly and was brought to completion in the usual way by further addition of $(\text{NH}_4)_2\text{SO}_4$ and acetic acid. The crude albumin was collected

at the centrifuge and redissolved in distilled water to a volume of 1 l. Recrystallization was effected by $(\text{NH}_4)_2\text{SO}_4$ and *N* acetic acid as before, the crystal mass being filtered off and washed by sodium acetate—ammonium acetate-acetic acid mixture [Adair & Robinson, 1930]. The filtrate from these two operations served for the preparation of seroglycoid (see later).

Hewitt's directions for the separation of crude globoglycoid were now followed, *N* NaOH being added dropwise to the mechanically stirred albumin solution (1 l.) until pH 7.06 (electrometric) was reached, and then an equal volume of saturated $(\text{NH}_4)_2\text{SO}_4$ which we had also adjusted with NaOH to pH 7.0. No precipitate formed, so to the stirred solution acetic acid, etc. were added and the crystalline albumin recovered at pH 4.7. It may be noted that Hewitt [1938] found it necessary to remove the bulk of the seroglycoid before separation of globoglycoid would occur under the conditions prescribed.

Our recovered crystalline albumin was redissolved in 500 ml. water and treated as before, with the result that at pH 7 and half-saturation with $(\text{NH}_4)_2\text{SO}_4$, a voluminous precipitate of crude "globoglycoid" now separated (Gg 1). This was filtered off and the filtrate adjusted to pH 4.7 to recover the albumin it contained (A). This was twice recrystallized and then dialysed, affording the "main albumin" fraction, and it was noticed that the mother liquors from these two recrystallizations were free from heat-coagulable protein.

The crude globoglycoid Gg 1 was redissolved in 400 ml. water and again brought to pH 7.0 and half-saturated, affording the precipitate Gg 2 and a mother liquor from which a crystalline albumin GA 1 in the form of stout prisms (Pl. 3, fig. 1) was obtained by adjustment to pH 4.7. GA 1 was once recrystallized before dialysis.

The globoglycoid Gg 2 was redissolved in 300 ml. water and the process repeated yielding Gg 3 and crystalline albumin GA 2 from the mother liquor. It was noticed that the suspension of Gg 3 exhibited, on stirring, a silky sheen, suggestive of crystallization or anisotropy of flow. On repetition of the process, therefore, Gg 3 was redissolved in 300 ml. of water, adjusted accurately to pH 7.0 and an equal volume of neutralized $(\text{NH}_4)_2\text{SO}_4$ added drop by drop whilst vigorous mechanical stirring was maintained. The globoglycoid (Gg 4) then separated only slowly and was beautifully crystalline, in the form of needles or long slender prisms (Pl. 3, fig. 2). Gg 4 was redissolved in 200 ml. and the process repeated, the globoglycoid separating (Gg 5) and albumin from the mother liquors (GA 4) both again crystallizing in the respective forms noted above. Since the amount of material available was now becoming small, the purification process was stopped at this stage and all fractions were dialysed till salt-free and then analysed, with the results shown in Table 1.

It will be seen from Table 1 that, at each reprecipitation of the globoglycoid fraction, there was left in the mother liquor a protein, which by adjustment of the pH to 4.7 separated in crystalline form, indistinguishable from albumin, leaving nothing in the filtrate which could be coagulated by heat. The globoglycoid itself became crystalline after the third precipitation at pH 7.0, but these crystals had the form of long needles, or prisms more slender than those of albumin. There is a progressive decrease in the specific rotations of the albumin fractions recoverable from the successive globoglycoid mother liquors, and the figures for the two final materials, globoglycoid 5 and GA 4—the albumin from its filtrate, are identical within the limits of experimental error. The rotatory power of the "main" albumin, separated at the beginning of the experiment, and only twice recrystallized, is appreciably higher. With regard to carbohydrate content, it may be noted that the final globoglycoid, like the final albumin

Table 1. *Analyses of main albumin, globoglycoid (Gg) and albumin recovered from globoglycoid mother liquors (GA) (lipin-free proteins)*

Fractionation 1				
Fraction	Designation	Crystalline form	$[\alpha]_{5461}$	Hexose/ total N
Crude globoglycoid	Gg 1	Amorphous	—	—
Main albumin (twice recrystallized) from mother liquors of Gg 1	A	Hexagonal prisms	-74.27°	0.0075
Once reprecipitated globoglycoid	Gg 2	Amorphous	—	—
Albumin from mother liquors of Gg 2	GA 1	Hexagonal prisms	-73.33°	0.0228
Twice reprecipitated globoglycoid	Gg 3	? Microcrystalline	—	—
Albumin from mother liquors of Gg 3	GA 2	Hexagonal prisms	-73.86°	0.0082
Thrice reprecipitated globoglycoid	Gg 4	Needles or fine prisms	-73.26°	Trace
Albumin from mother liquors of Gg 4	GA 3	Hexagonal prisms	-71.24°	Nil
Four times reprecipitated globoglycoid	Gg 5	Needles or fine prisms	-68.50°	Nil
Albumin from mother liquors of Gg 5	GA 4	Hexagonal prisms	-68.61°	Nil

fraction, was carbohydrate-free. The second GA fraction showed a lower carbohydrate : N ratio than the first, and no carbohydrate could be detected in the later ones.

These considerations strongly suggest that the Gg and GA fractions obtained by this procedure are modifications of the same substance, which may be obtained in part as globoglycoid by crystallization at pH 7.0, the remainder then appearing as the familiar crystalline albumin on readjustment of the filtrate to pH 4.7. It is conceivable that an equilibrium between the two forms of the same protein is established in solution, depending upon the protein, hydrogen ion and salt concentrations.

In addition to this, however, the experimental results suggest that a progressive fractionation was taking place in the direction of albumin fractions with lower specific rotatory powers [cf. Sørensen, 1930]. The main albumin with $[\alpha]_{5461} -74.27^\circ$, which had undergone a relatively limited purification, must be looked upon as intermediate with regard to this property, as the results of the second scheme of fractionation show.

Details of fractionation scheme no. 2

In this instance, 1 l. of native horse serum was employed as starting material and the crude albumin was purified by repeated treatments in half-saturated $(\text{NH}_4)_2\text{SO}_4$, first at pH 7.0 and then, after filtration, at pH 4.7. By this means, a series of globoglycoid fractions was obtained, each derived from the same albumin but in a series corresponding to progressive recrystallizations of the latter. The albumin finally obtained had thus undergone six successive crystallizations at pH 4.7, and five globoglycoid precipitates were obtained, of which the last three were crystalline and formed only slowly in the mechanically stirred solution.

Fig. 1. Crystalbumin from horse serum. $\times 300$.

Fig. 2. Globoglycoid from horse serum. $\times 220$.

All solutions, when dialysed salt-free, were analysed for N and carbohydrate contents and the refractive index and specific rotatory power determined with the following results (Table 2):

Table 2. *Analysis of globoglycoid fractions and final albumin (native serum)*

Fractionation 2				
Fraction	Designation	Crystalline form	$[\alpha]_{5461}$	Hexose/ total N
Globoglycoid from albumin:				
Once crystallized	CGg 1	Amorphous	-57.51°	0.321
Twice crystallized	CGg 2	Amorphous	-65.42°	0.160
Three times crystallized	CGg 3	Needles or slender prisms	-70.26°*	0.102
Four times crystallized	CGg 4	Needles or slender prisms	-66.45°	0.063
Five times crystallized	CGg 5	Needles or slender prisms	-67.78°	Nil
Final albumin	CA	Stout hexagonal prisms	-80.79°	0.0048

* ? High value owing to error.

It will be seen from Table 2 that this scheme of fractionation has resulted in a series of globoglycoid fractions with decreasing carbohydrate content and increasing specific rotatory power. Although CGg 1 had a hexose/N ratio of 0.321, equivalent to about 7.5% of glucosamino-di-hexose, which is similar to the figure quoted by Hewitt as the carbohydrate content of his globoglycoid, the subsequent fractions contained progressively less sugar until finally in CGg 5 the quantity was too small to be determined. The final albumin, recovered after the removal of CGg 5, also contained only a trace of carbohydrate, estimated at 0.005%.

Further experiments upon the relationship between globoglycoid and albumin

Hewitt described globoglycoid as a globulin present in the albumin fraction. One would expect therefore that physical, chemical and immunological differences should exist between these two substances which would serve to differentiate them. Our experiments, on the contrary, suggest that globoglycoid is formed from crystalalbumin under the conditions adopted for its isolation, or exists in equilibrium with crystalalbumin in solution. Had globoglycoid been a true globulin one would have anticipated its precipitation from native serum at the first treatment with half-saturated $(\text{NH}_4)_2\text{SO}_4$. The two substances differed in their crystalline forms, and the different specimens of globoglycoid redissolved much more rapidly when placed in contact with water than did the corresponding albumin fractions, although, after stirring for some time, these also gave absolutely clear solutions. The pH of the redissolved and dialysed globoglycoids remained in the region of 7.0, whilst that of the albumins approximated to 5. The whole behaviour of the two substances was reminiscent of the relationship between an acid and its somewhat more soluble salt. It seems possible that some such relationship may actually exist between these two compounds. Crystalline globoglycoid was also obtained from our albumin samples when $(\text{NH}_4)\text{OH}$ was used throughout instead of NaOH for adjustment of all solutions, so that it cannot be a sodium salt. More probably globoglycoid is a combination of albumin and ammonium ions. In this connexion, one may recall the well-

known behaviour of insulin when combined with salmine whereby the region of least solubility is removed from pH 5.5, the isoelectric point of insulin itself, to pH 7.3.

McMeekin [1939] has recently reported that by acidifying a solution of carbohydrate-free crystalalbumin to pH 4.0 with H_2SO_4 and then concentrating by ultrafiltration, he has obtained about one-half of the material in the form of crystals of an albumin sulphate; the sulphate ion was removable by electro-dialysis. When boiled, dialysed solutions of globoglycoid became opalescent like milk, but, even when salts were added, gave no precipitate comparable with the coarse, curdy coagulum of heat-denatured albumin. With a view to defining more precisely the nature of globoglycoid, we undertook another fractionation experiment, in which sufficiently large fractions of the crystalline material were obtained to render possible the determination of molecular weights by osmotic pressure measurements, the reconversion of globoglycoid into crystalline albumin etc.

Fractionation 3

A quantity of once crystallized albumin (about 80 g. in all) from different batches of normal horse serum was redissolved in water, filtered from insoluble material and recrystallized at half-saturation with $(NH_4)_2SO_4$ and pH 4.7. This material was again dissolved, and taken through a scheme of fractionation essentially similar to that followed in fractionation 2, five lots of globoglycoid being obtained corresponding to the five successive recrystallizations of the albumin fraction.

During the course of these operations, for the purpose of following more closely the relation between globoglycoids and their parent albumins, determinations of optical rotatory power were made upon the various solutions and, since calculation of specific rotations was required, a rapid means of determining protein concentration was desirable, which should be applicable if possible to undialysed solutions.

Such a method was found in the reaction with Folin and Ciocalteu's phenol reagent as employed by Greenberg [1929]. By determining the colour value of albumin solutions of known protein content (micro-Kjeldahl and specific refractive index) the factor 22.81 was obtained for conversion of colour values with our sample of the reagent in terms of tyrosine into protein.

A high concentration of ammonium salts interfered with the phenol reagent method by causing turbidities, but since most of the solutions analysed required a 50- or 100-fold dilution to bring them into the convenient range of protein concentration, the amount of salt carried over into the final determination was too small to result in interference from this cause.

One more consideration must be mentioned. Since the factor for each protein depends upon its tyrosine or phenol content, it was possible that different factors would have to be used for albumin and globoglycoid respectively. Determinations, to be reported in a later section, showed, however, that the tyrosine and tryptophan contents of these proteins were identical, within the limits of the analytical procedure employed [Folin & Marenzi, 1929].

The phenol reagent method, although convenient, is less accurate than the Kjeldahl procedure, and it would also appear from determinations upon solutions before and after dialysis that, in the former case, the values tend to be slightly high [compare Rimington, 1924], resulting in a lower value for specific rotatory power.

The analytical results of this experiment are given in Table 3.

Table 3. *For the preparation of globoglycoid specimens. Analytical data*

Fractionation 3		[α] ₅₄₆₁ before dialysis	[α] ₅₄₆₁ after dialysis
Fraction	Designation		
Globoglycoid fraction from first treatment, separating on standing from mother liquor of immediate precipitate: once recrystallized	Gg 1a	-67.72°	—
Globoglycoid from second treatment	Gg 2	—	-68.22°
The above, twice recrystallized	Gg 2ii	-70.47°	-71.40°
Albumin recovered from mother liquors of these recrystallizations	Gg (2-2ii) alb.	-68.42°	—
Extra globoglycoid from second treatment separating after adjustment of pH from 6.6 to 7.0: once recrystallized	Gg 2ai	—	-72.38°
Globoglycoid from third treatment	Gg 3	-69.80°	-72.47°
Main albumin filtrate after removal of above	A (3)	-71.45°	—
Globoglycoid from fourth treatment	Gg 4	-73.26°	—
Globoglycoid from fifth treatment	Gg 5	-74.54°	—
Main albumin after removal of above and two recrystallizations	Final alb.	-74.11°	-74.64°

Recrystallization of globoglycoid at pH 4.7

This was accomplished using a portion of the specimen Gg 3, the specific rotatory power of which had been determined both before and after dialysis, affording a mean value of -71.14° . To the protein solution was added an equal volume of saturated $(\text{NH}_4)_2\text{SO}_4$ (unneutralized) and the pH adjusted cautiously to 4.7 by the dropwise addition of 2.5% acetic acid to the mechanically stirred solution. The separation of crystals commenced after the addition of 1 ml. of acid and was complete after 3.5 ml. The crystals were stout hexagonal prisms indistinguishable from crystalalbumin, and the specific rotatory power determined upon a 3% solution was -70.89° . Thus, as in the examples noted in fractionation 1, and also in the case of the final globoglycoid and albumin fractions (Gg 5 and Final alb.) of fractionation 3, there was no difference between the specific rotatory powers of the globoglycoid and the crystalline product derived from it or from its mother liquors at pH 4.7. These two materials, obtained at pH 7.0 and at 4.7 respectively, did however exhibit the customary difference, the former being in the form of long, slender prisms or needles, and the latter being indistinguishable from ordinary crystalline albumin.

A determination of the osmotic pressure of globoglycoid at pH 7.0 was kindly carried out for us by Dr Elford and afforded values corresponding to a mol. wt. for the protein of approximately 60,000 uncorrected for ionic effects.

Preparation of seroglycoid and carbohydrate-rich albumin fractions

From a study of Hewitt's papers dealing with the protein fraction he has named seroglycoid, it is clear that several alternative routes are available for arriving at a concentration of the carbohydrate-rich material left in the filtrate

after the separation of crystalline albumin from serum. Thus the second communication [Hewitt, 1937, 1] describes under "Fractionation methods" a separation by means of 2/3 saturated $(\text{NH}_4)_2\text{SO}_4$, and trichloroacetic acid is there stated to give inconsistent results. Later much attention is devoted to the production of fractions rich in carbohydrate by a process involving treatment in the boiling water bath at two different hydrogen ion concentrations. In a subsequent paper [Hewitt, 1937, 2] dealing with sera from various animal species, salting out, trichloroacetic acid precipitation, heat coagulation and combinations of these processes are all employed. For comparison with our own fractions we decided to make material by both the heat coagulation technique and by a scheme of fractionation at the 66, 75 and 100% levels of saturation with $(\text{NH}_4)_2\text{SO}_4$ [see Lustig, 1930]. The latter preparations were used in the anaphylaxis experiments since the application of heat at any stage is undesirable when the preparations are intended for immunological work. Trichloroacetic acid, in our hands, proved to be unsatisfactory. Our preparations, though we make no claim for their homogeneity, should, at least, have contained appreciable quantities of seroglycoid.

The filtrate from fractionation 1, after the albumin had been removed as completely as possible by crystallization at pH 4.7, was brought to pH 6.0 by NaOH and then mechanically stirred in a large flask immersed in a water bath. The solution was maintained at 80° for 15 min., then brought to pH 4.7 and again heated for 15 min. Next day the coagulum was filtered off and the filtrate (11 litres), after being dialysed, was concentrated under reduced pressure to 300 ml. 2 vol. of alcohol were added and the precipitate produced was centrifuged off, and redissolved in water. The solution was filtered from some inorganic matter and then reprecipitated with alcohol. After a further reprecipitation the material was dissolved, dialysed and stored after filtration through a Seitz filter. This material gave an intense Molisch reaction and a purplish-pink biuret reaction.

Another preparation was made from native serum in an analogous manner, but, in order to avoid the lengthy process of concentration under reduced pressure, the filtrate from the albumin was fully saturated with $(\text{NH}_4)_2\text{SO}_4$ and the precipitate thus produced was redissolved and heated at 80° as described above. Trouble was encountered; when attempting to redissolve the alcohol-precipitated material, from the presence of insoluble, lipoid material giving strongly positive sterol reactions [compare Rimington, 1940].

Ammonium sulphate fractionation was carried out on the albumin filtrate derived from 2 l. of serum. By raising the concentration of this salt to 66% saturation a fairly bulky precipitate (Sg 66) resulted which, when redissolved and dialysed, gave a deep brown solution. Further raising of the concentration in the main filtrate to 75% saturation afforded Sg 75, redissolving to a straw-coloured solution, and finally by full saturation there resulted a scanty precipitate (Sg 100) giving a slightly opalescent but almost colourless solution. These fractions were dialysed and their hexose:N ratios determined. For comparison, the analytical results on preparations from two additional sera are also set out (Table 4).

Table 4. *Ratio $\frac{\text{Hexose}}{\text{total N}}$ for ammonium sulphate fractions*

	Sg 66	Sg 75	Sg 100
Serum 1	0.313	—	—
Serum 2	0.453*	0.548	1.370
Serum 3	0.315	0.538	1.370

* Reprecipitated at $\frac{2}{3}$ saturation.

Preparation of seromucoid fraction

Since the proper purification of seromucoid demands much labour and considerable quantities of starting material, preparations were employed which were obtained as follows: 5 l. serum were diluted with 15 l. of water and adjusted to pH 4.7 with acetic acid. The mixture was then heated in a steam-jacketed pan until coagulation was complete and the pH readjusted, if necessary, to 4.7. The coagulum was filtered off and the clear, yellow filtrate concentrated under reduced pressure to a small volume and then precipitated by 2 vol. of alcohol. The precipitate was centrifuged down and washed several times with absolute alcohol, which removes, among other things, much pigment, and then with ether. After resuspension in water, a small insoluble residue was removed, affording a water-clear solution giving an intense Molisch reaction. Analyses indicated a hexose : N ratio of 0.574, corresponding approximately to 11.2 % of glucosaminodihexose, since the N content of seromucoid is only about 13 %.

Determination of tyrosine and tryptophan contents of the various serum protein fractions

A further basis for the comparison of the different fractions was afforded by their tyrosine and tryptophan contents. The Folin & Marenzi [1929] method was used and the following results obtained:

Fraction	Tyrosine	Tryptophan	Hexose/ total N
Main albumin (A) from fractionation 1	4.17	0.29	0.0075
Globoglycoid (Gg 5) five times reprecipitated, according to fractionation 1	4.29	0.30	Nil
Precipitate (Sg 66) formed by $\frac{2}{3}$ sat., with $(\text{NH}_4)_2\text{SO}_4$ after removal of crystalline albumin	4.59	1.04	0.272
Precipitate (Sg 75) formed by $\frac{2}{3}$ sat. after removal of Sg 66	4.05	0.99	0.313

These figures for tryptophan content confirm those of Hewitt, who found 0.26–0.30 % in his rigorously recrystallized albumin samples of low carbohydrate content and values around 1.0 % for the carbohydrate-rich seroglycoid fractions. His tyrosine values are somewhat higher than ours, but this may perhaps depend upon the method of calculation, since the N content of the carbohydrate-rich fractions is appreciably lower than that of the recrystallized albumins. Hewitt, however, used the factor $\text{N} \times 7.0$ for calculation of the protein content of his materials.

IMMUNOLOGICAL INVESTIGATIONS

In the immunological comparison of several protein fractions from one serum, many difficulties arise, which are mainly due to the presence in any one fraction of traces of one or more of the others.

The preparation of antisera, and their absorption to render them specific, are laborious procedures. Nevertheless Hewitt [1937, 3] has succeeded by such methods in demonstrating immunological differences between seroglycoid and crystalbumin from horse serum. He concluded moreover that pure, carbohydrate-free crystalbumin was but poorly antigenic in comparison with seroglycoid or globulin. He attributed the main antigenic function of his crystalbumin

preparations to the presence of traces of seroglycoid and globulin. Dale & Hartley [1916] had previously shown that the sensitivity to the albumin fraction of horse serum requires a longer latent period for its development than sensitivity to horse serum globulins. If a mixture of antigens consisting of albumin, globulin and seroglycoid is used for immunization, it is to be expected that precipitins for the better antigens (e.g. globulin) will develop more rapidly than precipitins for albumin.

The test of the anaphylactic responses of isolated uterine strips from actively sensitized guinea-pigs affords a rapid, simple, and highly specific method for the comparison of the antigenic properties of serum fractions. The method has been successfully employed by Dale & Hartley in a study of the immunological differences between the globulins and albumins of normal horse serum. Hewitt did not succeed in demonstrating immunological differences between his serum fractions by anaphylactic methods. His experiments, however, were performed on the whole animal, and massive, shock-eliciting doses of the antigens were injected. A method using the isolated tissue is better suited to this type of experiment, since desensitization of the whole guinea-pig to several antigens is impracticable, whereas a single sensitized smooth muscle strip suspended *in vitro* can be used for several successive tests with different antigens.

In the majority of our experiments specific sensitiveness was induced by a single intraperitoneal injection, of 2 mg. of a separated protein fraction, into a normal virgin guinea-pig weighing 125–150 g. A latent interval of at least 3 weeks was allowed after the sensitizing injection. In agreement with the findings of Dale & Hartley, an interval of this duration was found to be essential for the development of adequate sensitiveness to our crystalbumin preparations. We have, indeed, in this way repeatedly succeeded in inducing specific sensitivity to crystalbumin, without at the same time producing sensitivity to globulin, and in some experiments with a preparation of very highly purified ten times recrystallized crystalbumin.

In a few cases guinea-pigs were passively sensitized by a single I.P. injection of 0.5 ml. of an antiserum from a rabbit immunized to a single protein fraction.

Groups of guinea-pigs were sensitized to the most highly purified samples of each of the five serum fractions to be compared. These were: (1) the six-times precipitated total globulins from horse serum; (2) the crystalbumin from lipin-free serum after it had been crystallized repeatedly and freed before each crystallization from "globoglycoid" (CA); (3) globoglycoid which had been repeatedly reprecipitated at pH 7.0 in 50% sat. $(\text{NH}_4)_2\text{SO}_4$ (Gg 5); (4) seroglycoid, represented by that fraction of normal horse serum which after complete removal of crystalbumin is precipitated at $(\text{NH}_4)_2\text{SO}_4$ concentrations between 66 and 75% saturation (Sg); (5) seromucoid (Sm).

For the comparison of any two of these fractions, the excised uterine strips, from a guinea-pig sensitized with one of them, were suspended in Locke's solution, according to Dale's [1913] method. One of the uterine strips was then tested for specific sensitiveness to the homologous (sensitizing) antigen, and the other for its ability to react to the other serum fraction under test.

With the possible exception of crystalbumin and globoglycoid, the serum fractions used in these investigations were not immunologically pure. In the comparison of any two serum fractions, therefore, the presence of traces of a third fraction as a common impurity could yield results which might erroneously suggest the existence of some degree of antigenic community between the two fractions. In a large number of the earlier experiments, much confusion resulted from the presence of such impurities. It became necessary therefore to

adopt a technique whereby the sensitiveness to those antigenic fractions which, for any particular experiment, could be regarded as impurities, was specifically neutralized, before the two antigens to be compared were tested. Weil [1914] has shown that specific desensitization is only relative. Sensitized smooth muscle, after reacting specifically with the sensitizing antigen in high dilution, is no longer capable of reacting to the antigen in that dilution, but still reacts to the same antigen in a higher concentration. In the majority of our experiments the test doses were applied at a constant dilution. The plain muscle sensitized with antigen A was first tested with antigen B, freedom of which from A could not be guaranteed. It might give a reaction to the first application of B and thereafter show itself insensitive to a further similar dose of B, but still react to the same dose of the specific antigen A. Such a result of the incomplete purity of several of the antigens must be kept in view in interpreting the results. The method evolved may be illustrated by the results of actual experiments. The interval allowed for sensitization after injection of the antigen is to be understood to have been 3 weeks, unless otherwise indicated.

(1) *Plain muscle from guinea-pigs sensitized to crystalbumin (CA)
or to globoglycoid (Gg)*

The earlier experiments of this group were directed to a comparison between crystalbumin and globoglycoid, which, according to the chemical evidence already presented, seemed to be interconvertible forms of the same protein. The initial responses given by the plain muscle to either of these proteins were, in fact, indistinguishable, whichever had been used for the sensitizing injection. When, however, attempts were made to desensitize the plain muscle to either Gg or CA by increasing and repeated applications of one of them, a phenomenon suggestive of some kind of antigenic disparity between the two presented itself. If, after desensitization to a relatively high concentration of one, the uterus was then tested with the other protein in low concentration, in a number of experiments it gave a pronounced reaction. Fig. 3 illustrates this phenomenon. It shows that the application of 1 mg. of either Gg or CA to the plain muscle, no longer capable of responding to 0.2 mg., resulted in a typical contraction, after which it was insensitive to further application of the larger dose. Each was still however capable of reacting to the other in the original lower concentration. We have records as clear as that shown in Fig. 3, from plain muscle of guinea-pigs sensitized with CA, giving large responses to 0.2 mg. of Gg or of CA, after complete desensitization to 1 mg. of the other. In other cases the phenomenon was less clear. For example, plain muscle from a guinea-pig sensitized with Gg, after being completely desensitized to 2 mg. of CA may give only a small residual reaction with even 2 mg. of Gg.

The explanation of this phenomenon is not easy. A supposition that CA and Gg are essentially the same protein with a common antigenic specificity, as the chemical evidence would lead one to expect, but that our preparations of them are incompletely pure, and that the contaminant is a different additional antigen in each case, is incompatible with the reaction observed; for application of the antigen which had been used for sensitization should remove also the sensitiveness to the other. Such a suggestion is moreover difficult to reconcile with the fact that the phenomenon under discussion is exhibited as clearly and strongly by plain muscle from guinea-pigs sensitized with our purest crystalbumin, which gives no reaction with globulin, as by that from guinea-pigs sensitized with less rigidly purified samples of CA. It would be possible on the other hand

to suppose that CA and Gg are separate sensitizing antigens and that our preparation of each of them contains the other as contaminant. Numerous experiments, however, and especially those in which plain muscle from guinea-pigs

Fig. 3.

Fig. 4.

Fig. 3. Response *in vitro* of the uterine muscle of a guinea-pig sensitized with globoglycoid. 1. First horn. 0.2 Gg, 0.2 mg. globoglycoid; CA, crystalbumin. 2. Second horn.

Fig. 4. Response *in vitro* of the uterine muscle of a guinea-pig sensitized with crystalbumin. 1. First horn. Sm, seromucoid; Sg, seroglycoid; G, globulin; Gg, globoglycoid; CA, crystalbumin. 2. Second horn. H, 0.002 mg. histamine.

Fig. 5. Response *in vitro* of the uterine muscle of a guinea-pig sensitized with crystalbumin. 1. First horn. 0.2 G, 0.2 mg. globulin; Sm, seromucoid; CA, crystalbumin; Sg, seroglycoid; H, histamine. 2. Second horn.

passively sensitized with rabbit antigloboglycoid or anticrystalbumin serum were used, and which will be described subsequently, have shown that Gg and CA cannot be regarded as separate antigens.

Fig. 6. Response *in vitro* of the uterine muscle of a guinea-pig passively sensitized with rabbit anti-albumin serum. 1. First horn. 2. Second horn. The test dose of protein was 0.2 mg. throughout, except in the case of the second application of seromucoid (Sm) to the second horn, when the dose was 1.5 mg. CA, crystalbumin; SG, seroglycoid; H, 0.002 mg. histamine.

Fig. 7. Response *in vitro* of the uterine muscle of a guinea-pig sensitized with seroglycoid. Test dose of 0.2 mg. protein used throughout. 1. First horn. 0.2 G, 0.2 mg. globulin; CA, crystalbumin; Gg, globoglycoid; Sg, seroglycoid; Sm, seromucoid; H, 0.002 mg. histamine. 2. Second horn.

Fig. 8. Response *in vitro* of the uterine muscle of a guinea-pig sensitized with seromucoid. Test dose of 0.2 mg. protein used throughout. 1. First horn. 0.2 G, 0.2 mg. globulin; CA, crystalbumin; Gg, globoglycoid; Sm, seromucoid; H, 0.002 mg. histamine. 2. Second horn. Sg, seroglycoid.

Fig. 9. Response *in vitro* of the uterine muscle of a guinea-pig sensitized with seromucoid. 1. First horn. 0.2 Sm, 0.2 mg. seromucoid. 2. Second horn. 0.2 Hs, 0.2 ml. normal horse serum; H, 0.002 mg. histamine.

Another possible explanation seems to be that CA and Gg, which the chemical evidence represents as different but interconvertible physical forms of the same protein, act as a single sensitizing antigen, so that either form produces the same degree of primary sensitiveness to both, but that the physical differences confer a minor degree of antigenic disparity, such that desensitization to CA will leave sensitiveness to Gg and *vice versa*.

In subsequent experiments we tested the reactions of plain muscle from guinea-pigs, actively sensitized with CA or Gg, first to seromucoid, seroglycoid and globulin, and its reaction to the specific sensitizing antigen, after desensitization to these others. Fig. 4 shows the result of such an experiment. There was a full reaction to Sm, and subsequent application of Sg also evoked a clear and characteristic but slightly smaller reaction. After these responses there was no reaction to globulin (G), so that, if G contained any of the antigens responsible for the responses to Sm and Sg, it contained not more, and probably less, than Sm or Sg. Further applications of these three non-specific antigens completed the desensitization to all, which was less easily effected to Sg than to the others. The plain muscle was then treated with Gg and CA in the standard doses. It still reacted well to either, although the reaction was not in either case as complete as would be expected if the muscle had not been partially desensitized by the earlier treatment with the other antigens; but it was at least as strong with Gg as with CA, the antigen with which the animal was sensitized. The plain muscle was then no longer capable of responding to either CA or Gg. On this evidence it would be reasonable to conclude that CA and Gg, if both could be obtained free from traces of other proteins, will be antigenically indistinguishable.

In the experiment illustrated in Fig. 5 both horns of a uterus actively sensitized to CA were first desensitized to G and Sm. One of the horns was then still capable of responding once with maximal reaction to CA but not to subsequent contact with CA or Sg. The other horn gave a typical reaction to Sg, after which it was no longer able to respond to the same dose of Sg, but application of CA was followed by a definite but submaximal reaction. Fig. 5 can be explained by the supposition that our G, Sm and Sg preparations all contain antigenic constituents in common with the sensitizing antigen CA. The fact that the response to our Sg significantly reduces the sensitivity and subsequent response to CA is probably explicable by the presence in the Sg of a significant contamination with CA which the method used for purifying Sg would not be likely to exclude.

A number of experiments was performed with the plain muscle from guinea-pigs passively sensitized with antiserum from rabbits which had received two courses of immunizing injections of our purest crystalbumin preparation. (The total amount of crystalbumin received by each rabbit during the two courses was 24 mg.) Such antisera could reasonably be expected to contain antibodies, the bulk of which are specific for the main antigenic constituent of the preparation used for immunization (CA). The sensitivity induced by such an antiserum will therefore be predominantly for CA, and sensitivity to the minor antigenic components (impurities) of our crystalbumin preparation will be less striking than after active sensitization with CA, when the respective degrees of sensitivity induced to the minor components of the antigen may be important and even as great as that to the main component. Fig. 6 shows the result of an experiment in which the uterine muscle of a guinea-pig sensitized by the intraperitoneal injection of 1.5 ml. of anticrystalbumin serum was used, and from it can be concluded that Sm contained no appreciable amount of crystalbumin but that our Sg contained a significant CA component.

The plain muscle from guinea-pigs passively sensitized with anticrystalbumin or antigloboglycoid serum was found, moreover, to react equally to both Gg or CA, and application of either of these preparations resulted in desensitization to both. The responses to G, Sg and Sm of plain muscle sensitized to CA or Gg are not unexpected in view of the fact that none of the other fractions could be regarded as purified to the same extent as the crystallizable fractions (CA and Gg). It is indeed evident that our Sg, for example, contained appreciable amounts of crystalbumin; the seromucoid preparation, on the other hand, was apparently free from CA. The majority of results have shown, in addition, that crystalbumin and globoglycoid are antigenically indistinguishable; and only in a number of the earlier experiments have minor immunological differences, the nature of which is at present obscure, been observed between them.

(2) *Plain muscle from guinea-pigs sensitized to seroglycoid (Sg)
or to seromucoid (Sm)*

The responses of uterine plain muscle of guinea-pigs actively sensitized by a single injection of 2 mg. seroglycoid have been variable. In the majority of cases, typical anaphylactic responses have been obtained by contact with any one of five serum fractions available. This finding was not unexpected in the light of evidence which has already been provided that our Sg, owing to its method of preparation, could not be regarded as immunologically pure. It has, however, occasionally been found that sensitiveness to Sg could be induced without at the same time producing demonstrable sensitiveness to CA, Gg or globulin. The result of such an experiment is shown in Fig. 7. From this result it must be concluded that seroglycoid contains an antigen, immunologically distinct from G, CA or Gg. This antigen is, moreover, present also in our Sm preparations, although in significantly smaller amounts than in Sg.

The antigenic difference between Sm and Sg could be demonstrated even more strikingly by the use of plain muscle from guinea-pigs actively sensitized to Sm. Fig. 8 shows the result of a typical experiment in which the uterine plain muscle after desensitization to the standard dose of 0.2 mg. of G, CA and Gg, was incapable of reacting to Sg, and application of Sg did not appreciably reduce the sensitivity to subsequent contact with Sm. Sm, therefore, owes its principal antigenic properties to a component different from the main antigen of Sg.

A few experiments were performed with the plain muscle of guinea-pigs passively sensitized by single intraperitoneal injections of an antiserum from rabbits immunized against our seroglycoid preparation. Such plain muscle was found capable of responding in a typical manner to contact with CA, but after desensitization to this antigen it was still capable of reacting to contact with either Sm or Sg. A single response to Sg was, however, followed by desensitization to both Sm and Sg, whereas the response to Sm resulted in desensitization to this antigen only, leaving the plain muscle capable of reacting to subsequent contact with Sg. The conclusions which can be drawn from these results with passively sensitized plain muscle are identical with those to be drawn from the experiments with actively sensitized material, namely that our Sg and Sm are immunologically different although our preparations of them contain a common antigen. This might be a common impurity, but it seems more probable that our Sg contained traces of Sm.

The observed immunological differences between our Sg and Sm preparations were apparently not brought about by the drastic methods of extraction and

purification involved in the preparation of the Sm. This was proved by the demonstration that sensitivity induced by Sm could be completely neutralized by the application of untreated horse serum (see Fig. 9).

DISCUSSION

We have prepared those fractions which Hewitt has designated "globoglycoid", "crystalbumin" and "seroglycoid" by following as closely as possible the methods he has described. We have also prepared seromucoid for comparison. By applying to normal serum different schemes of fractionation, we have in the one case proceeded to ever purer crystalbumin, by eliminating globoglycoid before each of the six successive crystallizations. In the other case, by subjecting the globoglycoid formed at the first treatment to repetition (three successive times) of the process of precipitation with half-saturated $(\text{NH}_4)_2\text{SO}_4$ at the neutral point, a purification of the globoglycoid was achieved, so that it became crystalline, and it was possible to remove albumin in crystalline condition from the mother liquors after each precipitation. The albumin from which the first crude globoglycoid had been separated was once recrystallized and gave us our main albumin fraction.

Hewitt has claimed that globoglycoid behaves like a globulin, is amorphous and contains carbohydrate, but we have repeatedly succeeded in obtaining it in a crystalline condition. The carbohydrate associated with the crude material was found to decrease with each successive reprecipitation or recrystallization, until finally a product was obtained which was carbohydrate-free, and chemically indistinguishable from crystalbumin (amino-acid content, mol. wt. etc.) and different only in certain physical properties, such as crystalline form, and appearance after heat coagulation, which may depend upon the state of dissociation and combination with anions or cations, as regulated by the *pH* of the medium with which the protein is in contact.

Particular interest attaches to the determination of optical rotatory power made on the materials resulting from the two main schemes of fractionation. It would appear that the crude globoglycoid separating from albumin upon the initial treatment at *pH* 7.0 possesses a relatively high optical rotatory power, but that as this globoglycoid is reprecipitated or recrystallized under the same conditions its optical rotatory power diminishes. The albumin recoverable at *pH* 4.7 from the final globoglycoid mother liquor has the same low value as the globoglycoid itself. When, on the other hand, successive crops of globoglycoid are taken off from the same initial albumin, these show a progressively increasing rotatory power, until finally a globoglycoid is obtained, having a moderately high optical rotatory power, identical with that of the albumin in its mother liquor. These observations can be most readily explained on the hypothesis of mutually interacting component systems as postulated by Sørensen [1930]. As has been pointed out in the experimental section of this paper the carbohydrate which is associated with the crude globoglycoid fraction is lost upon reprecipitation and recrystallization of the latter, just as is that from crude crystalbumin, and both these materials may ultimately be obtained practically carbohydrate-free.

On the immunological side, obvious difficulties of interpretation arise when anaphylactic sensitization and the subsequent testing of the specific sensitivity are carried out with proteins which cannot be regarded as rigidly purified from others having different antigenic specificities. For example, if plain muscle from an animal sensitized to crystalbumin is found to exhibit a minor degree of sensitiveness to globulin, and still to react to the albumin when desensitized to

a larger dose of the globulin, it is obvious that the condition might be attributable either to a trace of globulin in the albumin used for the sensitizing injection, or to the presence of albumin as a minor constituent of the globulin used for testing. The antigenic distinctness of globulin and albumin has been sufficiently established by previous work, and the method used by us for purifying our globulin fraction offers no good guarantee of its purification from the other serum antigens. It seems proper, therefore, to attach little weight, in our interpretations, to the reactions to globulin shown by plain muscle from guinea-pigs sensitized to the other proteins studied. Eliminating these, we are left with evidence of an almost complete community of antigenic specificity between globoglycoid and crystalbumin, corresponding to the chemical evidence that they represent essentially the same protein, probably in different combinations with inorganic ions producing two different points of minimum solubility.

Our seroglycoid preparations were certainly impure, and the method of preparation could hardly be expected to lead to the isolation of a single protein from so complex a mixture, but determinations of carbohydrate content indicated that in these fractions there accumulated a protein, the chemical properties of which are markedly different from those of crystalbumin. It is relatively rich in bound carbohydrate (hexose/N ratio approximately 0.3) and contains more than twice as much tryptophan and slightly more tyrosine than does crystalbumin. The seromuroid preparations were also carbohydrate-rich (hexose/N ratio 0.57). Immunologically, also, there was evidence that our seroglycoid and seromuroid were impure, but that each contains an antigen distinct from any present in the other proteins studied, presumably due to their characteristic glycoproteins, and antigenically distinct with respect to one another. Our results, however, give no indication as to whether the specificity is due to the association of two different protein antigens with a common antigenically indifferent carbohydrate complex (cf. Coghill & Creighton [1938]; Neuberger & Yuill [1940]), or to two distinct carbohydrate haptenes combined with the same protein.

Combining our conclusions with those from earlier work, it appears that the test based on anaphylaxis will detect the presence in normal serum of at least five protein antigens—euglobulin, pseudoglobulin, crystalbumin (existing also in the form of globoglycoid), a glycoprotein in “seroglycoid” and another in “seromuroid”. Our conclusions agree with those of Hewitt in regarding the carbohydrate complex found in crude crystalline serum albumin as associated with other protein constituents than the crystalbumin itself, which when rigidly purified contains no carbohydrate. They differ, in that we find that the fraction which Hewitt termed “globoglycoid”, regarding it as a globulin associated with carbohydrate, is not a globulin in any sense except its precipitation by $(\text{NH}_4)_2\text{SO}_4$ at neutral reaction; on the contrary we find it to be completely crystallizable, interconvertible with crystalbumin by appropriate methods, and similarly free from carbohydrate when pure.

SUMMARY

Methods of preparation of crystalbumin, globoglycoid, seroglycoid and seromuroid from horse serum are described.

Chemical and immunological investigations (anaphylaxis *in vitro*) on the separated fractions have been undertaken. From the results of these it is concluded that crystalbumin and globoglycoid are identical or closely similar. Seroglycoid is found to differ quantitatively from crystalbumin in its carbohydrate and amino-acid contents.

Immunological evidence is provided which suggests that seroglycoid preparations contain a carbohydrate antigen which is distinct from crystalbumin.

Seroglycoid and seromucoid, although both rich in bound carbohydrate, differ from one another when compared by anaphylactic methods.

We wish to express our thanks to Mrs M. E. Adair for kindly supplying several protein solutions which we used for comparison with our own preparations, and for her interest in this investigation.

*We are indebted to Sir Henry Dale for his invaluable and helpful criticism.

REFERENCES

- Adair & Robinson (1930). *Biochem. J.* **24**, 993.
Coghill & Creighton (1938). *J. Immunol.* **35**, 477.
Dale (1913). *J. Pharmacol.* **4**, 167.
— & Hartley (1916). *Biochem. J.* **10**, 408.
Folin & Marenzi (1929). *J. biol. Chem.* **83**, 89.
Greenberg (1929). *J. biol. Chem.* **82**, 545.
Hewitt (1934). *Biochem. J.* **28**, 2080.
— (1936). *Biochem. J.* **30**, 2229.
— (1937, 1). *Biochem. J.* **31**, 360.
— (1937, 2). *Biochem. J.* **31**, 1534.
— (1937, 3). *Biochem. J.* **31**, 1047.
— (1938). *Biochem. J.* **32**, 26.
— (1939). *Biochem. J.* **33**, 1496.
Lustig (1930). *Biochem. Z.* **225**, 247.
McMeekin (1939). *J. Amer. chem. Soc.* **61**, 2884.
Neuberger & Yuill (1940). *Biochem. J.* **34**, 109.
Rimington (1924). *Biochem. J.* **18**, 1297.
— (1940). *Biochem. J.* **34**, 931.
Sørensen (1930). *C.R. Lab. Carlsberg*, **18**, no. 5.
Weil (1914). *J. med. Res.* **30**, 299.