9. STUDIES IN DETOXICATION

9. THE SYNTHESIS OF SOME POSSIBLE BIOLOGICAL OXIDATION PRODUCTS OF SULPHANILAMIDE

BY W. V. THORPE AND R. TECWYN WILLIAMS

From the Physiology Department, Medical School, Hospitals Centre, Birmingham

(Received 19 December 1940)

IN the previous paper [Thorpe & Williams, 1941] we expressed the opinion that 1-amino-2-hydroxy-(II) and 1-amino-3-hydroxy-benzene-4-sulphonamide (V) and 2-oxybenzoxazole-6-sulphonamide (I) might be biological oxidation products of sulphanilamide. Here we describe the synthesis of these compounds. No difficulty was experienced in the preparation of the *o*-aminophenol and benzoxazole derivatives which were obtained at different stages in the same synthesis, but the synthesis of the *m*-aminophenol derivative presented unexpected difficulties:

We first attempted a synthesis of 1-amino-3-hydroxybenzene-4-sulphonamide (V), starting with the easily prepared *m*-nitraniline-sulphonic acid (III). It was hoped to obtain large amounts of 1-nitro-3-aminobenzene-4-sulphonamide (IV), and thence, by diazotization and reduction, the required *m*-aminophenol-sulphonamide. We found, however, after many trials using different chlorinating agents, that the nitranilinesulphonic acid formed little or none of the corresponding sulphonyl chloride. Small amounts of (IV) were eventually obtained in one experiment using chlorosulphonic acid, but this result could not be repeated. This proposed method of synthesis was therefore abandoned. One interesting point which emerged at this stage was that ammonium 1-nitro-3-aminobenzene-4-sulphonate was isolated in a dimorphous state, a pale orange and a red form. The orange form could be easily converted into the red form by gentle warming below 100°. The red form, once formed, was apparently stable.

Success was eventually attained starting with *m*-aminophenolsulphonic acid, which is easily prepared by heating *m*-aminophenol with conc. H_2SO_4 . The synthesis worked smoothly to the sulphonyl chloride stage, whilst the final stages were completed with some difficulty.

Although many workers (see previous paper for literature) state that they have used *p*-hydroxylaminobenzenesulphonamide (VI) in biological work, no mention is made in their papers of its method of preparation or chemical properties. This substance has been prepared by Bratton *et al.* [1939] and we have repeated their process and agree with their description of its chemical properties.

We have also prepared its *N*-acetyl derivative¹ which has m.p. 227° and gives no diazo reaction in agreement with Bratton *et al.* [1939]. We mention this since James [1940] claims to have isolated this acetyl compound from human urine after sulphanilamide medication, and states that the substance he isolated did not melt below 300°. We failed to prepare *p*-hydroxylaminobenzenesulphonamide using Bamberger's [1894] method with zinc dust at low temperatures, and Haworth & Lapworth's [1921] sodium hydrosulphide method for the preparation of arylhydroxylamines.

The phenolic sulphonamides (II) and (V) have not yet been tested for therapeutic activity. Of phenolic sulphonamides which have been tested for therapeutic activity, 1-hydroxy-2-aminobenzene-4-sulphonamide [Tréfouel *et al.* 1937; McLeod, 1938] and *p*-hydroxybenzenesulphonamide [Buttle *et al.* 1936; McLeod, 1938] were inactive, and 1-hydroxy-3-nitrobenzene-4-sulphonamide [McLeod, 1938] was active. Compounds related to our 1-nitro-3-aminobenzene-4-sulphonamide have also been tested and found inactive, namely 1-nitro-4-aminobenzene-3-sulphonic acid [Buttle *et al.* 1936] and 1-nitro-6-aminobenzene-3-sulphonamide [Tréfouel *et al.* 1937].

EXPERIMENTAL

2-Oxybenzoxazole-6-sulphonyl chloride. Benzoxazolone (2-oxybenzoxazole) was prepared from salicylamide and NaOCl according to Graebe & Rostovzeff [1902]. The only modification introduced was that the benzoxazolone was precipitated from the alkaline reaction mixture by neutralizing carefully, with cooling, with conc. HCl instead of by passing CO₂ into the solution. The neutralization with CO₂ may take several hours (on a 20 g. scale) whilst with HCl it can be done in under half an hour. Sodium benzoxazolonesulphonate was prepared according to the patent directions [Friedländer, 1911]. The dry salt (10 g.) was stirred with 12 g. PCl₅. The mass warmed up and became pasty, and after standing half an hour with occasional stirring, it was treated with ice water in the usual manner. The precipitated *chloride* was filtered, washed with water and dried (yield 6 g.). It crystallized from toluene in colourless rhombic plates, m.p. 186–187°.² (Found: C, 36.2; H, 1.8; N, 6.1%. C₇H₄O₄NClS requires C, 36.0; H, 1.7; N, 6.0%.)

2-Oxybenzoxazole-6-sulphonamide (I) was prepared by adding the powdered chloride to strong ammonia. The deposit formed overnight was filtered and dissolved in warm dilute NaOH. On cooling, the *sodium salt* of the sulphonamide separated as needles, which were recrystallized from water. (Found: N, 10.2; Na, 8.4; H₂O, 13.4%. C₇H₅O₄N₂SNa, 2H₂O requires N, 10.3; Na, 8.45; H₂O, 13.2%.) The *sulphonamide* itself immediately separates as needles if an aqueous solution of the sodium salt is acidified. On recrystallization from water, the sulphonamide formed needles, m.p. 263°. (Found: C, 39.4; H, 2.8; N, 13.3%. C₇H₆O₄N₂S requires C, 39.25; H, 2.8; N, 13.1%.)

1-Amino-2-hydroxybenzene-4-sulphonamide (II). 2.8 g. of the foregoing sodium salt were boiled for 20 min. with 25 ml. of 30% NaOH. Crystals of

¹ We experienced some difficulty in this acetylation. Using acetic anhydride as recommended by Bratton *et al.* we persistently obtained a product melting about 160° from which Bratton's acetyl derivative was only obtained in small yield by fractional crystallization. We did not examine the other product in detail except to note that it gave a diazo reaction, and prove that it was not unchanged *p*-hydroxylaminobenzenesulphonamide. It may be the substance responsible for Rosenthal and Bauer's diazo reaction (see Thorpe & Williams [1941]).

² All melting-points are uncorrected.

ERRATA

Volume 35, Nos. 1 & 2

p. 62, line 13 of text

for **1-hydroxy-3-nitrobenzene-4-sulphonamide**

read **1-hydroxy-2-nitrobenzene-4-sulphonamide**

Na_2CO_3 separated in the solution, and were liable to cause violent bumping. The solution was cooled, carefully neutralized with conc. HCl and then made alkaline with solid Na_2CO_3 . The solution was now extracted for 8 hr. with ether in a continuous extractor and on evaporation of the ether the *sulphonamide* was obtained as a crystalline residue (2 g.). The compound was purified by dissolving in hot absolute alcohol and adding sufficient carbon tetrachloride to produce a faint cloudiness; on cooling the sulphonamide separated as clusters of long needles, m.p. 164° . It was soluble in water and alcohol, sparingly soluble in ether and insoluble in CCl_4 . It reduced ammoniacal silver nitrate on warming and gave a purple colour with FeCl_3 turning a deep red in about 30 sec. It slowly caused the formation of methaemoglobin in a 1 in 50 solution of blood at 37° . (Found: C, 38.6; H, 4.3; N, 15.6%. $\text{C}_6\text{H}_5\text{O}_3\text{N}_2\text{S}$ required C, 38.3; H, 4.3; N, 14.9%.) The *hydrochloride* was prepared by adding the requisite amount of conc. HCl to an alcoholic solution of the sulphonamide. On stirring the solution the hydrochloride separated; more alcohol was now added and the solution boiled until the hydrochloride dissolved. Carbon tetrachloride was then poured into the solution until the hydrochloride just began to crystallize and the crystallization was completed at 0° . It formed fine matted needles, which were very soluble in water and which decomposed without melting above 300° . It was purified by crystallization from absolute alcohol and CCl_4 . (Found: C, 32.3; H, 3.9; N, 12.6%. $\text{C}_6\text{H}_5\text{O}_3\text{N}_2\text{S} \cdot \text{HCl}$ requires C, 32.1; H, 4.0; N, 12.5%.)

Treatment of an aqueous solution of the sulphonamide with benzoyl chloride and NaOH gave *1-benzamino-2-benzoyxybenzene-4-sulphonamide*, which formed needles from aqueous alcohol, m.p. 191° . (Found: C, 60.0; H, 3.9; N, 7.4%. $\text{C}_{20}\text{H}_{16}\text{O}_5\text{N}_2\text{S}$ requires C, 60.6; H, 4.1; N, 7.1%.)

1-Amino-3-hydroxybenzene-4-sulphonic acid was prepared by sulphonation of *m*-aminophenol [cf. Friedländer, 1896]. 15 g. of *m*-aminophenol were heated on the water bath with 25 ml. conc. H_2SO_4 and a few drops of fuming H_2SO_4 for an hour. The solid product was broken up in about 150 ml. of water and 10% NaOH added with stirring until a clear solution was obtained. The solution was now filtered, acidified with HCl and set aside to crystallize at 0° . The acid (tablets) was filtered, washed with water and dried over CaCl_2 ; yield 24 g.

Pyridine 1-acetamino-3-acetoxybenzene-4-sulphonate. The above acid (10 g.) was shaken with 25 ml. pyridine and then 25 ml. acetic anhydride added. Addition of the anhydride caused the evolution of heat; the solution was therefore cooled under the tap. The acid dissolved in about an hour, the solution being occasionally shaken to facilitate solution. The mixture solidified on keeping at room temperature overnight. The solid was filtered under suction, pressed and washed with small amounts of methylated spirit. It was recrystallized from spirit, ether being added to facilitate the crystallization. On evaporating the mother liquor from the acetylation mixture *in vacuo*, a small amount of the salt was recovered; yield 15 g. m.p. 170 – 172° . The substance gave no colour with FeCl_3 and no diazo reaction. (Found: N, 7.9%. $\text{C}_{15}\text{H}_{16}\text{O}_6\text{N}_2\text{S}$ requires N, 7.95%.)

1-Acetamino-3-acetoxybenzene-4-sulphonyl chloride. The foregoing pyridine salt (5 g.) was stirred with 5 g. PCl_5 . The mixture soon became pasty and warm. After 15–20 min., the mixture was heated for a few minutes on the water bath. The product was then thrown into ice-water and, on stirring, the *chloride* separated as a white solid (yield almost quantitative). It crystallized from toluene as long rods, m.p. 169° . (Found: C, 41.4; H, 3.4; N, 4.85%. $\text{C}_{10}\text{H}_{10}\text{O}_5\text{NClS}$ requires C, 41.2; H, 3.45; N, 4.8%.)

1-Acetamino-3-hydroxybenzene-4-sulphonamide. The above chloride was added to 5 times its weight of 40% ammonia (40 ml. NH_4OH sp.gr. 0.880 made

up to 100 ml. with water). The chloride gradually dissolved, the solution becoming warm. After an hour the opalescent solution was filtered and evaporated to a thin paste on the water bath. After cooling, the paste was stirred with about 5 vol. water and the sparingly soluble crystalline material collected and washed with water. The *1-acetamino-3-hydroxybenzene-4-sulphonamide* was recrystallized from 50 % aqueous alcohol as colourless needles, m.p. 280° decomp. (The melting-point was observed after raising the temperature fairly rapidly up to about 270°; if heated very slowly decomposition sets in about 250°, observed melting-point may be as low as 265° and it is difficult to get consistent results.) Yield of crude product 30 % weight of chloride. The pure substance gave no diazo reaction. A violet colour with FeCl_3 was obtained with a solution of the amide in 50 % aqueous alcohol. (Found: C, 41.8 %; H, 4.5; N, 12.2 %. $\text{C}_8\text{H}_{10}\text{O}_4\text{N}_2\text{S}$ requires C, 41.75; H, 4.4; N, 12.2 %.)

The mother liquor from the separation of this amide gave on further evaporation crystalline material (yield up to 10 % weight of chloride) rather more soluble than the main product. It crystallized from 30 % aqueous alcohol in apparently homogeneous needles, m.p. 235°. It gave no diazo reaction, but readily gave a violet colour with FeCl_3 with an aqueous solution. On hydrolysis with 2*N* NaOH, it yielded 1-amino-3-hydroxybenzene-4-sulphonamide. C, H and N values (C, 42.6; H, 4.5; N, 11.1 %) gave no clue to the nature of this substance (mixture?) and we did not investigate it further.

The filtrate from the separation of the unidentified substance appeared to contain small amounts of this unidentified substance or the above amide (or both) and also some acetylated 1-amino-3-hydroxybenzene-4-sulphonic acid, since after hydrolysis with 2*N* NaOH some 1-amino-3-hydroxybenzene-4-sulphonamide could be obtained by extraction with ether, and 1-amino-3-hydroxybenzene-4-sulphonic acid (yield up to 10 % weight chloride) separated in tablets on acidifying with HCl.

1-Amino-3-hydroxybenzene-4-sulphonamide (V). 1-Acetamino-3-hydroxybenzene-4-sulphonamide (1 g.) was dissolved in 10 ml. 2*N* NaOH and heated in a boiling water bath for 3½ hr. Water lost by evaporation was replaced at intervals. (Hydrolysis with 2*N* HCl was equally effective.) The mixture was then made neutral or just acid by addition of conc. HCl and finally rendered just alkaline by addition of solid Na_2CO_3 . The solution was extracted with ether in a continuous extractor for at least 8 hr. *1-Amino-3-hydroxybenzene-4-sulphonamide* separated from the extract in clusters of long colourless plates (yield 0.5 g.). A further crop (0.1 g.) was obtained by evaporation of the ether, but the product obtained in this way was liable to turn brown on keeping unless it was immediately purified. The amide was recrystallized by taking up in a small amount of absolute alcohol followed by addition of CCl_4 until cloudy. The amide gradually crystallized in long plates which had m.p. 152° and gave a strong diazo reaction and a weak red-violet colour with FeCl_3 . It was readily soluble in water and alcohol, sparingly soluble in ether, insoluble in CCl_4 . (Found: C, 38.7; H, 4.4; N, 14.6 %. $\text{C}_6\text{H}_8\text{O}_3\text{N}_2\text{S}$ requires C, 38.3; H, 4.3; N, 14.9 %.) On adding conc. HCl to a concentrated aqueous solution, the *hydrochloride* separated in elongated rhombic plates which could be recrystallized from a small volume of 80 % aqueous alcohol. By shaking an aqueous solution of the amide with acetic anhydride, the monoacetyl derivative m.p. 280° already described separated from the solution. The amide did not cause formation of methaemoglobin when incubated with a 1 in 50 blood solution.

1-Nitro-3-aminobenzene-4-sulphonamide (IV). *m*-Nitraniline was sulphonated according to Friedländer [1923] and 1-nitro-3-aminobenzene-4-sulphonic acid isolated as the sparingly soluble sodium salt. Treatment of the dry Na salt or of

the free acid itself with PCl_5 , thionyl chloride or chlorosulphonic acid failed to give any indication of the formation of the sulphonyl chloride except in one case. An account of this isolated case is given, but we must emphasize that this experiment could not be repeated. 20 g. of the dry *m*-nitranilinesulphonic acid were heated on the water bath for 3.5 hr. with 20 ml. of chlorosulphonic acid. The resulting thick liquid was poured into ice-water with stirring. A yellow precipitate separated which was filtered and washed twice with water. After pressing as dry as possible, it was stirred into 200 ml. ammonia (sp. gr. 0.880), in which it immediately dissolved. After 24 hr. at room temperature a deposit (6 g.) of clear light brown prisms of large size separated, having M.P. 213–214°. On evaporating the mother liquor ammonium 1-nitro-3-aminobenzene-4-sulphonate was obtained. This salt crystallized at room temperature as large orange prisms, but if the temperature of crystallization was above room temperature, then a mixture of bright red and orange crystals was obtained. The orange form could be easily converted into the red form on heating in a test tube at about 80°. The red form once formed appeared to be stable. (Found, orange form: C, 30.8; H, 3.8; N, 17.6%; red form, C, 30.5; H, 3.8; N, 17.8%. $\text{C}_6\text{H}_5\text{O}_5\text{N}_3\text{S}$ requires C, 30.6; H, 3.9; N, 17.9%.) The light brown crystals of M.P. 213–214° appeared to be mainly 1-nitro-3-aminobenzene-4-sulphonamide, contaminated with some of the above ammonium salt. The crystals were recrystallized from water and formed yellow prismatic needles, M.P. 215°. (Found: C, 33.3; H, 3.2%. $\text{C}_6\text{H}_7\text{O}_4\text{N}_3\text{S}$ requires C, 33.2; H, 3.25%.)

SUMMARY

The synthesis of 1-amino-2-hydroxybenzene-4-sulphonamide, 2-oxybenzoxazole-6-sulphonamide, 1-amino-3-hydroxybenzene-4-sulphonamide and several related compounds is described. Their preparation was necessitated because they may be biological oxidation products of sulphanilamide.

1-Amino-2-hydroxybenzene-4-sulphonamide is a methaemoglobin-former *in vitro*, but 1-amino-3-hydroxybenzene-4-sulphonamide is not.

We wish to express our gratitude to Prof. W. N. Haworth and Dr M. Stacey, who arranged for the C, H and some of the N estimations to be done by their micro-analyst.

REFERENCES

- Bamberger (1894). *Ber. dtsh. chem. Ges.* 27, 1548.
Bratton, White & Marshall (1939). *Proc. Soc. exp. Biol., N.Y.*, 42, 847.
Buttle, Gray & Stephenson (1936). *Lancet*, 1, 1289.
Friedländer (1896). *Fortschr. Teerfarb.* 3, 59 (D.R.P. 71,229).
— (1911). *Fortschr. Teerfarb.* 9, 148 (D.R.P. 197,496).
— (1923). *Fortschr. Teerfarb.* 13, 242 (D.R.P. 294,547).
Graebe & Rostovzeff (1902). *Ber. dtsh. chem. Ges.* 35, 2751.
Haworth & Lapworth (1921). *J. chem. Soc.* 119, 768.
James (1940). *Biochem. J.* 34, 640.
McLeod (1938). *Biochem. J.* 32, 1770.
Thorpe & Williams (1941). *Biochem. J.* 35, 52.
Tréfouel, Tréfouel, Nitti & Bovet (1937). *Ann. Inst. Pasteur*, 58, 39.