XLV. THE OXYTOMIC HORMONE OF THE POSTERIOR LOBE OF THE PITUITARY GLAND.

V. RECOGNITION AS AN OXIDATION-REDUCTION SYSTEM.

BY JOHN MASSON GULLAND
AND SYDNEY STEWART RANDALL¹.

From the Biochemical Department, The Lister Institute, London.

(Received December 20th, 1934.)

IN continuation of the study of the effects of chemical reagents on the physiological activity of the oxytomic hormone, it was important to examine its behaviour with reducing agents. Amongst such reagents electrolytic hydrogen and hydrogen sulphide suggested themselves as being peculiarly suitable for trial on account of the mildness of the conditions under which they can operate. This communication describes the partial inactivation of the hormone by electrolytic hydrogen and by hydrogen sulphide; the changes which the hormone then undergoes when freed from the reagents or when treated with oxidising agents such as molecular oxygen or the oxidised forms of certain dyes; and finally the examination of the equilibria developed in mixtures of the hormone and indicators of known oxidation-reduction potentials.

DESCRIPTION OF RESULTS.

The physiological activity of the hormone was diminished to about 55 % of its initial value by electrolytic reduction for 4 hours. No immediate change in activity occurred when the reduced solution was preserved from contact with

Fig. 1.

Fig. 2.

air for some hours, but when oxygen was passed into the solution, the activity first rose to about 85 % and then fell to some 60 % of the initial value (Fig. 1).

¹ Lister Institute Research Student in Biochemistry.

Hydrogen sulphide, when added to a solution of the hormone at p_H 3.5, caused the physiological activity to fall at a moderate rate to 42 % of its initial value. At this point the activity then remained constant (Fig. 2 and Table I). Aeration of such solutions with currents of moist air to remove excess of hydrogen sulphide was followed in acid and in alkaline solutions by a slow temporary recovery of oxytomic activity, which probably rose to its maximum (about 75 % of the initial value) in the course of 100 hours. The activity then slowly decreased again, and the experiments were stopped. It might have been expected that the progressive gain in activity would have complicated the assays, but in practice no difficulty was experienced, since the rate of reactivation was so slow.

A different picture was presented by the results of the next series of experiments. Hitherto, no precautions had been taken to exclude air from the reduction, and all the solutions had been prepared from distilled water containing dissolved air. The reductions with hydrogen sulphide were therefore repeated using solutions which had been boiled to expel air. This modification did not greatly affect the value to which the oxytomic activity fell as the result of the reduction (about 45 %), but when samples were diluted with distilled water containing air, the physiological values rose so rapidly that it was quite impossible to assay the solutions by the customary method of arranging the contractions of the guinea-pig uterus in groups of four [Gulland and Newton, 1932]. A method of assay, elaborated in the experimental section, demonstrated clearly (Fig. 3) that the activity rose during 1 hour from about 45 % to about 80–90 %, that it then fell to about 55 % during the course of a further hour, and that it then remained apparently constant.

Fig. 3.

Fig. 4.

Both in this and the preceding series of experiments, the dilution of the reduced hormone solutions with ordinary distilled water resulted in a temporary gain in activity followed by a decrease. In the first series (reduction in presence of air) the gain and fall were slow, occupying together about 200 hours. In the second series, the gain and fall were rapid, occupying together about 2 hours. These divergences must be attributed to the presence or absence of air during the reduction, but no information is at present available as to the identity, or otherwise, of the changes responsible for the increases and subsequent decreases of activity in these experiments.

In view of the complicated results described above which might have been due to the effects of oxygen on the products of the reduction by hydrogen sulphide, it was necessary to study the reduction and subsequent behaviour of the hormone under conditions which remained anaerobic until the doses were

introduced into the testing-bath for assay. Such conditions were realised by the use of boiled solutions cooled in nitrogen and by carrying out the reductions and dilutions in atmospheres of pure nitrogen. Under these circumstances and when the assay was performed immediately, the oxytotic value of a partially purified hormone solution decreased to about 55–60 % of its value after the reduction had proceeded for 40 hours and thereafter began slowly to increase again (Fig. 4). In exactly similar experiments with a less highly purified hormone solution containing more proteinoid material, the reduction proceeded less rapidly; the experiment was not prolonged sufficiently to ascertain the final value or to determine whether the activity again increased (Fig. 4, curve *A*). Evidently the course of the reduction was influenced by the presence of other substances.

Experiments were then made to ascertain whether the oxytotic activity was affected by the mere removal under anaerobic conditions of hydrogen sulphide from the reduced hormone solutions (55–60 %, above) and its replacement by pure nitrogen. Assays were made of diluted samples of reduced solutions immediately after the dilution. The solutions were then freed from hydrogen sulphide by repeated evacuations and fillings with nitrogen, and second assays made at intervals varying between 1 and 2½ hours from the first assays gave values for the activities which were different from those of the first assays. At first sight, it appeared as if hydrogen sulphide were affecting the response of the uterus to doses of the hormone, but this explanation was ruled out by demonstration that comparable and even much greater doses of hydrogen sulphide, introduced directly into the bath, did not modify the response to standard doses of hormone solution.

A study was therefore made of the changes in oxytotic value which occurred with passage of time when reduced hormone solutions were freed from hydrogen sulphide and kept under anaerobic conditions (Figs. 5 *a* and 5 *b*). The activity

Fig. 5 *a*. Hormone solution 2.

Fig. 5 *b*. Hormone solution 1.

first increased, and the extent and rate of this reactivation in the case of a hormone solution which contained relatively much physiologically inactive material (Fig. 5 *a*) were greater than in the case of a purer solution (Fig. 5 *b*), indicating that reactivation was possibly due, at any rate in part, to interaction of the hormone with other substances present in the solution. The value to which the activity rose in the reactivations was less than 100 %, and this change was followed by a loss of activity, of which the final result was a value of about 50 %. In the case of the less pure hormone solution (Fig. 5 *a*), however, the

exact progress of the changes of activity after the peak of reactivation is not clear. The curves of Fig. 5a have been drawn as closely as possible to the experimental points, and it may be seen that when the duration of reduction was short there is evidence of a second peak, which became smaller and finally disappeared as the duration of the reduction with hydrogen sulphide was increased. It has not been possible to carry out sufficient experiments to establish this fact, and it is not necessary to investigate it further at present. If this peculiar behaviour is correct and is a property of the hormone, it must be investigated later with a purer specimen; if it is due to the action of other substances present in the solution, its full investigation is not a matter of immediate importance for this communication.

The experiments described above indicate that oxidation or reduction of at least one group in the hormone molecule was accompanied by, and presumably caused, variations in the oxytomic activity. A preliminary investigation of the oxidation-reduction potential of this system was made by studying the action of oxidised dyes on the hormone reduced by hydrogen sulphide. Methylene blue, for example, completely restored the activity of the reduced hormone, presumably by oxidation of the reduced group. Neutral red, on the other hand, effected only partial reactivation at p_H 3.6 and seemed to have little action at p_H 6.6 and 8.5. The examination of the redox system by this method was unreliable, however, partly because hydrogen sulphide itself is unsatisfactory as an indicator of redox potential, and partly because it might attack the molecule otherwise than by straightforward reduction. Moreover, neutral red is unsuitable as an indicator since its reduced form is unstable [Clark and Perkins, 1932]. Reliable results, however, were to be expected from assays of the oxytomic activity at equilibrium of mixtures of hormone solutions and the oxidised and leuco-forms of reversible oxidation-reduction indicators. Fig. 6

Fig. 6.

Fig. 7.

shows how the oxytomic activity of the hormone varies with changes in the electrode potential of the medium between +0.04 and -0.26 v. at p_H 6. The left-hand portion of the curve represents the theoretical relationship between the percentage of the oxidised form of a redox system and the potential in volts, drawn in this case to a scale such that the values representing the oxidised and reduced forms correspond to those denoting the highest and lowest oxytomic activities determined. The right-hand section of the curve is the mirror image of the left-hand portion. Equilibrium was rapidly attained, except at potentials of -0.12 and -0.16 v., when an upward drift towards increased activity oc-

curred so rapidly that assay by the normal procedure was impossible. These drifts (Fig. 7) were followed by the method of assay developed in the reactions after reduction by hydrogen sulphide, and the values denoted by open circles in Fig. 6 were estimated by extrapolating these curves to zero time.

At potentials of $+0.35$ (phenolindophenol) and -0.45 v. (methyl viologen) (not shown in Fig. 6) the oxytocic activity was 100 %.

DISCUSSION.

The experiments described above demonstrate that the oxytocic hormone contains one oxidation-reduction system, having $E_0' - 0.025$ v. at p_H 6. When the system is oxidised the oxytocic activity is 100 %, when reduced, 50 %. It is interesting to recall that Johnson *et al.* [1927] concluded that oxidation-reduction processes constitute an important factor in the regulation of the tone of uterine muscle.

The results might also be interpreted as showing that the hormone molecule contains a second oxidation-reduction system having $E_0' - 0.190$ v. at p_H 6. On this view, the oxytocic activity would be 100 % when both systems are oxidised or reduced, and 50 % when one is oxidised and the other reduced. The existence of the second system is possible, but there are certain objections to its unqualified acceptance, and these are discussed below.

The oxytocic activity remains at 100 % after treatment of the hormone with methyl viologen at a potential of -0.45 v. or with alkaline glucose. On the other hand, electrolytic reduction for 4 hours lowers the activity to 56 %. These facts are not in harmony if the reagents only attack the oxidation-reduction systems. If the hormone contains two redox systems, as seems to be indicated by the experiments with redox potential indicators and alkaline glucose, electrolytic reduction should not have decreased the activity. Possible explanations of the apparent anomaly are that the duration of the electrolytic reduction was insufficient to allow both systems to be reduced or that its effect was complicated by exposure of the reduced hormone to air at the surface of the liquid in the reduction vessel. In the hope of settling this point, the hormone was reduced catalytically with hydrogen and platinum black for many hours; the activity fell rapidly, and remained at 50 %. This fact seemed at first to support the results of the electrolytic reduction and to show that the hormone contained only one redox system, the activity being 50 % when this was reduced. In that case, the increasing oxytocic activities corresponding to increasingly negative potentials, as shown by the right-hand portion of Fig. 6, might be the result of oxidation of the reduced system of the hormone through a rapid transfer of oxygen by the more negative redox potential indicators when the dose is introduced into the oxygenated Ringer solution in the testing bath.

It was demonstrated in the following ways that transfer of oxygen is not alone responsible for the higher activities at the more negative potentials. The hormone was reduced to an activity value of 50 % by indigodisulphonate at a potential of -0.09 v. (Fig. 6) and was then mixed anaerobically with an air-free solution of oxidised methyl viologen. No marked change in the potential of the solution occurred, as would have been shown by a large increase in the blue colour of the solution, but the oxytocic activity rose to 74 %. In an exactly similar experiment in which rosinduline replaced methyl viologen the activity rose to 59 %. These results indicate that transfer of oxygen to the hormone by the dye may occur to some extent, and the mutual relationship of the values in the two experiments supports this view. It was expected, however, that

complete reactivation of the hormone would have occurred if transfer of oxygen were the sole cause of the increasing oxytomic activity with the more negative indicators.

Further attempts were made to investigate the question of oxygen transfer by making assays under conditions in which the supply of available oxygen was considerably reduced. Johnson *et al.* [1927] record that when the isolated guinea-pig uterus is suspended in Ringer solution through which nitrogen, instead of oxygen, is passing, it contracts and remains in that state of tonus until oxygen is admitted. Attempts to repeat these observations were unsuccessful, even when nitrogen, freed from oxygen as described in this communication, was used and when the Ringer solution was covered with a layer of medicinal paraffin to exclude air. Spontaneous contractions of the uterine horn did occur, only to be followed by relaxation in the usual manner. We do not wish to stress this difference between our observations and those of Johnson *et al.*, or to offer any adverse criticism of their results. It has, however, been possible to make completely satisfactory assays of hormone solutions using a uterine horn suspended in Ringer solution covered with paraffin and saturated by a rapid stream of pure nitrogen. The only oxygen available must either (a) have been absorbed by the horn during its momentary exposure when the bath was emptied between doses or subsequently from the fresh Ringer solution, or (b) have remained dissolved in the Ringer solution in the bath at 37° for about 8 or 10 minutes in a rapid stream of nitrogen. This method was utilised to assay some hormone solutions which had been reduced with hydrogen sulphide or used in electrode potential experiments. No difference within the limits of experimental error could be detected between the activities of solutions assayed in this and in the normal way, except in the case of the indigodisulphonate-methyl viologen experiment described above; in this case, the activity was 61 %, instead of 74 % determined by the normal method.

It seems therefore that transfer of oxygen by the more negative indicators may cause oxidation of the hormone and consequent increase in oxytomic activity, but that this transfer is insufficient to account for the apparent activity of 100 % exhibited by the hormone at potentials of -0.247 v. and lower.

This conclusion strengthens the possibility that the hormone molecule contains two oxidation-reduction systems, and support for this view might be derived from the course of anaerobic reduction by hydrogen sulphide—initial decrease of activity to 50–60 %, followed by a slow increase. The chief evidence to the contrary was that catalytic or electrolytic reduction lowered the activity to about 50 %, instead of leaving it unchanged at 100 %. It was shown as follows that this was probably due to adsorption of the hormone or to its destructive reduction. A hormone solution which had been reduced by platinum black and hydrogen and exhibited 50 % of its initial activity was mixed anaerobically with oxidised methylene blue in order to convert the redox system completely into its oxidised form as in the experiments after reduction with hydrogen sulphide. The activity first fell, a value of about 33 % being recorded, and then rose again to 50 %, where it remained constant. These changes would have been predicted on the assumption that two redox systems in half the hormone molecules present in the solution had been fully reduced, resulting in an activity of 100 % as regards those molecules, and that the other half of the hormone molecules had been adsorbed on the catalyst or destroyed. In agreement with this view, it was found that oxidised rosinduline, added anaerobically to the reduced hormone solution (activity 50 %), did not alter the activity. Further, an attempt to restore full activity to an electrolytically reduced solution

(value 70 %) by adding oxidised methylene blue only raised the activity to 83 %, indicating that some hormone molecules had been destroyed. These facts are in agreement with the existence of two redox systems. On the other hand, the addition of reduced rosinduline to the catalytically reduced hormone solution lowered the activity from 50 % to about 20 %, a fact which is not in harmony with these views.

Our present opinion is that the hormone contains one oxidation-reduction system, that it may also contain a second group which readily undergoes reduction, and that the states of oxidation of these groups are intimately connected with the physiological activity of the hormone. In addition, transfer of oxygen to the reduced hormone may occur in the testing bath through the agency of the more negative redox potential indicators.

The decision as to whether the drift of the oxytocic activity of the hormone at electrode potentials between about -0.11 and -0.18 v. or after reduction by hydrogen sulphide is an inherent property of the molecule or results from the presence of other substances must be postponed until purer hormone solutions are available. The subsequent inactivations which follow these increases in activity are also intimately linked with this question. Whatever the explanation may be, it is clear that molecular oxygen is not a necessary factor, since both these changes can occur in its absence. On the other hand, oxygen may well take part in the aerobic reactivations, and undoubtedly it is responsible for the existence of the hormone in its oxidised state in posterior pituitary extracts. It would be unwise to attempt an explanation of these activations and inactivations without a knowledge of the simultaneous electrode potentials of the solutions in which they were occurring. Possibly measurements of these may throw light on these questions.

EXPERIMENTAL DETAILS.

Hormone solutions.

Three solutions of the hormone were used in this investigation. One, called hormone solution 1, was prepared according to Kamm *et al.* [1928], and corresponded in its state of purification to that used by Gulland and Newton [1932]. It contained 300 oxytocic units per ml., associated with 5.65 mg. of solid when dried to constant weight.

The second solution, called hormone solution 2, was prepared by extracting dried, powdered posterior lobe (1.5 units per mg.) with 0.25 % acetic acid, concentrating the extract and precipitating by saturation with ammonium sulphate as described by Kamm *et al.* [1928]. The precipitate was then thoroughly extracted with 98 % acetic acid, and the extract was mixed with ether (2.5 vols.) and light petroleum (5 vols.). The solid thus precipitated was washed with a mixture of ether and light petroleum and was dissolved in a convenient volume of water at p_H 3. This solution therefore contained more inactive material than hormone solution 1, the actual content being 28.3 mg. of solid when dried to constant weight associated with 450 units per ml. The amount of ammonium sulphate was 3.9 mg. per ml., based on estimations of sulphate ion as barium sulphate, so that the weight of glandular material was 24.4 mg. per ml.

The third solution, hormone solution 3, prepared as described by Kamm *et al.*, contained 900 units per ml., associated with 3.67 mg. of solid.

Electrolytic reduction and reactivation by oxygen (Fig. 1).

The apparatus used was a rectangular three-compartment cell of the type used for electrodialysis. The electrodes were platinum foil, each measuring

2.5 × 2.5 cm. and the compartments were cooled by glass coils containing flowing water. The dividing membranes consisted of cellophane No. 400, and each compartment contained 10 ml. of liquid. Experiments on the electrodialysis of the hormone will be described at an early date in another communication.

The following description is typical of several experiments. The cathode chamber contained a mixture of hormone solution 2 (0.2 ml. containing 90 units) and *M*/10 sodium acetate solution (0.1 ml.) diluted to 10 ml. The centre compartment was filled with water (9 ml.) and *M*/10 acetate buffer at p_H 3.5 (1.0 ml.), and the anode chamber contained *N*/1000 acetic acid (10 ml.). The current was maintained at 50 milliamps, the voltage across the cell being 60 v. in the early part of the experiment and somewhat higher after the first half-hour. The temperature remained at 15°. The cathode solution was kept adjusted to p_H 9 by passing in carbon dioxide as required.

After the reduction had proceeded for 1 hour, the cathode solution was transferred quantitatively to a 25 ml. flask, brought to p_H 3.5 by the addition of 0.5 ml. of *N*/10 hydrochloric acid and diluted to 25 ml. When diluted and assayed, it had 66 % of the total activity, and this value remained unchanged when the undiluted solution, which still contained dissolved hydrogen, was kept in a cooled vessel for 18 hours.

When the reduction was prolonged for 4 hours and the cathode solution was treated as before, it contained only 56 % of the total activity. In both cases the liquid in the anode compartment exhibited no activity, and the centre compartment contained less than 1 % of the total number of units.

A stream of moist oxygen was passed for 1 hour through the cathode solution which had been reduced for 4 hours. The assay of a sample, taken at the end of this period and tested at once, showed that the activity had risen to 76 %, but after this oxygenated solution had been kept at 0° for 18 hours, the activity had fallen to 60 %. In a parallel experiment, the activity after oxygenation for 1 hour was 75 % and 3 hours later 83 %.

Aerobic reduction by hydrogen sulphide and reactivation by aeration (Fig. 2).

Mixtures of hormone solution 1 (0.1 ml. containing 30 units), *M*/10 acetate buffer at p_H 3.5 (0.5 ml.) and hydrogen sulphide solution (1.0 ml. saturated at 20°) were diluted to 2.5 ml. and kept at room temperature. The results are summarised in Table I.

Table I.

EXPERIMENT 1.			EXPERIMENT 2.		
Time hours	Treatment	Activity %	Time hours	Treatment	Activity %
2.5	Sample taken	76	2	Sample taken	75
26.5	"	46	7	"	50
50.5	"	42	25	"	42
75	"	43	25.5	Aeration by moist air for 1 hour at p_H 3.5	—
80	Aeration by moist air for 45 mins.	—	26.5	Sample for $Ce(OH)_2$ exp.	—
98	Sample taken	45	50.5	Sample taken	50
102	p_H adjusted to 8.2	—	75	"	53
119	Sample taken	45			
170	"	68			
218	"	67			

It is evident that the reactivation occurs very slowly in acid or alkaline solution. In order to ascertain whether the rate of reactivation would increase and the activity be restored completely if the formation of hydrogen peroxide were avoided or its amount, if formed, were diminished, a portion (1.0 ml.) of

the solution from Exp. 2, which had been reduced for 25 hours and then aerated was adjusted to p_H 8.2 with $N/10$ sodium hydroxide and then mixed with borate buffer at p_H 8.2 and a suspension of cerous hydroxide prepared from $M/50$ cerous sulphate (1.0 ml.). The suspension was shaken, and samples (not shown in Table I) were taken at 49 and 73 hours, centrifuged, diluted and assayed; the values for activity were 60 and 46 % respectively. The results of this experiment seemed to indicate that reactivation was more rapid in presence of cerous hydroxide. This point must not be stressed, however, since it is possible that the rate of oxidation was accelerated at the surface of the finely divided cerous hydroxide. In addition, no information could be obtained as to the adsorptive effect of cerous hydroxide on the reduced hormone or on the oxidised hormone formed in its presence.

Anaerobic reduction with hydrogen sulphide and reactivation.

The oxygen-free nitrogen used in all the anaerobic experiments was prepared by storing nitrogen (from a cylinder) over alkaline pyrogallol, scrubbing it thoroughly with alkaline pyrogallol immediately before use, passing it through a roll of reduced copper gauze in an electric furnace at a low red heat, and then washing it with air-free water. When purified in this way, the nitrogen gave no coloration with alkaline pyrogallol.

All solutions used in the anaerobic experiments were rendered free from oxygen as follows: they were either boiled and cooled quickly while a rapid stream of nitrogen passed through them, or if that procedure were unsuitable, they were repeatedly evacuated and saturated with nitrogen.

Reduction. The hormone solution (0.1 ml.) was mixed with $M/10$ acetate buffer at p_H 3.5 (0.5 ml.) and water (0.9 ml.). Part (0.3 ml.) of this solution was then freed from air and mixed with a saturated solution (0.2 ml.) of hydrogen sulphide in air-free water in a small tube swept out by nitrogen. The tube was then tightly stoppered and placed in a thermostat at 20°. A number of tubes prepared in this way at the same time formed the material for one experiment. After the reduction had proceeded for the desired time, one tube was introduced into a separating funnel containing air-free water (24.5 ml.) in an atmosphere of nitrogen. By evacuating the funnel, the stopper of the small reduction tube was released, care being taken not to allow hydrogen sulphide to escape from the funnel; the contents of the tube and funnel were thoroughly mixed, and the funnel was again filled completely with nitrogen. These experiments were repeated many times using hormone solutions 1 and 2, and some of the results, which differed slightly, are recorded in Fig. 4. These assays were not complicated by changes due to reactivation, presumably because these, if they occurred at all, were slow in presence of hydrogen sulphide.

Aerobic reactivation. In the early experiments on the reduction described above, the contents of the small tubes were diluted with air-containing water (24.5 ml.) and not as described above. Under these conditions it was observed that the activity seemed to have increased during the first hour after the dilution and then to have fallen at the end of 2 hours to approximately the original value of the reduced solution. The rate of change was too rapid to allow assays to be made by the customary procedure of arranging contractions of the uterus in groups of four [Gulland and Newton, 1932], but it was possible to obtain a moderately accurate estimate of the activity at any given time after dilution (a) by making assays simultaneously on two uterine horns previously brought to a satisfactorily sensitive condition by doses of standard hormone solution; and (b) by arranging doses of the solutions either in the order experimental,

experimental, standard, standard, or in the order experimental, standard, standard, experimental. From the charts thus obtained, the activity values of the experimental solutions were estimated by direct comparison of the heights of contractions evoked by them and by the adjacent doses of standard solution. It is evident that the results obtained by this procedure cannot be so reliable as those reached by the customary method of assay.

A number of experiments were made in this way, and in some cases the assays were made by two observers working simultaneously, but independently, with different uterine horns. These changes are illustrated by Fig. 3, from which it is clear that after reduction with hydrogen sulphide and dilution with aerated water the activity rose during 1 hour and then fell to a steady value close to that of the original reduced solution.

Anaerobic reactivation. A series of reductions of hormone solutions 1 and 2 was carried out as described above, and the reduction tubes were exploded as before in air-free water in an atmosphere of nitrogen. As soon as the contents of the separating-funnel had been mixed, a dose was withdrawn without exposing it to the air and inserted immediately into the testing-bath. After some practice it was possible to arrange that this dose should cause a contraction of which the height lay between two adjacent doses of standard solution. The activity of the reduced sample was thus ascertained with some measure of certainty, since in these experiments only those uterine horns were used which at the beginning of the assay were not only giving uniform responses to identical standard doses but which also exhibited a satisfactory degree of sensitivity.

The anaerobic removal of doses of the experimental solutions from the tube below the tap of the separating funnel was easily accomplished by means of a 1 ml. hypodermic syringe graduated in 1/100 ml.

The remainder of the solution was at once freed from hydrogen sulphide by four successive evacuations and replacements by nitrogen, and the changes in activity were followed simultaneously in two uterine horns as described in the case of the aerobic reactivation experiments. Some results are given in Figs. 5 *a* and 5 *b*.

Action of dyes on the hormone after anaerobic reduction (Table II).

Hormone solution 2 (0.1 ml.), *M*/10 acetate buffer at p_H 3.5 (0.5 ml.), and hydrogen sulphide solution (1.0 ml., saturated at 20°) were mixed in a flask swept out by nitrogen. The mixture was diluted to 2.5 ml. with air-free water, and kept in a thermostat at 20°; the period of reduction varied between 24 and 48 hours in different experiments.

Methylene blue at p_H 4.2 and neutral red at p_H 3.6. The reduced hormone solution (1.0 ml.) was transferred without being exposed to air to a specially constructed tube swept out by nitrogen and containing air-free water (8 ml.). The tube was then evacuated and refilled with nitrogen four successive times, and whilst it was still under slightly reduced pressure an air-free, *M*/50,000 aqueous solution (1.0 ml.) of the oxidised form of the dye was added from a burette fixed in the stopper. Some minutes later (accurately timed), the p_H was determined, and a sample was removed anaerobically by a tap in the bottom of the tube and diluted for assay. Difficulties due to changes in activity were not experienced in making assays in the presence of these dyes, which seemed to stabilise the oxytocic value.

A control experiment was carried out simultaneously on another portion of the same reduced solution; the dye solution was replaced by water, and care was taken to ensure that the conditions as to time during which the dye acted

and rates of evacuation and refilling were copied as faultlessly as possible. These precautions were necessary in view of the reactivation which occurs when the hydrogen sulphide is removed.

Neutral red at p_H 6.6 and 8.6. These experiments were carried out similarly to those described above, except that the dye was added as a mixture of $N/10$ sodium hydroxide (0.1 ml.) and 2.0 ml. of $M/50,000$ neutral red in $M/5$ sodium phosphate buffer at p_H 6.6 or in Palitzsch's borate buffer [Clark, 1928] at p_H 8.6 respectively. Control experiments were made in the same way.

Table II.

Dye	p_H	Time of standing with dye mins.	% activity of control	% activity with dye
Methylene blue	4.2	10	69	104*
"	4.2	5	50	100
Neutral red	3.6	4	55	71†
"	3.6	5	64	78‡
"	6.6	5	67	65
"	6.6	5	63	69
"	8.5	5	42	57
"	8.5	4	70	69

* 100 % after remaining 1.5 hours in nitrogen.

† 88 % " 24 "

‡ 80 % " 21 "

Table III. *The relationship between oxidation-reduction potential and oxytocic activity.*

Dye	p_H	% oxidised form	E'_0	Correction K	E_h	% activity
Phenolindophenol	6.0	99	+0.29	+0.060	+0.35	96
Methylene blue ¹	6.0	25	+0.047	-0.014	+0.033	100
	6.0	14	+0.047	-0.024	+0.023	97
Indigotetrasulphonate ²	6.0	20	+0.006	-0.018	-0.012	87
Indigotrisulphonate ²	6.0	20	-0.028	-0.018	-0.046	59
	6.0	12.5	-0.028	-0.026	-0.054	53
Indigodisulphonate ²	3.5	15	+0.080	-0.023	+0.057	55
	6.0	13.5	-0.069	-0.025	-0.094	51
	6.0	15	-0.069	-0.023	-0.092	50
Brilliant alizarin blue ³	6.0	30	-0.112	-0.011	-0.123	53*
Anthraquinone- β -sulphonate ⁴ †	6.0	70‡	-0.170	+0.011	-0.159	59*
	3.5	20	-0.040	-0.018	-0.058	73
	5.6	20	-0.160	-0.018	-0.178	74
Rosinduline 2G ⁵	6.0	70‡	-0.221	+0.011	-0.210	87
	6.0	14	-0.221	-0.025	-0.246	97
	6.0	12.5	-0.221	-0.026	-0.247	100
	6.0	20	-0.221	-0.018	-0.239	106§
Methyl viologen ⁶	10.0	40	-0.446	-0.005	-0.45	100

approx.

¹ Clark *et al.* [1925].

² Sullivan *et al.* [1923].

³ Michaelis [1930].

⁴ Conant *et al.* [1922].

⁵ Michaelis [1931].

⁶ Michaelis and Hill [1933].

* During the assays of these experiments the activity drifted upwards, and these values were estimated by extrapolating to zero time the curves representing the drifts.

† Anthraquinone- β -sulphonate is practically colourless, and it was assumed that the oxidised form was present to an extent of 20 % of the total dye in the reaction mixture, since this was the average figure for a series of experiments made with coloured indicators at the same period.

‡ 0.5 ml. of the oxidised dye solution was mixed with the hormone solution, and 0.5 ml. of the leuco-dye solution was added from the burette.

§ This assay was made under as completely anaerobic conditions as possible. Nitrogen, free from oxygen, was substituted for the oxygen passing through the Ringer solution in the testing-bath, and a layer of medicinal paraffin covered the surface of this Ringer solution.

Oxidation-reduction potential of the hormone system (Table III).

These experiments were made in an apparatus in which the following series of operations could be carried out in an atmosphere of air-free nitrogen with solutions which had previously been freed from air. A $M/5000$ aqueous solution of the leuco-dye, prepared by reduction with platinised asbestos and hydrogen, was filtered through a sintered glass plate into a burette. 1.0 ml. was added to a mixture of the hormone solution (1.0 ml. of a $1/100$ dilution of solution 2 or 0.5 ml. of a $1/100$ dilution of solution 3, 4.5 units in each case), $M/10$ buffer solution (1.0 ml.) and water (7.0 or 7.5 ml.), the final volume being 10.0 ml. In every case oxidation of the leuco-dye occurred, and the percentage of oxidised form present in the reaction mixture was estimated colorimetrically. The total molarity of the dye in the mixture was $1/50,000$. Temperature was 18° approximately. The samples for assay were removed at a tap in the reaction vessel without being exposed to air and were introduced immediately into the testing bath, the dilutions having been suitably arranged to enable this to be done. Equilibrium appeared to be attained immediately in most cases, but at potentials of -0.12 and -0.16 v. the activity value drifted rapidly (see footnote to Table III), and a slow drift was also observed at -0.19 v.

The effects of adding oxidised potential indicators to the hormone reduced with indigodisulphonate.

A $M/5000$ aqueous solution (1.0 ml.) of leuco-indigodisulphonate was added to an air-free mixture of a dilution of hormone solution 3 (0.5 ml., 4.5 units) $M/10$ acetate buffer at p_H 6 (1.0 ml.) and water (6.5 ml.) in the apparatus used in the previous experiments. An air-free $M/5000$ aqueous solution (1.0 ml.) of methyl viologen was then added, and assays of oxytomic activity were made in the normal way and in a stream of nitrogen (see Discussion). The values were 74 and 61 % respectively.

Addition of a $M/5000$ aqueous solution (1.0 ml.) of oxidised rosinduline 2 G instead of methyl viologen gave a solution which assayed at 59 %.

Reduction with hydrogen and platinum black.

The catalyst was a strip of platinum foil (8×0.5 cm.) coated electrolytically with a thin film of platinum black and washed by electrolysis in dilute sulphuric acid and then distilled water. A mixture (20 ml.) of a dilution of hormone solution 3 (2.22 ml., 20 units), $M/10$ acetate buffer at p_H 6 (2.0 ml.) and water was shaken mechanically with the catalyst in an atmosphere of hydrogen. After $1\frac{1}{2}$ hours an assay showed the activity to be 59 %, and after a further period of reduction of 11 hours the activity was 50 %.

Equal volumes of this reduced hormone solution and $M/5000$ solutions of potential indicators were mixed anaerobically and the results are summarised in Table IV.

Table IV.

Potential indicator	% of initial activity	Time elapsed from mixing indicator with reduced hormone mins.
Oxidised rosinduline	50	3
	50	30
Oxidised methylene blue	33	6
	50	60
Reduced rosinduline	20	6
	20	60

Electrolytic reduction and reactivation by methylene blue.

Hormone solution 2 was reduced electrolytically, as described previously, until an assay showed that the activity was 70 %. The reduced solution (5 ml.) was freed from dissolved hydrogen by repeated evacuation and saturation with nitrogen and was mixed anaerobically with a $M/5000$ aqueous solution (2.0 ml.) of methylene blue. An assay showed the activity to be 83 % of its initial value.

SUMMARY.

1. The oxytocic activity of the hormone falls to about half its initial value when the molecule is reduced by hydrogen sulphide. The reduced hormone is completely reactivated by oxidised methylene blue.

2. The hormone is also reduced by catalytic or electrolytic hydrogen, the activity again falling to about 50 %. Part of this decrease in activity is probably due to destruction of the hormone or to its adsorption by the catalyst or electrode.

3. The reduced hormone is partially reactivated when it is exposed to air or when the hydrogen sulphide is removed under anaerobic conditions. These reactivations are followed by inactivations. These processes vary according to the circumstances in which they occur and the conditions under which the hormone was reduced.

4. Assays of the oxytocic activity of the hormone in presence of indicators of known oxidation-reduction potential indicate that the hormone molecule contains a redox system having $E_0' + 0.025$ v. at p_H 6. The activity is 100 % when this is oxidised and 50 % when it is reduced.

5. Assays of the oxytocic activity of the hormone at potentials between -0.10 and -0.25 v. show that the activity increases from 50 to 100 % over this range. This may indicate the presence in the molecule of a second group which readily undergoes reduction, but the interpretation of the results is complicated by the possibility that some oxidation of the reduced hormone in the testing bath may occur through a rapid transfer of oxygen to it by the more negative redox potential indicators.

We are much indebted to Dr B. C. J. G. Knight and Dr G. M. Richardson for valued advice and criticism and to Dr K. G. Stern and Mr G. D. Greville for specimens of rosinduline and methyl viologen respectively.

REFERENCES.

- Clark (1928). The determination of hydrogen ions, 3rd ed. (Baillière, Tyndall and Cox.)
— Cohen and Gibbs (1925). *U.S. Pub. Health Rep.* **40**, 1131.
— and Perkins (1932). *J. Amer. Chem. Soc.* **54**, 1228.
Conant, Kahn, Fieser and Kurtz (1922). *J. Amer. Chem. Soc.* **44**, 1382.
Gulland and Newton (1932). *Biochem. J.* **26**, 337.
Johnson, McClosky and Voegtlin (1927). *Amer. J. Physiol.* **83**, 15.
Kamm, Aldrich, Grote, Rowe and Bugbee (1928). *J. Amer. Chem. Soc.* **50**, 573.
Michaelis (1930). *J. Biol. Chem.* **87**, 713.
— (1931). *J. Biol. Chem.* **91**, 369.
— and Hill (1933). *J. Gen. Physiol.* **16**, 859.
Sullivan, Clark and Cohen (1923). *U.S. Pub. Health Rep.* **38**, 1669.

ERRATUM

Vol. XXIX, page 390, in the Summary No. 4

for $E_0' + 0.025$ read $E_0' - 0.025$