XCI. A PRELIMINARY NOTE ON THE APPEAR- ANCE OF PARESIS IN ADULT RATS SUFFERING FROM CHRONIC AVITAMINOSIS E.

By AXEL RINGSTED.

*From the University Institute of Hygiene and the State Vitamin Laboratory,
Copenhagen, Denmark.*

(Received January 24th, 1935.)

THIS paper deals with the observation of some characteristic neuropathic disturbances which have developed in adult albino rats fed on various vitamin E-free food mixtures. Similar observations have not, to my knowledge, been previously dealt with, though Evans and Burr [1928] have observed the appearance of paresis in suckling rats bred from vitamin E-free mothers.

Evans and Burr [1928] found that the progeny of mothers fed on a diet very poor in vitamin E developed paresis in the hind quarters during the latter part of the lactation period. The paresis appeared as a rule on about the 20th day after birth and always before the 25th day. Otherwise the animals did not become paretic. An initial relaxed paretic stage appeared in some of the animals, whilst the majority showed signs of spasticity in the hind legs from the outset. A claw-like condition of the feet developed later. No disturbance of sensibility was evident. In the course of the following months, symmetrical baldness of the hind quarters and femur and atrophy of the skin occurred. Paresis spontaneously disappeared in scarcely 20 % of the diseased animals. Some individual animals suffering from a mild attack of paresis were cured by a liberal supply of substances containing vitamin E. The great majority appeared to be quite unaffected by treatment with various vitamins—even vitamin E.

Evans and Burr were able to show, by means of a wide range of experiments, that this paresis was due to lack of vitamin E in the diet, and that other organic and mineral ingredients of the food had, *per se*, no direct causal significance in the appearance of paresis. The experiments showed that it was possible completely to prevent the appearance of paresis by giving the mothers, prophylactically, substances rich in vitamin E, such as yellow corn, corn oil and wheat germ oil. It was evident that it was the vitamin E component of wheat germ that was the conclusive factor in experiments with many different fractions of wheat germ oil. These experiments showed a quantitatively exact agreement in the properties of the wheat germ in curing sterility and preventing paresis.

Evans and Burr's observation on this paresis was later confirmed by Mason [1933]. Mason succeeded in producing this paralysis in suckling rats but not in keeping the latter alive. He is of the opinion that that was due to the difficulty of finding just the right quantity of vitamin E which is sufficient to maintain life and insufficient for the vigorous development of the nervous system in the lactation period.

Evans and Burr and Mason give no information as to the morbid anatomical changes in the nerve system of these paretic rats.

Experimental methods.

The animals used for experiment comprised three groups of albino rats, A, B and C, 27 animals in all, 26 female and 1 male. The mothers of these animals had all had a uniform diet, Gudjonsson's diet 4 for breeding animals. The animals were used for the experiment at the age of 24–35 days (see Tables II, III, IV). The cage of each rat was provided with a wire mesh base so that the faeces fell through.

The rats were constantly supplied with fresh drinking water, were given fresh food three times a week and were weighed once a week, and during pregnancy every day.

From puberty, a vaginal smear was taken six times a week from all the females. The purpose of this was to ascertain the regularity of the period of oestrus; to decide the optimum time for copulation; to find out whether copulation had taken place (bouchon vaginale, sperm in the vaginal smear) and whether conception had occurred (placental sign). The vaginal smear provides, furthermore, information as to the period of the cessation of the resorption-pregnancy and as to the reserves of vitamin A in the organism (see below).

The rats were kept at a constant temperature of 22°. No sign of acute or chronic infection was ever observed among the animals that were killed. The composition of the food mixtures, the content and method of administering the vitamin preparations and also their preservation and manner of distribution are given in the two following sections.

As to the composition of the breeding diet and further data as to fertility and the suitability of the experimental animals for vitamin A experiments, the reader is referred to Gudjonsson [1930, 1, 2].

Table I. *Food mixtures.*

Group of animals	...	A	B	C
Food mixture		Vitamin E-free I %	Vitamin E-free III %	Vitamin E-free II %
Caseinogen		18	20	20
Rice starch		50	53	53
Oxidised lard		20	15	15
Dried autolysed top yeast		7	7	7
Salt mixture (185)		5	5	5
Daily additions of vitamin A		Cod-liver oil	Carotene	Cod-liver oil
Daily additions of vitamin D		Cod-liver oil	Vit. D-standard	Cod-liver oil

The vitamin E-free food mixtures may all be taken together under the same heading.

Caseinogen. To avoid any possibility of its containing vitamin E, ether-extracted caseinogen was used during the first 1½ months of the experiment. As contemporaneous experiments showed that caseinogen treated with acid did not contain any appreciable quantity of vitamin E, this was used for the rest of the experiment.

Rice starch was an ordinary commercial product, as also was the lard. The lard was oxidised by being kept at a temperature of 110° for 10 hours with simultaneous aeration. This prevented the possible presence of vitamins A and E.

McCollum's salt mixture (185) was used.

At first, a peanut oil solution of a biologically standardised cod-liver oil was used as a source of vitamin A for Group A, so as to give a daily dose of 8 I.U. of vitamin A. As some of the rats about 8–12 weeks old evinced a prolonged period of oestrus¹, up to 3–4 days' duration, the

¹ Evans and Bishop [1922] showed that at the commencement of deprivation of vitamin A the period of oestrus—confirmed by the vaginal smear—became longer and dioestrus shorter. Later on the stage with cornification becomes continuous. This was subsequently confirmed by Hohlweg und Dohrn [1930] and Aberle [1934].

daily dose was increased to 40 i.u. of vitamin A of fresh standardised cod-liver oil, after which this phenomenon disappeared.

From the beginning of the experiment the animals in Group B were given 20 γ of standard carotene dissolved in sesame oil daily.

The same cod-liver oil solution in the same dose as mentioned for Group A was used for Group C from the commencement of the experiment, i.e. 40 i.u. of vitamin A.

Dried autolysed top yeast mixed with the diet so as to form 7 % of it was used as a source of vitamin B for all three groups.

The yeast was autolysed at 38° for 24 hours, after which it was dried on flat trays at 30° and then powdered. As the increase in weight began to diminish, certain of the animals were given an addition of 0.5 g. dry yeast for a longer period.

We gave no vitamin C, as rats do not need this supplied in their food.

The animals in Group A received in their daily addition of cod-liver oil 0.4, and later 2, i.u. of vitamin D, those in Group C had 2 i.u. of vitamin D throughout the whole experiment. A sesame oil solution of standard vitamin D was used for Group B, mixed with food so that the mixture contained 0.8 i.u. of vitamin D per 10 g. of food.

The vitamin E content of the food mixture is certainly very low or nil. The phenomenon of initial fertility, which is due partly to the presence of reserves of vitamin E and partly to the food containing small quantities of vitamin E, never appeared in animals fed on these food mixtures. Even animals that became pregnant after having only been on the experimental diet for 20–30 days had so exhausted their vitamin E reserves that they resorbed the foetus. The reason for the low content of vitamin E in the diet lies certainly in the thorough oxidation of the lard by which the whole of the vitamins A and E are effectively destroyed [Fridericia, 1924; Mattill and Crawford, 1931; Cummings and Mattill, 1930].

We lay the greatest stress on the very low vitamin E content of the diet which is certainly of decisive significance for the appearance of paresis (see later).

A preparation of wheat germ oil "*Fertilan*"¹ was used as a source of vitamin E. The wheat germ oil was kept in hermetically sealed ampoules at 0°. The contents of the ampoule were warmed and shaken up before use and kept at 0° after use again. An ampoule lasted for 2–3 days. The oil was measured with the same standard pipette for all the experimental groups. One drop weighed on an average 25.4 mg.

Clinical observations.

The animals in the three groups, except No. 58, thrived well for the first 8–12 weeks.

At about the 10th week of the experiment the rate of increase in weight appreciably diminished and in spite of a doubling, shortly followed by a quadrupling, of the daily dose of cod-liver oil, the animals put on weight very slowly. The average weight curve of the animals was, meanwhile, very near the average weight curve for rats of the same breeding fed on "diet 4 for breeding animals" [Gudjonsson, 1930, 2] (Fig. 1).

The females were mated with normal males between the 80th and the 100th experimental days. All pregnancies resulted in resorption. The females were, thereupon, available for use as test animals and were used for investigation as to the vitamin E content of various substances, by the curative method exclusively [Evans and Burr, 1927]. Ten females (see Tables II, III, IV) were given varying quantities of wheat germ oil at different periods after conception had been verified. They all gave birth to live progeny except No. 55 which was first given wheat germ oil on the 7th day of pregnancy² and No. 78 which had

¹ "*Fertilan*" wheat germ oil was very kindly supplied to us by the Danish Medicinal Company's Experimental Laboratory. For information as to its preparation reference may be made to Vogt-Møller's [1931] articles on this subject.

² This confirms Evans and Burr's observation that vitamin E should be given before the 5th or 6th day after conception has taken place so that the fertile ovum may develop fully.

had too small doses. None of these 10 animals later showed signs of neuropathic disturbances.

The 16 other females were used for investigation of the vitamin E content of various foodstuffs. All these and the only male later developed paresis.

Fig. 1. Average weight curves for the first 20 experimental weeks for animals in groups A, B and C compared with Gudjonsson's average weight curve (G) for female rats on "diet 4 for breeding animals".

As the details concerning these tests have no relation to the pareses described later on they are left out of this preliminary communication.

It should however be pointed out that the vitamin E content of the substances was so low that all the pregnancies resulted in resorption. In about the 19th week of the experiment it was noticed that one animal (No. 58) was beginning to drag its hind legs a little. Simultaneously the legs began to straddle slightly. About 8 weeks later, when similar disturbances in gait were observed in three other rats (Nos. 51, 53 and 56), the conclusion was come to that this was no accidental phenomenon.

It was further confirmed by a closer examination of the gait of all the animals in Groups B and C during the following weeks. It appeared that several of these animals had already begun to show disturbances in their gait. We were hereby in a position to observe the very first stage in the development of the disease.

These observations have provided the basis of the following description in which, for practical reasons, the disease has been divided into three stages, or grades, as all the animals did not develop the disturbances to the highest degree, even though they continued on the diet for several weeks after the appearance of the first symptoms.

Stage I. The first clinical symptoms are a dragging of the legs and slight degree of inco-ordination in the gait. The gait is slower and a little cautious and the feet are placed slightly out to the side¹. The strength of the hind legs is good, and the animal can jump with ease (7-8 cm.). In some cases the fur has begun to loosen on the hind quarters and the hind legs. The animal is lively and eats well. This condition changes with no definite break into stage II.

¹ The disturbance in the gait is most evident when the animal is allowed to walk over the wide-meshed wire netting of the roof of the cage.

Table II. *Group A. Vitamin E-free diet I. (Female rats.)*

Serial no.	Litter no.	Age at commencement of exp. days	Addition of W.G.O. during 1st test pregnancy	Time of appearance of paresis weeks	Stage of paresis	Treatment of paresis	Result of treatment
50	1135	26	+	—	—	—	—
51	1135	26	—	30	II	1000 i.u. of vit. A daily for 4 weeks 50 i.u. of vit. D daily for 4 weeks	Continued development of paresis to stage III
52	1136	26	+	—	—	—	—
53	1136	26	—	27	III	125 mg. W.G.O. daily for 12 weeks	No effect
54	1136	26	+	—	—	—	—
55	1136	26	+	—	—	—	—
56	1136	26	—	27	II	None*	—
58	1134	29	—	19	III	1 g. autolysed dried top yeast daily for 6 weeks	No effect
59	1134	29	+	—	—	—	—
61	BfR2	23	+	—	—	—	—

Table III. *Group B. Vitamin E-free diet III. (Female rats and one male.)*

75	1810	26	—	24	II	None	—
76	1810	26	+	—	—	—	—
77	1810	26	—	24	I	600 i.u. of vit. A daily for 3 weeks 30 i.u. of vit. D daily for 3 weeks	Continued development of paresis to stage II
78	1810	26	+	—	—	—	—
79	1810	26	—	21	II	ca. 100 mg. W.G.O. daily for 4 weeks	No effect
80	1810	26	—	20	III	None	Condition worse
82	1146	24	+	—	—	—	—
83	1146	24	—	20	I	1.5 g. autolysed yeast daily for 3 weeks	Continued development of paresis to stage II
84	1830	25	—	23	I	1.0 g. autolysed dried yeast daily for 6 weeks	Continued development of paresis to stage III
85	1830	25	—	20	III	80 i.u. of vit. A daily for 30 days 4 i.u. of vit. D daily for 30 days	No effect

Table IV. *Group C. Vitamin E-free diet II. (Female rats.)*

88	1147	35	+	—	—	—	—
89	1147	35	—	22	I	1 g. autolysed dried yeast for 6 weeks	Increased baldness and paresis to stage II
90	1147	35	—	23	I	75 mg. W.G.O. daily for 6 weeks	No effect
91	1147	35	—	34	I	None	—
103	BfR 46	24	—	16	II	None	—
104	BfR 46	24	—	17	I	50 mg. W.G.O. daily for 6 weeks	No effect
105	BfR 46	24	—	15	II	250 mg. W.G.O. daily for 6 weeks	No effect

* Nos. 57 and 60 were killed during the first test pregnancy long before the paresis had appeared. No. 56 was operated on shortly after the commencement of paresis and died a few days after the operation with symptoms of ileus.

Fig. 2. BfR. 58. Noticeable straddling of the hind legs. Low position of the body. Stage II.

Fig. 3. BfR. 58. Noticeable straddling. Soles of the feet flexed. Toes flexed. The animal treading on the lateral edge of the foot.

Fig. 4. BfR. 85. Hind quarters atrophied. Baldness most pronounced on the legs. Distinct straddling gait. Hind quarters dragging along the floor. Flexion of feet and toes. Stage III.

Stage II. This stage (Figs. 2 and 3, Plate I) is distinguished by a pronounced straddling of the hind legs when walking. A healthy rat keeps the foot under the femur so that it is covered by the latter when seen from above [Aberle, 1934]. In this stage there is a considerable diminution in the functions of the adductors. There is flexion of the sole of the foot and extension of the knee joint. The inco-ordination is pronounced. The animals frequently walk upon the side of the foot (Fig. 3) and on the tail without noticing it. Paresis of the adductor is clearly apparent when the animals are taken out of the small opening in the cage. They must be taken out in a slanting position as, otherwise, the straddling leg is caught. When moving, the hind quarters are kept low. There is incipient atrophy of the muscles of the hind quarters and hind legs. The fur begins to fall out, especially from the extremities, but to a less degree in the centre. Susceptibility to touch seems also to be lessened. If the tail and hind legs be taken in a pair of forceps, there is an impression that there is hypoaesthesia in most of the animals in this stage and hypoaesthesia in many of them.

The fore legs seem to be quite normal at this stage. The animals are lively, but move round the cage slowly.

Stage III. The gait is more characteristic in this stage (Fig. 4, Plate I). The hind legs are kept very highly abducted, and cannot be adducted. The hind quarters are dragged along the floor (Fig. 4). The animals in reality move forwards on their stomachs. The hind quarters and tail swing from side to side at every step. The sole of the foot is highly inflexed when walking, as are the small joints of the toes. If the animals are taken up by the nape of the neck, there is no dorsal flexion of the foot, as is the case with normal rats [Aberle, 1934]. Ataxia is still more pronounced and gives rise to the most grotesque attitudes and attempts to walk. These phenomena were most pronounced when the rat tries to jump up in the cage and in walking tests on the wide-meshed netting roof of the cage. There is anaesthesia or hypoaesthesia of the tail and hind legs and nearly always hypoaesthesia of the tail. The fur is thinned out to a very great extent, mostly on the legs and nates, to a less degree on the other parts of the hind quarters. In most cases the baldness was found to be diffused symmetrically. The animals are less lively and constantly sit quietly in a corner and sleep. Their appetite is diminished and a slow falling off in weight of some few g. a week is noticed in a few of them.

Therapeutic experiment.

As part of the attempt to find the causes of the neuropathic disturbances, therapeutic experiments were carried out with large doses of vitamins A, B, D and E. Animals in different stages were selected, as well as those which had shown the symptoms for various lengths of time. The details of dosage and the results obtained are shown in Tables II, III and IV.

Since all the observations indicated that animals that had been given a potent preparation of vitamin E were protected against paresis, it seemed also desirable to investigate whether the paretic syndrome was affected by wheat germ oil. Treatment was begun in four animals (Nos. 53, 90, 104, 105) 2-4 weeks after the paresis was observed. Unfortunately there was no more of the same wheat germ oil as was used earlier on in the experiment. The new preparation, which was also called "Fertilan" was prepared in a different way—not otherwise further elucidated [Vogt-Møller, 1934]. It was supposed to be of the same strength as the wheat germ oil which had been given previously in the experiment. From later experiments, however, it has shown itself to be ineffective

with regard to the preservation of pregnancy in doses of as much as 20 drops (500 mg.) daily.

As a result of these experiments we may conclude that large doses of vitamins A, B and D have no curative effect on the paresis, while nothing definite can be said as to the effect of vitamin E therapy.

DISCUSSION.

The clinical phenomena which have been described have a general similarity to the neuropathic disturbances in rats suffering from vitamin A deficiency described by Aberle [1934] and to the phenomenon in suckling rats from mothers on a diet low in vitamin E described by Evans and Burr [1928] but differ from them in essential points. The characteristic in the paralysis in the suckling young [Evans and Burr, 1928] is the initial stage with a sudden development of a spastic paralysis of the hind legs, feet and toes in a period of pronounced growth, from the 20th to the 25th days. The few that primarily showed a relaxed paresis also ended in a spastic condition.

All my paretic animals were almost adult when the paresis set in. The paresis was of the relaxed type. It developed slowly so that it was possible to differentiate the various stages. The paresis was accompanied by disturbances of the sensibility affecting especially the deep-seated and to a less degree the cutaneous sensibility of the legs and tail.

Since the vitamin E preparation given in our therapeutic experiment was shown to be quite inactive—in some control experiments—we are not able to compare the negative curative effect in our paretic animals with the beneficial effect obtained by Evans and Burr in some of their mildly affected animals.

The neuropathic disturbances described above bear a stronger resemblance to the neuropathic disturbances in rats fed on a diet low in vitamin A described by Aberle [1934].

In her animals the clinical phenomena appeared successively in the following order: spreading of the hind legs when in motion increasing to pronounced straddling, ataxia, later on spasticity and finally helpless trailing of the hind quarters and hind legs when walking. Aberle did not observe disturbances of sensibility or atrophy of the hind quarters or hind legs, or even atrophic disturbances of the skin or fur. On the other hand, Aberle observed incontinency of urine in several of her animals, a phenomenon which I have not observed in mine. On the contrary, I have got the impression that many of the paretic animals have had a greatly increased capacity of the bladder. Finally, the paresis in Aberle's animals was cured by large doses of vitamin A, a result which is quite different from my experience.

I wish to emphasise that the clinical phenomena in the skin and nervous system do not resemble those seen in rats with experimental beriberi [Hofmeister, 1922; Stern and Findlay, 1929] or rat pellagra [Goldberger and Lillie, 1926; Clemmensen, 1933]. The fact that the content of yeast in the food mixture corresponds with a daily intake of 0.7 g. dried yeast speaks strongly against the supposition of a lack of vitamin B₁ or B₂ as an aetiological factor.

Similarly lack of vitamins A and D may be left out of consideration since more than sufficient daily doses of these vitamins were given. The negative therapeutic result of greater doses of cod-liver oil points also in the same direction.

The following facts make it probable that lack of vitamin E is the cause of the paresis. Vitamin E is the only definitely known vitamin that was lacking in

the food mixture. If the animals in the three groups be compared, it is a striking fact that all the non-paralytic animals had an addition of vitamin E (wheat germ oil) during one or more pregnancies. None of the paretic animals was given wheat germ oil. They were given substances free from vitamin E or containing very little of it, far less than sufficient to permit of a successful pregnancy. All the paretic animals have, then, suffered for a long period of time—4–7 months—from a pronounced lack of vitamin E. The non-paretic animals were given vitamin E preparations relatively early and thereby replenished their deposits. By reason of this effect of protection against paresis by early dosing with vitamin E, in conjunction with the above-mentioned conditions, I consider that the origin of the paresis is lack of vitamin E of long standing and very high degree.

REFERENCES.

- Aberle (1934). *J. Nutr.* **7**, 445.
Clemmensen (1933). *Acta Pathol. Microbiol. Scand.* **10**, 304.
Cummings and Mattill (1930). *J. Nutr.* **3**, 421.
Evans and Bishop (1922). *Anat. Rec.* **23**, Proc. 17.
—— and Burr (1927). *Mem. Univ. Calif. Monograph*, 8.
—— (1928). *J. Biol. Chem.* **76**, 273.
Fridericia (1924). *J. Biol. Chem.* **62**, 471.
Goldberger and Lillie (1926). *Publ. Health Rep. Wash.* **41**, 1025 (quoted in Vitamins
(1932) *Med. Res. Council. Spec. Rep. Series*, No. 167).
Gudjonsson (1930, 1). *Dissertation. Acta Pathol. Microbiol. Scand. Suppl.* **4**, 1930.
—— (1930, 2). *Biochem. J.* **24**, 1591.
Hofmeister (1922). *Biochem. Z.* **129**, 477.
Hohlweg und Dohrn (1930). *Z. ges. exp. Med.* **71**, 762.
Mason (1933). *Amer. J. Anat.* **52**, 153.
Mattill and Crawford (1931). *Ind. Eng. Chem.* **22**, 341.
Stern and Findlay (1929). *J. Path. Bact.* **32**, 63.
Vogt-Møller (1931). *Hospitalstidende*, **74**, 567.
—— (1934). *Hospitalstidende*, **77**, 674.