mone. An increase in serum protein-bound iodine in humans due to the administration of iodide has been described by Danowski, Johnston & Greenman (1950). By extracting with butanol, Man *et al.* (1951) have been able to separate the protein-bound iodine fraction, which presumably represents hormone from the unknown protein-bound iodine compound and which appears during the administration of inorganic iodine. In future studies of the distribution of exogenous hormone in tissues it would probably be advisable to employ butanol extraction as an integral part of the method.

SUMMARY

1. A method for the determination of protein-bound iodine in extrathyroidal tissues is described. Data regarding the limits of error of the method are given.

2. In normal dogs the concentration of protein-bound iodine in the serum is higher than in skeletal

muscle and lower than in liver and kidney. No significant difference was found between the concentration of protein-bound iodine in the serum and that in the heart, lung, spleen, small intestine, pancreas and testis.

3. The feeding of powdered thyroid or iodide to normal dogs caused an increase in the concentration of protein-bound iodine of all tissues studied.

4. Protein-bound iodine is probably partly intracellular in the dog. In rat liver, protein-bound iodine occurs in approximately equal concentrations in the nuclear, mitochondrial, and supernatant (cytoplasm plus extracellular fluid) fractions.

This work was supported in part by a grant from the United States Public Health Service and in part by funds received from the Eugene Higgins Trust. We wish to acknowledge the co-operation of Dr A. K. Solomon of the Department of Biophysics, Harvard Medical School. Miss D. E. Graham gave most valuable technical assistance in this work over a period of 3 years.

REFERENCES

- Albert, A. & Keating, F. R., jun. (1949). *J. clin. Endocrin.* **9**, 1406.
 Albert, A., Rall, J. E., Keating, F. R., jun., Power, M. H. & Williams, M. M. D. (1949). *J. clin. Endocrin.* **9**, 1392.
 Barker, S. B. (1948). *J. biol. Chem.* **173**, 715.
 Carr, E. A., jun., Graham, D. E., Ober, S. & Riggs, D. S. (1950). *Science*, **111**, 552.
 Chaney, A. L. (1940). *Industr. Engng Chem. (Anal.)*, **12**, 179.
 Danowski, T. S., Johnston, S. Y. & Greenman, J. H. (1950). *J. clin. Endocrin.* **10**, 519.
 Gross, J. & Leblond, C. P. (1947). *J. biol. Chem.* **171**, 309.
 Hogeboom, G. H., Schneider, W. C. & Pallade, G. E. (1948). *J. biol. Chem.* **172**, 619.
 Justus, J. (1904). *Virchows Arch.* **176**, 1.
 Lipmann, F. A. (1951). Private communication.
 Man, E. B., Kydd, D. M. & Peters, J. P. (1951). *J. clin. Invest.* **30**, 531.
 Pratt, D. (1950). Private communication.
 Riggs, D. S. (1947). *Trans. Amer. Ass. Goiter*, p. 137.
 Riggs, D. S., Lavietes, P. H. & Man, E. B. (1942). *J. biol. Chem.* **143**, 363.
 Riggs, D. S., Stanbury, J. B. & Carr, E. A., jun. (1952). In preparation.
 Sandell, E. B. & Kolthoff, I. M. (1936). *Mikrochem. Acta*, **1**, 9.
 Taurog, A. & Chaikoff, I. L. (1946). *J. biol. Chem.* **163**, 323.
 Taurog, A. & Chaikoff, I. L. (1948). *J. biol. Chem.* **176**, 639.

Studies in Detoxication

48. URINARY METABOLITES OF 4:6-DINITRO-*O*-CRESOL IN THE RABBIT

By J. N. SMITH, R. H. SMITHIES AND R. T. WILLIAMS
Department of Biochemistry, St Mary's Hospital Medical School, London, W. 2

(Received 17 October 1952)

4:6-Dinitro-*o*-cresol (DNOC) has been used as an insecticide and fungicide for many years, and more recently has become important in agriculture as a selective weed killer (Shepard, 1951). Its use for this purpose has been the cause of several deaths and its former use in the treatment of obesity has been discontinued because of its high toxicity (Bidstrup & Payne, 1951; Horner, 1942).

Little is known about the metabolism of DNOC; only a small amount is eliminated unchanged in the urine, although it is readily absorbed (Spencer, Rowe, Adams & Irish, 1948). In man, it seems to act as a cumulative poison and is eliminated slowly over a period of weeks after ingestion (Harvey, Bidstrup & Bonnell, 1951; King & Harvey, 1952). In laboratory animals, however, there appears to be no

evidence of accumulation, although the DNOC eliminated in the urine is still only a small proportion of the dose (Parker, Barnes & Denz, 1951).

The toxic action of DNOC can probably be attributed to its blocking of 'oxidative phosphorylation' (Judah, 1952) and it is likely that two of its possible metabolites, 6-amino-4-nitro-*o*-cresol and 4-amino-6-nitro-*o*-cresol have no such inhibitory properties since the two corresponding metabolites of 2:4-dinitrophenol have no such action (Judah & Williams-Ashman, 1951). The object of the present work was to find out whether or not these compounds were metabolites of DNOC and to see if the formation of such metabolites could account for the reported low elimination of DNOC in the urine.

MATERIALS AND METHODS

Reference compounds. 4:6-Dinitro-*o*-cresol, DNOC (Pest Control Ltd.), had m.p. 86° after recrystallization from 95% ethanol and was chromatographically homogeneous. 6-Amino-4-nitro-*o*-cresol, m.p. 178°, was prepared according to Cazeneuve (1897). 6-Acetamido-4-nitro-*o*-cresol had m.p. 209° (see Friedländer, 1927-9). 4-Acetamido-6-nitro-*o*-cresol had m.p. 219° and on deacetylation gave 4-amino-6-nitro-*o*-cresol, m.p. 122° (Nietzki & Ruppert, 1890).

Since DNOC could undergo oxidation of its methyl group *in vivo*, the following derivatives of salicylic acid were prepared; 3:5-dinitrosalicylic acid, m.p. 172°, 3-amino-5-

nitrosalicylic acid, m.p. 229°, and 3-acetamido-5-nitrosalicylic acid, m.p. 246° (Meldola, Foster & Brightman, 1917).

Chromatography. Downward paper chromatograms were run in the usual way at room temperature in rectangular glass tanks closed at the top with plate-glass sheet. No special precautions were taken to make the tanks completely airtight or to equilibrate paper with the solvent. Under these conditions the R_F was not exactly the same in different tanks but the general pattern and order of separation were reproducible. All identification of compounds by their R_F values were by comparison with an authentic sample run on the same paper at the same time.

For most purposes Whatman no. 1 or 4 paper was used except where larger amounts were handled, when the thicker no. 3 was found more suitable.

Solvents were prepared by shaking the components together in the proportions by volume shown in Table 1 and the top layer used for chromatography. Values of R_F obtained with the different solvents are quoted in this table to illustrate the degree of separation achieved.

Colour reactions. The following reagents were used to locate compounds on the paper chromatograms (see also Table 2): 2N-NaOH gave yellow, pink or orange colours with nitro compounds. Ammoniacal AgNO₃, prepared from 10 ml. 0.1N-AgNO₃ and 1 drop 40% NaOH with enough dilute ammonia to give a clear solution, gave brown or black spots with aminophenols or di-amino compounds on standing at room temperature for a few minutes. Diazo reagents: a spray of 1% NaNO₂ in dilute acetic acid followed a few minutes later by a spray of 1% β -naphthol in N-NaOH,

Table 1. *Paper chromatography of DNOC and its possible metabolites. Relative R_F values with Whatman no. 1 paper**

	Benzene/acetic acid/water 1 : 1 : 2	<i>n</i> -Butanol/acetic acid/water 4 : 1 : 5	Benzene/ <i>n</i> - butanol/ammonia (sp.gr. 0.88) 2 : 5 : 2
	R_F	R_F	R_F
4:6-Dinitro- <i>o</i> -cresol	0.96	0.95	0.50
6-Acetamido-4-nitro- <i>o</i> -cresol	0.84	0.95	0.44
4-Acetamido-6-nitro- <i>o</i> -cresol	0.80	0.95	0.38
6-Amino-4-nitro- <i>o</i> -cresol	0.56	0.95	0.20
4-Amino-6-nitro- <i>o</i> -cresol	0.88	0.95	0.22
3-Amino-5-nitrosalicylic acid	0	0.64	0.13
3-Acetamido-5-nitrosalicylic acid	0	0.56	0.07
6-Acetamido-4-nitro- <i>o</i> -cresyl glucuronide	0	0.68	0.07

* Chromatograms run until front had moved 18 in. from the starting point.

Table 2. *Colour reactions of DNOC and related compounds*

Compound	Colour given in aqueous solution with		
	Aqueous FeCl ₃	N-NaOH	Diazo test*
DNOC	N.C.	Yellow	—
6-Amino-4-nitro- <i>o</i> -cresol	Solution darkens	Red	Yellow with HNO ₂ , no coupling
6-Acetamido-4-nitro- <i>o</i> -cresol	N.C.	Orange-red	Yellow with HNO ₂ , no coupling
4-Amino-6-nitro- <i>o</i> -cresol	N.C.	Pink	Red-violet slowly
4-Acetamido-6-nitro- <i>o</i> -cresol	N.C.	Orange-yellow	N.C.
3:5-Dinitrosalicylic acid	Red	Yellow	N.C.
3-Amino-5-nitrosalicylic acid	Red-brown	Orange	Yellow with HNO ₂ , violet on coupling
3-Acetamido-5-nitrosalicylic acid	Red-brown	Yellow	N.C.

* Diazo test is carried out by acidifying the solution with dilute HCl, adding a drop of 0.1% NaNO₂, followed by 1% ammonium sulphamate, then coupling with 1% ethanolic dimethyl α -naphthylamine. N.C. means no change on adding reagent.

gave red or red-violet colours with amino compounds, but with compounds containing the *o*-aminophenol structure, a yellow or brown colour appeared with the first spray. 10% NaCN: on spraying with NaCN and then heating to 100° for 10 min., DNOC and *m*-dinitro compounds gave pink colours.

Spectroscopy. Wherever amounts of material were sufficient, the spots were examined in the Unicam SP 500 spectrophotometer after elution by 0.1 N-NaOH or ethanol as previously described (Robinson, Smith & Williams, 1951). In many cases, however, the amount of material available on the chromatogram was very small and difficulty was experienced with a considerable irrelevant absorption at wavelengths below about 350–400 μ . This was not appreciably reduced by washing the paper, either in the chromatographic solvents or ethanol. As far as possible the effect of irrelevant absorption was minimized by eluting, with the same volume of eluent, a strip cut from the paper alongside the spot under investigation and using this in the compensating cell in the spectrophotometer. Absorbing material from the paper, however, may account for only part of the irrelevant ultraviolet absorption, for the extracts of rabbit urine contain much material not connected with DNOC which may underlie the yellow spots eluted. In general, using the methods of extraction described below, amounts of the order of 200 μ g. were found necessary to give satisfactory spectra after elution, since the eluate could then be diluted to reduce the relative amount of interfering absorption.

Wherever the spectra had characteristic bands at a wavelength not masked by the irrelevant absorption, identification of the eluted compounds was made by comparison with spectra of authentic material in 0.1 N-NaOH and 0.1 N-HCl.

Spectra of reference compounds and those of metabolites are summarized in Table 3 and typical identifications are shown in Figs. 1–3.

Animal experiments

Animals and dosing. All compounds were administered orally with water by stomach tube. In all, some forty chinchilla rabbits were given 20–30 mg./kg. of DNOC and survived. Two rabbits which had received respectively 31 and 32 mg./kg. of DNOC died within 24 hr. of dosing. Heymans & Casier (1935) have reported that rabbits tolerated subcutaneous injections of 10–20 mg./kg.

6-Amino-4-nitro-*o*-cresol and its *N*-acetyl derivative were administered in the same way but larger doses were tolerated, no deaths occurring after doses of 600 mg./kg. while some deaths occurred at a dose level of 700 mg./kg. of the amino compound.

Examination of DNOC urine. The urine was normal in appearance and did not react with Benedict solution or ammoniacal AgNO₃. The naphthoresorcinol reaction was not different from that of normal urine. In the diazo reaction a slight increase of yellow colour was noted on addition of HNO₃ and on coupling with dimethyl- α -naphthylamine a weak red colour developed. This diazo reaction was still positive 2 days after feeding and was detectable on the third and fourth days.

Since most of the material appeared to be present in the urine of the first 2 days, this was often pooled and worked up together.

Three extracts of the urine were made using peroxide-free ether in a continuous extractor.

Table 3. Absorption spectra of authentic DNOC and related compounds and those of various fractions from the urine of rabbits dosed with DNOC

Compound or extract	Absorption spectra					
	In 0.1 N-HCl		In 0.1 N-NaOH			
	$\lambda_{\max.}$ (μ .)	$\epsilon_{\max.}$ or $E_{\max.}$	$\lambda_{\max.}$ (μ .)	$\epsilon_{\max.}$ or $E_{\max.}$	$\lambda_{\max.}$ (μ .)	$\epsilon_{\max.}$ or $E_{\max.}$
DNOC						
Authentic	268	12 600	265	5 000	{ 370 ~ 408–410	12 800 approx. 10 000
From extract A	—	—	265	0.36	{ 370 ~ 408–410	0.61 0.48
From extract B	—	—	—	—	370	—
6-Acetamido-4-nitro- <i>o</i> -cresol						
Authentic	280	6 400	285	6 700	425	13 900
From extract A	—	—	285	0.35	425	0.39
6-Amino-4-nitro- <i>o</i> -cresol						
Authentic	{ 225 310	8 600 8 500	272	6 100	460	14 600
From extract B	—	—	272	0.47	460	0.56
From extract C	{ 225 310	0.37 0.35	272	0.31	460	0.64
3-Amino-5-nitrosalicylic acid						
Authentic	—	—	292.5	5 750	457.5	16 000
From extract B	—	—	~ 290	—	455	—

~ Denotes an inflexion.

Extract *A* was prepared from the urine which had been made acid (pH 2–5). The residual urine was then made 2*N* with H_2SO_4 , boiled for 3 hr., cooled and then extracted to give extract *B*. The hydrolysate was then neutralized to pH 7–8 and extracted again to give extract *C*. Each extract was reduced to a small volume on the water bath, taken up in ethanol and small portions used for the paper chromatograms.

Fig. 1. Absorption spectra of authentic DNOC in 0.1*N*-NaOH (broken line; ordinate scale, ϵ , on the right) and of DNOC obtained by paper chromatography of extract *A* of DNOC urine (full line; ordinate scale, $E_{1\text{ cm.}}$, on left).

Fig. 2. Absorption spectra of authentic 6-acetamido-4-nitro-*o*-cresol in 0.1*N*-NaOH (broken line; ordinate scale, ϵ , to the right) and of the same compound obtained by paper chromatography of extract *A* of DNOC urine (full line; ordinate scale $E_{1\text{ cm.}}$, to the left).

Extract *A*. This was found to contain unchanged DNOC and metabolites with an unconjugated OH group.

Unchanged DNOC was separated by chromatography in benzene/*n*-butanol/ammonia and identified by comparison with a reference spot of the authentic compound in this solvent and in *n*-butanol/acetic acid/water and benzene/*n*-butanol/ammonia, by its reaction with NaCN, by its yellow colour with 2*N*-NaOH, and by its absorption spectra after elution with ethanol (cf. Fig. 1). By calculation from the extinction of the absorption band in 0.1*N*-NaOH at 370 $m\mu$, it was found, in two experiments, that 3.5 and 4% of the

dose was eliminated as unchanged DNOC on the first day after feeding. A further 1.5% of the dose was excreted unchanged on the second day. Known amounts of DNOC applied to paper chromatograms and estimated in this way could be recovered quantitatively.

6-Acetamido-4-nitro-*o*-cresol was identified by comparing it chromatographically with an authentic sample in benzene/acetic acid/water, *n*-butanol/acetic acid/water and benzene/*n*-butanol/ammonia, by its yellow colour with 2*N*-NaOH and by its absorption spectrum after elution from the paper in 0.1*N*-NaOH (cf. Fig. 2). Its identity was further checked by hydrolysing a portion of the eluate by boiling in 2*N*-HCl for 2 hr. when the spectrum in 0.1*N*-NaOH corresponded with that of 6-amino-4-nitro-*o*-cresol.

Calculations from the extinctions in alkaline solution at 425 $m\mu$. showed that about 1.5% of the dose of DNOC was eliminated as unconjugated 6-acetamido-4-nitro-*o*-cresol on the first 2 days after dosing.

Fig. 3. Absorption spectra of authentic 6-amino-4-nitro-*o*-cresol in 0.1*N*-NaOH (broken line; ordinate scale, ϵ , to the right) and of the same compound obtained by paper chromatography of extract *C* of DNOC urine (full line; ordinate scale, $E_{1\text{ cm.}}$, to the left).

6-Amino-4-nitro-*o*-cresol also appeared to be present in small amounts in extract *A* since a spot with the same R_F value as the authentic material in benzene/acetic acid/water and in benzene/*n*-butanol/ammonia was obtained. This spot gave an orange colour with 2*N*-NaOH and a brown colour with HNO_3 . Insufficient amounts were present for spectroscopic identification.

Extract *B*. This contained compounds liberated from conjugates by hydrolysis and extractable from acid solution.

DNOC was identified, as in extract *A*, by colour reactions, R_F values and spectroscopy, and accounted for 0.7% of the dose as estimated by the extinction in alkaline solution at 370 $m\mu$.

3-Amino-5-nitrosalicylic acid also appeared to be present in some experiments, since a spot was obtained with the same R_F value as the authentic material, in benzene/*n*-butanol/ammonia, *n*-butanol/acetic acid/water and benzene/acetic acid/water and which gave the same red colour with 2*N*-NaOH. Two-dimensional chromatograms in the second and third of these solvents led to a clear separation of this compound from 6-amino-4-nitro-*o*-cresol which gives similar colour reactions. Elution of the spot in methanol and examination of its spectrum in 0.1*N*-NaOH showed that

it had a maximum at 450 $m\mu$. and an inflexion at 295 $m\mu$. The authentic compound had a maximum at 457.5 and 292.5 $m\mu$., but the amount of urinary material available was small and its spectrum was considerably distorted by irrelevant absorption.

Extract C. This contained compounds liberated from conjugates on hydrolysis, and extractable at neutral pH.

6-Amino-4-nitro-*o*-cresol was separated by chromatography in benzene/acetic acid/water and benzene/*n*-butanol/ammonia and identified by comparison with an authentic spot, by its brown colour in 2*N*-NaOH, by its brownish reaction with HNO_3 and by its ultraviolet spectrum in 0.1*N*-NaOH after elution with ethanol. This metabolite, which could arise by the hydrolysis of *O*-conjugated 6-acetamido or 6-amino-4-nitro-*o*-cresol, accounted for 9.0–10.6 % of the dose as determined spectroscopically in alkaline solution at 460 $m\mu$. (cf. Fig. 3).

4-Amino-6-nitro-*o*-cresol appeared also to be present in very small amounts, since a spot was found with the same chromatographic properties as the authentic material in benzene/butanol/ammonia and benzene/acetic acid/water. It gave a violet colour with 2*N*-NaOH, reduced ammoniacal $AgNO_3$ and gave a red-violet colour in the diazo reaction,

similar to those given by the authentic material. The amount present, however, was insufficient for spectroscopy.

In some experiments, this extract also appeared to contain some 3-amino-5-nitrosalicylic acid as indicated by separation of a spot with the same R_F value as the authentic material, a light brown colour with 2*N*-NaOH, a pink colour with HNO_3 , a purple-brown colour with 1 % $FeCl_3$ and an acid reaction to 1 % aqueous bromocresol purple. Amounts were insufficient for spectroscopic examination.

*The metabolites of 6-amino-4-nitro-*o*-cresol*

The cresol (0.6 g./kg.) was fed to a rabbit and a 24 hr. urine collected. Suitable small volumes (0.01–0.05 ml.) of the urine were chromatographed directly on the paper without previous extraction and the following compounds identified.

Unchanged 6-amino-4-nitro-*o*-cresol was identified by its chromatographic behaviour in benzene/acetic acid/water, *n*-butanol/acetic acid/water and benzene/*n*-butanol/water, by its brown colour in alkali and by its spectrum in 0.1*N*-NaOH after elution in 0.1*N*-NaOH. From the extinction at 460 $m\mu$. it was calculated that in different experiments 7–18 % of the dose was excreted unchanged over 3 days.

6-Acetamido-4-nitro-*o*-cresol was identified by its chromatographic behaviour in benzene/acetic acid/water, *n*-butanol/acetic acid/water and benzene/*n*-butanol/ammonia, by its yellow colour in alkali and by its spectrum in 0.1*N*-NaOH after elution from benzene/acetic acid/water chromatograms. From the value of the extinction at 425 $m\mu$. it was calculated that, in different experiments, from 1 to 7 % of a dose of 600 mg./kg. was excreted as this acetyl derivative.

The chromatograms also showed a large spot giving a yellow colour with 2*N*-NaOH. This was taken to be 6-amino-4-nitro-*o*-cresol glucuronide since it was present in the gum obtained by separating the glucuronides as the basic lead salts. It was not 6-acetamido-4-nitro-*o*-cresyl glucuronide (see below) since this does not give an immediate yellow colour with NaOH.

*The metabolites of 6-acetamido-4-nitro-*o*-cresol.* The urine (200 ml.) collected 24 hr. after feeding 2 g. of this compound was normal in appearance and gave a strong naphthoresorcinol reaction. The Benedict and diazo reactions were negative. Paper chromatography of the urine in benzene/acetic acid/water and *n*-butanol/acetic acid/water showed only two spots, one corresponding to the unchanged compound and the other to its glucuronide. The latter gave no immediate reaction with 2*N*-NaOH, but a yellow colour developed several hours after spraying the paper.

The glucuronic acid fraction of the urine was prepared through the basic lead salt and was obtained as a yellow syrup. This was neutralized with solid $NaHCO_3$, a little absolute ethanol was added and the mixture warmed. On cooling the hydrated sodium salt of 6-acetamido-4-nitro-*o*-cresyl glucuronide crystallized. It was recrystallized from ethanol/water to give 300 mg. of pale yellow needles decomposing at 160–170° and showing $[\alpha]_D^{20} - 29.2^\circ$ (in water, 2). (Found: C, 38.4; H, 5.8; N, 5.8; Na, 4.9. $C_{15}H_{17}O_{10}N_2Na \cdot 3.5H_2O$ requires C, 38.2; H, 5.9; N, 5.9; Na, 4.9 %. Light absorption: in 0.1*N*-HCl λ_{max} . 240, 290, ϵ_{max} . 8900, 5500; in 0.1*N*-NaOH λ_{max} . 245, 290, ϵ_{max} . 9200, 5500.)

DISCUSSION

Our experiments suggest that in 2 days after dosing with 20–30 mg./kg. of DNOC (I) rabbits excrete less than 20 % of the dose as metabolites in the urine. About 5 % of the dose is excreted as unchanged DNOC and a much smaller amount (about 1 %) as conjugated DNOC (II). The main urinary metabolites are derivatives of 6-amino-4-nitro-*o*-cresol

which account for about 11–12 % of the dose. Some 1–1.5 % of the dose is excreted as 6-acetamido-4-nitro-*o*-cresol (III) and about 10 % as *O*-conjugates (IV) of this compound. Other compounds which seem to be excreted in small amounts appear to be 3-amino-5-nitro-salicylic acid (V) and derivatives of 4-amino-6-nitro-*o*-cresol (VI).

These results may be compared with those of Parker *et al.* (1951) in rabbits and dogs, where a 4–10 % excretion of DNOC was found, and those of Harvey *et al.* (1951) in man where a 1–2 % excretion was found. Both these groups of workers estimated DNOC in urine by the method of Parker (1949). This method probably estimates some 6-acetamido-4-nitro-*o*-cresol as well as DNOC, since the acetamido compound possesses appreciable absorption at 430 m μ , the wavelength used in Parker's method, and is also extractable by ethyl methyl ketone, the solvent used. The absence of the Derrien (diaz) test in DNOC urine noted by Parker *et al.* (1951) and the weak diazo tests found by us suggest that the bulk of the aminonitrocresols excreted are acetylated.

From our results it appears that the main detoxication of DNOC in rabbits is by reduction of the 6-nitro group and not by conjugation of the phenolic hydroxyl of DNOC alone. Much more free than conjugated DNOC is excreted. The main detoxication product, 6-acetamido-4-nitro-*o*-cresol, is a relatively innocuous substance and is at least 20 times less toxic than DNOC.

It is interesting to note that, in the intact rabbit, the 6-nitro group, which is ortho to the phenolic group, appears to be more readily reduced than the 4-nitro group of DNOC. Similar observations have been made in intact animals on 2:4-dinitrophenol

(Parker, 1952) which is reduced in rats to 2-amino-4-nitrophenol, and on picric acid (Walko, 1901; Kozumi, 1914–15) which forms largely picramic acid (2-amino-4:6-dinitrophenol). Rat-liver suspensions, on the other hand, reduce 2:4-dinitrophenol to the other isomer, 4-amino-2-nitrophenol (Parker, 1952), which is apparently not formed in the intact animal.

SUMMARY

1. Using the techniques of paper chromatography and absorption spectroscopy, the urinary metabolites of 4:6-dinitro-*o*-cresol (DNOC) in rabbits have been examined.

2. After feeding doses of 20–30 mg./kg., less than 20 % of the dose could be accounted for as metabolites in the urine 2 days after dosing. Some 6–7 % was excreted as DNOC (about 5–6 % free and 1 % *O*-conjugated) and about 12 % as 6-acetamido-4-nitro-*o*-cresol (about 1.5 % free and 10–11 % *O*-conjugated). Traces of derivatives of 4-amino-6-nitro-*o*-cresol and 3-amino-5-nitro-salicylic acid were also excreted.

3. The conversion of DNOC into the 6-acetamido-4-nitrocresol resulted in a detoxication of DNOC since the latter compound is at least 20 times less toxic than DNOC.

4. 6-Amino-4-nitro-*o*-cresol and its *N*-acetyl derivative were also fed. The former was partly excreted unchanged and partly acetylated. The acetyl derivative formed a glucuronide which was isolated from the urine.

The expenses of this work were partly defrayed by a grant from the Medical Research Council. One of us (R. H. S.) is grateful to the Council for a studentship.

REFERENCES

- Bidstrup, P. L. & Payne, D. J. H. (1951). *Brit. med. J.* **2**, 16.
Cazeneuve, M. P. (1897). *Bull. Soc. chim., Paris*, **17**, (3) 204.
Friedländer, P. (1927–9). *Fortschr. Teerfarb.* **16**, 2611.
Harvey, D. G., Bidstrup, P. L. & Bonnell, J. A. L. (1951). *Brit. med. J.* **2**, 13.
Heymans, C. & Casier, H. (1935). *Arch. int. Pharmacodyn.* **50**, 20.
Horner, W. D. (1942). *Arch. Ophthalm., N.Y.*, **27**, 1097.
Judah, J. D. (1952). *Proc. R. Soc. Med.* **45**, 574.
Judah, J. D. & Williams-Ashman, H. G. (1951). *Biochem. J.* **48**, 33.
King, E. & Harvey, D. G. (1952). *Biochem. J.* **51**, vii.
Kozumi, T. (1914–15). *Zbl. Biochem. Biophys.* **17**, 615.
Meldola, R., Foster, H. S. & Brightman, R. (1917). *J. chem. Soc.* **111**, 533.
Nietzki, R. & Ruppert, F. (1890). *Ber. dtsch. chem. Ges.* **23**, 3477.
Parker, V. H. (1949). *Analyst*, **74**, 646.
Parker, V. H. (1952). *Biochem. J.* **51**, 363.
Parker, V. H., Barnes, J. M. & Denz, F. A. (1951). *Brit. J. industr. Med.* **8**, 226.
Robinson, D., Smith, J. N. & Williams, R. T. (1951). *Biochem. J.* **50**, 221.
Shepard, H. H. (1951). *The Chemistry and Action of Insecticides*. New York: McGraw-Hill Book Co. p. 281.
Spencer, H. C., Rowe, V. K., Adams, E. M. & Irish, D. D. (1948). *J. industr. Hyg.* **30**, 10.
Walko, K. (1901). *Arch. exp. Path. Pharmac.* **46**, 181.