urine can be removed by precipitation of the urine with barium chloride and may be attributed to phosphate and sulphate ions.

4. Total inhibition of limpet arylsulphatase by 0.2M-phosphate allows the use of the crude viscera

concentrates as β -glucuronidase preparations free from arylsulphatase.

5. The dissimilarity in the characteristics of arylsulphatases from various sources is commented on.

REFERENCES

- Butenandt, A. & Hoffstetter, H. (1939). *Hoppe-Seyl. Z.* **259**, 222.
- Clarke, W. G., Akawie, R. J., Pógrund, R. S. & Geissman, T. A. (1950). *18th Int. Congr. Physiol. Abstr.* p. 159.
- Cohen, H. & Bates, R. W. (1949). *Endocrinology*, **44**, 317.
- Cohen, H. & Bates, R. W. (1950). *Endocrinology*, **45**, 86.
- Dodgson, K. S., Lewis, J. I. M. & Spencer, B. (1952). *Biochem. J.* **51**, xlii.
- Dodgson, K. S., Lewis, J. I. M. & Spencer, B. (1953a). *Biochem. J.* **55**, 254.
- Dodgson, K. S. & Spencer, B. (1953). *Biochem. J.* **53**, 444.
- Dodgson, K. S., Spencer, B. & Thomas, J. (1953b). *Biochem. J.* **53**, 452.
- Dzialeszynski, L. M. (1947). *Nature, Lond.*, **160**, 464.
- Dzialeszynski, L. M. (1950). *Nature, Lond.*, **166**, 157.
- Noguchi, J. (1923). *Biochem. Z.* **143**, 186.
- Robinson, D., Smith, J. N., Spencer, B. & Williams, R. T. (1952). *Biochem. J.* **51**, 202.
- Samuels, L. T. & Reich, H. (1952). *Annu. Rev. Biochem.* **21**, 129.
- Seligman, A. M., Chauncey, H. H. & Nachlas, M. M. (1951). *Stain Tech.* **26**, 19.
- Soda, T. & Egami, F. (1941). *J. chem. Soc., Japan*, **62**, 256.
- Whitehead, J. E. M., Morrison, A. R. & Young, L. (1952). *Biochem. J.* **51**, 585.

The Oxidation of Glycerol by Mycobacteria

BY G. J. E. HUNTER

Travis Laboratory, Medical School, University of Otago, Dunedin, New Zealand

(Received 8 January 1953)

Nocard & Roux (1887) discovered that the growth of *Mycobacterium tuberculosis* var. *hominis* was stimulated by addition of glycerol to culture media, but Hammerschlag (1891) was the first to demonstrate utilization of glycerol. Several investigators have since shown that glycerol is a favourable source of carbon for the cultivation of many species of mycobacteria. Baudran (1910) drew attention to the simultaneous disappearance of glycerol and phosphoric acid from culture media, and suggested that glycerophosphoric acid might be formed. Merrill (1930, 1931a, b) studied the fate of glycerol in broth cultures and in synthetic media, and concluded that glycerol was completely oxidized without the accumulation of breakdown products, a view which was shared by Dingle & Weinzirl (1932), Wedum (1936) and Friedemann & Seibert (1939).

The investigation of washed cell suspensions has furnished additional information (see Edson, 1951). Nakamura (1938) reported that glycerol increased the oxygen consumption of *Myco. tuberculosis* var. *hominis*; but no direct relationship between respiration and the growth-promoting ability of glycerol could be found by Bloch, Matter & Suter (1947). Although it is clear that many acid-fast organisms oxidize glycerol, there has been little advance in knowledge of the intermediate metabolism. This

paper shows that the use of washed-suspension techniques and cell-free extracts has provided means of elucidating the mechanism of glycerol degradation.

METHODS

Organisms. The organisms used were *Myco. stercoris* (no. 3820), *Myco. smegmatis* (no. 523) and *Myco. butyricum* (no. 337). These strains were obtained from the National Collection of Type Cultures, Lister Institute, London. Stock cultures were maintained on glycerinated Dorset's egg medium, and cultivated on meat infusion broth containing 1% (w/v) peptonum B.P.C. (British Drug Houses) and 5% (v/v) glycerol. On the fifth or sixth day of incubation at 38°, the bacteria were washed and suspended in distilled water by methods described earlier (Edson & Hunter, 1943).

Bacterial extracts. Extracts were prepared by disrupting the organisms either in the wet crushing mill of Booth & Green (1938) or in the Mickle disintegrator (H. Mickle, Hampton, Middlesex). In the Mickle apparatus 4 g. of washed cells, 4 g. of no. 12 ballotini glass spheres (0.18–0.25 mm.) and 4 ml. of phosphate buffer (0.1M, pH 7.4) were mixed, precooled in each container and shaken at maximum amplitude for 30 min. in a cold room (2–3°). The suspension of disintegrated organisms and glass beads was diluted with 4 ml. of buffer per tube, the extracts clarified by centrifugation and the supernatant dialysed for 24–48 hr. against distilled water at 2–3°.

Manometric measurements. Oxygen consumption was measured by the Warburg technique at 38°, and the method of Warburg & Yabusoe (1924) was used for the determination of the respiratory quotient (R.Q.). Substrates were added from the side bulb after equilibration, and inhibitors were in contact with bacteria from the start of an experiment. The CO₂ absorbent consisted of appropriate mixtures of KOH and KCN (Krebs, 1935) when HCN was used as inhibitor. Acid substrates were used as sodium salts.

Materials. Adenosine triphosphate (ATP) was a purified sample supplied by Boots Pure Drug Co. Ltd. The barium salt was about 94% pure as judged by 7-min. P, 7-min. P/total P and P/N values. A typical analysis of the solution (sodium salt) used in experiments gave: inorganic P, <0.05 mg.; acid-labile 7-min. P, 0.71 mg.; total P, 1.10 mg.

Diphosphopyridine nucleotide was obtained from Schwartz Laboratories Inc., New York. Its purity (50%) was assayed by the method of LePage (1947), using the value $E_{1\text{cm}}^{1\%} = 8500$ at 340 m μ .

Analytical methods

Estimation of glycerol. At the end of an experiment with intact cells, the Warburg flasks were cooled in ice and the contents quantitatively transferred to a graduated test tube by means of a Pasteur pipette. Deproteinizing reagent (0.4 ml. of 7% (w/v) metaphosphoric acid solution) was added, the volume adjusted to 10 ml. with distilled water, the tubes centrifuged and 2.0 ml. portions of supernatant used for analysis by the periodate method of Voris, Ellis & Maynard (1940).

Determination of phosphate and phosphate esters. After experiments with clarified and dialysed extracts, the flasks were cooled in ice, 1 ml. of 30% (w/v) trichloroacetic acid was added and the volume made up to 10 ml. with distilled water. After centrifugation 2 ml. portions of the supernatant were used for phosphorus determinations. Triose phosphate was determined as alkali-labile P (Lohmann & Meyerhof, 1934) and ATP as readily hydrolysable P (7 min., 1.0N-HCl, 100°). For total-P determinations the wet incineration was carried out by heating 1 ml. of P-containing solution with 0.5 ml. of H₂SO₄ (sp.gr. 1.84) and 0.5 ml. of HNO₃ (sp.gr. 1.42) in a micro-Kjeldahl flask. Additional HNO₃ was added if required. All phosphate determinations (inorganic P, hydrolysable P and total P) were performed by the method of Fiske & Subbarow (1925). Phosphoenolpyruvic acid and pyruvic acid were measured by the methods of Lohmann & Meyerhof (1934) and Friedemann & Haugen (1943) respectively.

RESULTS

Respiration

The oxygen consumption of *Myco. smegmatis*, *Myco. stercoris* and *Myco. butyricum* in the presence and absence of relevant substrates is shown in Table 1. The endogenous respiration of each organism is substantial and fairly constant. For a series of different cultures the mean Q_{O_2} ($Q_{O_2} = \mu\text{l. O}_2$ consumed/mg. dry wt./hr.) values (with s.d.) in the absence of added substrate were 13.4 ± 0.9 (twenty-two cultures) for *Myco. smegmatis*, 10.1 ± 0.6 (nine cultures) for *Myco. stercoris*

and 13.7 ± 0.8 (twenty-three cultures) for *Myco. butyricum*. In the presence of glycerol the oxygen consumption was doubled. Working with a strain of *Myco. tuberculosis* var. *hominis* virulent for guinea pigs, Bloch (1944) and Bloch *et al.* (1947) reported that the augmentation of oxygen uptake due to glycerol was little affected by the pH of the medium. *Myco. smegmatis* and *Myco. butyricum* also behave in the same way. When the medium was buffered at pH 7.8, 6.8 or 6.2, there was little effect on the extra oxygen uptake due to glycerol, although the endogenous respiration was diminished at the lower pH.

Table 1. *Oxygen consumption of mycobacteria in the presence of various substrates*

(Washed bacteria, 12.0–13.0 mg. dry wt./cup suspended in 3.3 ml. of 0.03M-phosphate buffer, pH 7.4, at 38°; gas, air. $Q_{O_2} = \mu\text{l. O}_2$ uptake/mg. dry wt. bacteria/hr.)

Substrate (0.02M unless stated otherwise)	Q_{O_2}		
	<i>Myco. smegmatis</i>	<i>Myco. stercoris</i>	<i>Myco. butyricum</i>
Nil	15.0	11.9	13.4
Glycerol	32.0	20.5	25.3
DL-Glyceraldehyde	54.8	42.8	21.9
Dihydroxyacetone	41.1	22.8	18.0
Glycerate	32.7	24.3	17.4
α -Glycerophosphate (0.01M)	18.5	13.8	13.6
D-3-Phosphoglycerate (0.01M)	15.0	12.7	12.9
Pyruvate	33.2	23.6	25.0
Acetate	38.7	23.1	33.8
Glucose	20.0	15.9	16.6

Oxygen consumption is considerably increased by the presence of DL-glyceraldehyde, dihydroxyacetone, pyruvate and acetate. Although the suspensions do not oxidize α -glycerophosphate or phosphoglycerate significantly, it will be shown later that cell extracts are able to bring about the degradation of these substances. Bacterial suspensions which had been heated in a boiling-water bath for 10 min. did not take up oxygen either in the presence or absence of glyceraldehyde or dihydroxyacetone. Hence the high oxygen consumption in the presence of these relatively unstable substrates is not due to spontaneous oxidation.

Disappearance of glycerol in relation to oxygen uptake. The relationship between oxygen consumption and the disappearance of glycerol was investigated by two methods: (1) The rate of oxygen uptake in washed suspensions was measured in the presence of small amounts of glycerol and compared with the endogenous respiration. The resulting curves for *Myco. smegmatis* are given in Fig. 1. It will be seen that the graphs are rectilinear up to the point where glycerol disappears completely from the medium, and thereafter become parallel with the curve for the control suspension to which sub-

strate was not added. The excess oxygen uptake at the point where the rate of gas absorption became equal to the endogenous rate was used in calculating the ratio oxygen consumed/glycerol disappearing. In Fig. 1 the molar ratios of oxygen consumed/glycerol disappearing are 1.6, 1.5 and 1.4 for curves A, B and C respectively. *Myco. butyricum* gave

Fig. 1. Oxygen consumption of washed suspension of *Myco. smegmatis* (9.0 mg. dry wt. bacteria/3.4 ml. 0.03M-phosphate buffer, pH 7.4, at 38°; gas, air). Initial quantity of glycerol: A, 4.0 μ moles; B, 8.0 μ moles; C, 16.0 μ moles; D, 40.0 μ moles.

similar curves with the same amounts of glycerol, the molar ratios being 1.0, 1.1 and 1.2 respectively. (2) Residual glycerol was estimated after suspensions of the organisms had respired for a fixed time in a buffered medium containing excess of glycerol (Table 2).

The results of both methods agree in demonstrating that glycerol is incompletely oxidized, the molar ratio of oxygen uptake/glycerol consumed being less than half the theoretical value for complete oxidation in one case (*Myco. smegmatis*) and less than a third in the other (*Myco. butyricum*). The assimilation of glycerol resembles that of glucose, lactate (Edson & Hunter, 1947), pyruvate

and fatty acids (Lindsay, O'Donnell & Edson, 1950), all of which are incompletely oxidized by washed suspensions of mycobacteria.

Oxygen consumption and carbon dioxide production. Table 3 records the results obtained when the oxygen consumption and carbon dioxide production of washed suspensions were measured in the presence and absence of glycerol. The 'extra' carbon dioxide/oxygen ratios, which are lower than the theoretical R.Q. for complete combustion of glycerol, suggest that oxidative assimilation may lead to the formation of carbohydrate and/or fatty acids.

Influence of inhibitors on respiration. Table 4 shows the effect of various reagents on the oxidation of glycerol by *Myco. smegmatis*, *Myco. stercois* and *Myco. butyricum*. Typical data illustrating the influence of fluoride and fluoroacetate on the oxidation of glycerol, pyruvate and acetate are recorded in Table 5. The inhibition of respiration referred to below is calculated on the basis of oxygen uptake in excess of that due to endogenous respiration.

It will be seen that the endogenous respiration is strongly inhibited by cyanide and arsenite, the profound effect of cyanide on the oxidation of glycerol indicating that the oxidation may be linked to the cytochrome system. Arsenite blocks the oxidation of pyruvate almost completely. It also inhibits the oxidation of glycerol by 72 to 80% in *Myco. smegmatis* and *Myco. stercois*, and to a lesser degree (49 to 62%) in *Myco. butyricum*. Glyceraldehyde oxidation is likewise sensitive to arsenite, the effect being more pronounced in *Myco. butyricum*. Iodoacetate inhibits glycerol oxidation by 63–80%. Malonate has no significant effect. Fluoride, which has no inhibitory action at pH 7.4, strongly inhibits the oxidation of glycerol at pH 6.0 in *Myco. smegmatis*, but does not affect this oxidation in *Myco. butyricum*. Fluoride also affects the metabolism of pyruvate and acetate with results which depend on pH and on the species of organism (cf. the observation of Fitzgerald & Bernheim (1948) on the tubercle bacillus). In *Myco. smegmatis* fluoride causes partial inhibition of pyruvate (24%) and acetate (62%) oxidation at pH 6.0 but not at pH 7.4; whereas in *Myco. butyricum* at pH 6.0 there is partial inhibition of acetate oxidation (21%) and a slight stimulation of pyruvate oxidation. These differences in the sensitivity of species to selective metabolic inhibitors are well illustrated in the case of fluoroacetate, the action of which is not greatly affected by pH.

The interpretation of these facts is difficult, but taken together they suggest that glycerol may be phosphorylated and broken down to pyruvate via the relevant portion of the Embden-Meyerhof scheme.

Table 2. *Oxidation of glycerol by washed suspensions of mycobacteria*

(Bacteria suspended in 3.4 ml. 0.03 M-phosphate buffer, pH 7.4, at 38°. Gas, air.)

Time (min.)	Initial glycerol (μmole)	Oxygen uptake (μmole)	'Extra' oxygen consumed (μmole)	Glycerol disappearing (μmole)	μmole 'extra' oxygen used / μmole glycerol disappearing
<i>Myco. smegmatis</i> (7.0 mg. dry wt.)					
60	0.00	8.40			
	79.43	20.76	12.36	8.98	1.4
120	0.00	15.90			
	79.18	41.57	25.67	16.87	1.5
180	0.00	22.77			
	78.79	63.89	41.12	27.50	1.5
<i>Myco. butyricum</i> (15.7 mg. dry wt.)					
60	0.00	7.72			
	78.24	15.00	7.28	7.22	1.0
120	0.00	15.31			
	77.75	31.30	15.99	13.07	1.2
180	0.00	22.32			
	78.61	47.73	25.41	21.60	1.2
Theoretical ratio for complete combustion					3.5

Table 3. *Oxygen consumption and carbon dioxide production by washed suspensions of mycobacteria*(Medium: 3.3 ml. 0.03 M-phosphate buffer, pH 7.4, at 38°. Total CO₂ (free and bound) measured by the method of Warburg & Yabuseo (1924).)

Glycerol (0.02 M)	Oxygen uptake (μl.)	Total CO ₂ (μl.)	CO ₂ /O ₂	'Extra' CO ₂ / 'Extra' O ₂
<i>Myco. smegmatis</i> (11.0 mg. dry wt./cup; time 120 min.)				
Absent	220	215	0.98	—
Present	447	387	—	0.76
<i>Myco. butyricum</i> (15.0 mg. dry wt./cup; time 75 min.)				
Absent	203	183	0.90	—
Present	392	308	—	0.66
Theoretical R.Q.				0.86

Table 4. *Effect of reagents on the oxidation of glycerol and other substrates by mycobacteria*

(Washed suspensions (10–18 mg. dry wt. bacteria) in 3.5 ml. 0.03 M-phosphate buffer, pH 7.4. Gas, air. Time 60–120 min. in different experiments.)

	<i>Myco. smegmatis</i> Substrate		<i>Myco. stercoris</i> Substrate		<i>Myco. butyricum</i> Substrate	
	Absent	Present	Absent	Present	Absent	Present
Substrate, glycerol (0.02 M)						
Nil	11.3	28.1	12.6	26.8	15.6	26.6
Arsenite (0.5 × 10 ⁻³ M)	6.4	10.9	7.9	10.8	6.7	12.3
Arsenite (1.0 × 10 ⁻³ M)	6.1	10.8	—	—	6.3	10.5
Nil	11.3	28.1	10.5	20.5	12.1	29.9
Iodoacetate (1.0 × 10 ⁻³ M)	11.8	17.4	9.9	13.5	7.6	9.7
Nil	9.3	21.3	8.4	11.8	14.9	27.3
HCN (1.0 × 10 ⁻³ M)	7.5	8.9	4.3	4.9	5.0	5.2
HCN (1.0 × 10 ⁻³ M)	3.5	3.6	—	—	3.4	3.5
Nil	14.8	31.2	10.5	20.5	12.1	29.9
Malonate (1.0 × 10 ⁻³ M)	15.0	32.3	11.0	19.4	11.3	31.8
Substrate, glyceraldehyde (0.02 M)						
Nil	11.3	41.7	—	—	15.6	24.2
Arsenite (0.5 × 10 ⁻³ M)	6.4	19.3	—	—	6.7	7.6
Arsenite (1.0 × 10 ⁻³ M)	6.1	19.6	—	—	6.3	6.6
Substrate, pyruvate (0.02 M)						
Nil	15.9	37.6	—	—	11.8	27.2
Arsenite (0.5 × 10 ⁻³ M)	6.9	8.4	—	—	5.3	7.1
Arsenite (1.0 × 10 ⁻³ M)	6.8	8.5	—	—	5.3	7.2
						21.2

Table 5. *Influence of fluoride and fluoroacetate on the oxidation of glycerol, pyruvate and acetate by mycobacteria*

(Washed suspensions (13–17 mg. dry wt. bacteria) in 3.5 ml. 0.03 M-phosphate buffer, pH's 7.4 and 6.0. Gas, air. Time 60–120 min. in different experiments.)

Inhibitor (0.01 M)	<i>Myco. smegmatis</i>				<i>Myco. butyricum</i>			
	pH 7.4 Substrate		pH 6.0 Substrate		pH 7.4 Substrate		pH 6.0 Substrate	
	Absent	Present	Absent	Present	Absent	Present	Absent	Present
Substrate, glycerol (0.02 M)								
Nil	14.5	37.2	14.4	34.3	17.8	19.7	13.6	21.2
NaF	15.3	36.8	10.0	11.2	18.4	20.4	11.7	19.2
Nil	16.9	31.6	10.7	31.6	16.9	24.2	14.6	19.6
CH ₂ FCOONa	18.3	37.5	16.2	36.2	10.2	12.9	7.9	8.2
Substrate, pyruvate (0.02 M)								
Nil	13.5	33.5	10.0	33.7	20.2	35.2	17.8	33.0
NaF	14.1	35.4	8.8	26.9	21.0	35.6	18.4	38.9
Nil	16.9	37.5	10.7	31.4	16.9	29.6	14.6	29.7
CH ₂ FCOONa	18.3	32.6	16.2	31.6	10.2	11.6	7.9	10.8
Substrate, acetate (0.02 M)								
Nil	15.0	29.4	13.5	21.3	16.5	46.2	13.6	39.8
NaF	14.8	32.8	9.8	12.7	16.8	47.0	11.7	32.4
Nil	16.9	37.7	10.7	22.9	16.9	37.7	14.6	31.1
CH ₂ FCOONa	18.3	31.0	16.2	23.4	10.2	9.8	7.9	6.7

Products of oxidation

The fact that the Q_{O_2} values for the oxidation of pyruvate and glycerol are approximately equal (Table 1) is consistent with the suggestion that pyruvate is an intermediate in the aerobic breakdown of glycerol. If this is the case, pyruvate should accumulate when glycerol is oxidized in the presence of arsenite which inhibits pyruvate metabolism (Table 4). The Rothera test was applied to the contents of Warburg flasks (Table 4) after the bacteria and soluble protein had been removed by addition of 0.1 vol. of 30% (w/v) trichloroacetic acid. Pyruvate (blue ring) accumulated when glycerol or glyceraldehyde was metabolized in the presence, but not in the absence, of arsenite. The test was negative when the metabolism of glycerol was blocked by fluoride or iodoacetate (Table 5). These facts are consistent with the working hypothesis stated above.

Although the endogenous respiration is inhibited by arsenite, pyruvate did not accumulate in heavy suspensions of *Myco. smegmatis*; on the other hand, small amounts of pyruvic acid and traces of α -ketoglutaric acid (identified chromatographically (Cavallini, Frontali & Toschi, 1949) after crystallization of the 2,4-dinitrophenylhydrazones) were formed endogenously by *Myco. butyricum* respiring in the presence of arsenite. This difference is reflected in the cultural behaviour of the two organisms (Hunter, 1952). In washed suspensions of both species the yield of pyruvic acid was considerably increased when glycerol was metabolized in the presence of arsenite.

Experiments with bacterial extracts

Clear cell-free extracts of the bacteria were used to elucidate the intermediate steps in the breakdown of glycerol to pyruvic acid.

Anaerobic reduction of methylene blue. The possibility of glycerophosphate being an intermediate was tested by the Thunberg technique with methylene blue as hydrogen acceptor. Table 6 shows that dialysed extracts of *Myco. smegmatis* and *Myco. butyricum* buffered with inorganic phosphate do not oxidize glycerol unless ATP is added, suggesting that the first step in glycerol metabolism is a phosphorylation. This notion is supported by the fact that α -glycerophosphate is oxidized anaerobically in the absence of ATP (Table 6). In some extracts subjected to prolonged dialysis the oxidation of α -glycerophosphate was dependent on the addition of magnesium ions but the effect was not observed regularly. Control extracts which had been heated in a water bath for 15 min. at 100° did not reduce methylene blue in the presence or absence of substrates.

Since α -glycerophosphate is probably oxidized to triose phosphate, an attempt was made to detect triose phosphate by the sensitive method of Sibley & Lehninger (1949). At the end of an experiment the solutions were deproteinized with trichloroacetic acid and the methylene blue removed by filtration with charcoal. Several brands of charcoal produced precipitates which interfered with the colour test, but iron-free 'Norite A' (Pfanstiehl Chemical Co.) and Merck's medicinal charcoal gave satisfactory clear filtrates. The test reagents were added to the

Table 6. *Anaerobic triose phosphate formation from glycerophosphate and glycerol*

(Evacuated Thunberg tubes contained 2.0 ml. 0.1 M-phosphate buffer, pH 7.4 and 0.5 ml. methylene blue (1:1000) at 38°.)

		Dialysed cell-free extracts from								
Additions (ml.)		<i>Myco. smegmatis</i>					<i>Myco. butyricum</i>			
Extract		1.0	1.0	1.0	1.0	1.0	0.2	0.2	0.2	0.2
α -Glycerophosphate (0.1 M)		0.3	0.3	—	—	—	0.3	0.3	—	—
Neutralized hydrazine hydrochloride (M)		—	0.2	0.2	—	0.2	—	0.2	0.2	0.2
Glycerol (0.2 M)		—	—	0.3	0.3	0.3	—	—	0.3	0.3
ATP (0.04 M)		—	—	—	0.2	0.2	—	—	—	0.2
Water		0.2	—	0.2	0.2	—	0.2	—	0.2	—
Reduction time (min.)		60	55	>300	100	75	>300	85	>300	65
Sibley-Lehninger Test		N.	P.	N.	N.	P.	N.	P.	N.	P.

N. = negative; P. = positive.

Table 7. *Oxygen consumption of dialysed extracts from Mycobacterium smegmatis*

(Each Warburg cup contained 2.5 ml. 0.03 M-phosphate buffer, pH 7.4. Temp., 38°. Gas, air. Time, 100 min. 1 ml. bacterial extract added from side bulb after equilibration.)

Additions	Oxygen uptake (μ l.)	Sibley-Lehninger test
Nil	15	Negative
Neutralized hydrazine hydrochloride (0.05 M)	23	Negative
Methylene blue (0.003 %)	20	Negative
Hydrazine (0.05 M); methylene blue (0.003 %, w/v)	28	Negative
α -Glycerophosphate (0.01 M)	24	Negative
α -Glycerophosphate (0.01 M); diphosphopyridine nucleotide (0.1 mg., 50 % purity)	14	—
α -Glycerophosphate (0.01 M); hydrazine (0.05 M)	33	Trace
α -Glycerophosphate (0.01 M); methylene blue (0.003 %, w/v)	80	Negative
α -Glycerophosphate (0.01 M); methylene blue (0.003 %, w/v); diphosphopyridine nucleotide (0.1 mg. 50 % purity)	77	—
α -Glycerophosphate (0.01 M); methylene blue (0.003 %, w/v); hydrazine (0.05 M)	90	Positive

water-clear filtrates. When glycerol was the substrate, it was necessary to keep the concentration of magnesium ions to a minimum since precipitation of the hydroxide occurred on making the solution alkaline in the final stages of the Sibley-Lehninger test. Table 6 shows the anaerobic formation of triose phosphate from glycerol and glycerophosphate, the triose ester accumulating only when hydrazine was used as a fixative. The colour test is not specific for triose phosphate but is also given by ketonic acids. Since the nitroprusside test for pyruvate was negative, it is probable that the positive Sibley-Lehninger tests indicated the presence of triose phosphate.

Aerobic oxidation of glycerophosphate with methylene blue as hydrogen carrier. Dialysed extracts showed a small spontaneous oxygen uptake in the absence of added substrate. This blank was slightly increased by α -glycerophosphate (Table 7), but when methylene blue was added the oxygen consumption was increased fourfold. When the substrate was oxidized in the presence of hydrazine, triose phosphate was detectable by the Sibley-Lehninger test applied to the fluid from the manometric cups (after deproteinization and removal of

methylene blue). The slow rate of oxygen consumption was not accelerated by the addition of diphosphopyridine nucleotide (Table 7).

When the experiments were repeated on a larger scale, the triose phosphate was determined as alkali-labile phosphate in deproteinized filtrates from which the methylene blue was removed by adsorption on charcoal as before (Table 8). Again it will be seen that triose phosphate accumulates only when the artificial hydrogen carrier is present and hydrazine is used as a fixative. Control experiments with boiled bacterial extracts and the requisite additions gave no alkali-labile phosphate; and analytical controls showed that hydrazine did not interfere with the Fiske and Subbarow phosphate determinations. It will be noticed (Table 8) that free inorganic phosphate increased during the experiment, especially in the presence of hydrazine. Since this did not occur with boiled extracts, the effect is attributed to phosphatase action.

Breakdown of phosphoenolpyruvate and D-3-phosphoglycerate. Dialysed extracts of *Myco. smegmatis* and *Myco. butyricum* are capable of dephosphorylating phosphoenolpyruvic acid (Table 9). The breakdown of D-3-phosphoglycerate was studied

Table 8. *Formation of triose phosphate from glycerophosphate*

(Clarified and dialysed bacterial extract (4 ml.) shaken in stoppered 50 ml. conical flasks for 3 hr. at 38°, unbuffered initial pH 6.8.)

Additions (total volume made up to 6 ml. with distilled water)	<i>Myco. smegmatis</i>		<i>Myco. butyricum</i>	
	Inorganic P	Alkali-labile P (μ moles/flask)	Inorganic P	Alkali-labile P
Nil	0.3	0.0	5.8	0.3
1 ml. 0.1 M- α -glycerophosphate (100 μ moles)	3.2	2.3	6.7	1.3
1 ml. 0.1 M- α -glycerophosphate (100 μ moles) + 0.4 ml. 1.0 M-hydrazine hydrochloride (neutralized)	8.7	2.3	11.9	2.3
1 ml. 0.1 M- α -glycerophosphate (100 μ moles) + 0.5 ml. 1:200 methylene blue	3.8	2.9	7.1	0.3
1 ml. 0.1 M- α -glycerophosphate (100 μ moles) + 0.4 ml. 1.0 M-hydrazine + 0.5 ml. methylene blue	8.8	23.2	10.0	11.6

in the presence and absence of fluoride by measuring simultaneously the formation of phosphoenolpyruvate and the release of inorganic phosphate and pyruvic acid (Table 10). The degradation of phosphoglycerate does not occur unless magnesium ions are added and it is inhibited by fluoride, showing that the enzyme concerned has the characteristic properties of enolase. The amount

Table 9. *Enzymic cleavage of phosphopyruvate*

(4.0 ml. dialysed extract and 1.0 ml. 0.1 M-phosphoenolpyruvate (100 μ moles) incubated in stoppered 50 ml. conical flasks for 3 hr. at 38°, pH 6.8.)

	<i>Myco. smegmatis</i> (μ moles/flask)	<i>Myco. butyricum</i> (μ moles/flask)
Initial inorganic P	0.0	0.0
Final inorganic P	3.5	2.2
Pyruvic acid	2.8	1.9

of phosphoenolpyruvate accumulating is small in relation to the amount of pyruvate liberated. Control experiments showed that the disappearance of pyruvate was not significant under the conditions used. The activity of extracts was variable.

Phosphorylation of glycerol. When neutral unbuffered extracts are incubated with ATP, phosphate is liberated, the enzymic hydrolysis being complete in 3 hr. with *Myco. smegmatis* but incomplete with *Myco. butyricum*. Spontaneous liberation of phosphate was negligible when ATP was incubated with heat-inactivated extracts. Although the amount of inorganic phosphate set free is substantially reduced when glycerol is added to extracts of *Myco. smegmatis*, the residual ATP (7-min. acid-labile P) is negligible (Table 11). The change in the distribution of phosphate can be explained if glycerophosphate is formed. The esterification of glycerol is slower in extracts of *Myco. butyricum*. This interpretation of ATP breakdown is consistent with the observations which show that glycerophosphate is oxidized in extracts, whereas glycerol is not oxidized unless ATP is added (Table 6). Since the transphosphorylation experiments were performed in the absence of methylene blue, significant amounts of triose phosphate were not formed (Tables 7 and 8). Hence triose ester did not interfere with the determination of residual 7-min. acid-labile P. The inorganic phosphate content of the bacterial extracts was too small to be determined by the method used.

Table 10. *Enzymic breakdown of phosphoglycerate*

(Dialysed extract and substrate incubated in stoppered 50 ml. conical flasks for 3 hr. at 38°, pH 6.8. A control experiment without MgSO_4 gave zero results for all three types of P measured and for pyruvic acid.)

Contents (ml.)	Flask 1		Flask 2	
D-3-Phosphoglycerate 100 μ moles/ml.	1.0		1.0	
Bacterial extract	4.0		4.0	
MgSO_4 (0.04 M)	0.1		0.1	
NaF (0.2 M)	—		0.5	
Water	0.5		—	
	<i>Myco. smegmatis</i> (μ moles/flask)	<i>Myco. butyricum</i> (μ moles/flask)	<i>Myco. smegmatis</i> (μ moles/flask)	<i>Myco. butyricum</i> (μ moles/flask)
Initial inorganic P	0.0	0.0	0.0	0.0
Final inorganic P	46.0	9.7	4.5	0.0
Phosphopyruvate P	2.3	2.6	0.1	0.0
Pyruvic acid	42.0	9.6	0.1	0.0

Table 11. *Phosphorylation of glycerol*

(Reaction mixtures in 50 ml. conical flasks for 3 hr. at 38°.)

Contents (ml.)	Flask 1		Flask 2	
Glycerol (200 μ moles/ml.)	—		1.0	
Bacterial extract	4.0		4.0	
ATP (53.2 μ moles acid-labile 7-min. P/ml.)	0.4		0.4	
Water	1.0		—	
	<div> <div><i>Myco. smegmatis</i></div> <div><i>Myco. butyricum</i></div> <div>(μmoles/flask)</div> </div>		<div> <div><i>Myco. smegmatis</i></div> <div><i>Myco. butyricum</i></div> <div>(μmoles/flask)</div> </div>	
Initial inorganic P	1.6	1.6	1.6	1.6
Final inorganic P	22.9	16.1	11.6	9.7
Final acid-labile 7-min. P + inorganic P	22.9	22.9	12.6	18.4
Initial acid-labile 7-min. P	21.3	21.3	21.3	21.3
ATP disappearing (as acid labile 7-min. P)	21.3	14.5	20.3	12.6
Glycerophosphate P	0.0	0.0	10.3	4.5

DISCUSSION

Glycerol is readily metabolized by washed suspensions of the mycobacteria used in this investigation. The oxidation is incomplete and accompanied by assimilation (Fig. 1, Tables 2 and 3); and yet significant amounts of intermediate metabolites do not appear in the suspension medium. In these respects the metabolism of glycerol conforms to a pattern characteristic of the acid-fast bacteria.

Arsenite, which blocks the metabolism of pyruvate, interferes with glycerol oxidation (Table 4), reducing oxygen consumption by more than 70 % and causing accumulation of pyruvate in the surrounding medium. Iodoacetate and fluoride, which also inhibit respiration in the presence of glycerol, operate at different intermediate stages since pyruvate does not accumulate extracellularly under their influence (Tables 4 and 5). These facts are compatible with the view that glycerol is normally broken down to pyruvate by a mechanism which involves phosphorylation; and they are not inconsistent with the meagre increase in oxygen uptake which occurs in the presence of α -glycerophosphate and D-3-phosphoglycerate (Table 1), because these substrates are known to penetrate cells with difficulty. Dialysed extracts, free from cells and insoluble cell debris, can phosphorylate glycerol (Tables 6 and 11) in the presence of ATP and can break down certain phosphorylated intermediates of carbohydrate metabolism. The activity of these extracts is weak and somewhat variable and usually lower with *Myco. butyricum* than with *Myco. smegmatis*. The extracts contain an α -glycerophosphate dehydrogenase which oxidizes α -glycerophosphate to triose phosphate anaerobically with methylene blue as a hydrogen acceptor (Table 6) or aerobically with methylene blue as a hydrogen carrier (Table 7). Prolonged dialysis has

not revealed the presence of a dissociable coenzyme for this reaction. The extracts possess an enolase (inhibited by fluoride) which converts D-3-phosphoglycerate to phosphoenolpyruvate (Table 10) and an enzyme which splits phosphoenolpyruvic acid into pyruvic acid and inorganic phosphate (Table 9).

Since triose phosphate dehydrogenase has been found in extracts of *Myco. smegmatis* (Faine, Whiteside & Edson, 1951), the following scheme is put forward to represent the catabolism of glycerol in that organism and probably in other mycobacteria:

Both pyruvate and acetate are readily oxidized by intact cells (Table 1). Thus the catabolism of glycerol converges on the chief oxidative machinery of the mycobacteria, the initial phosphorylation being analogous to that which occurs in suspensions of homogenized mammalian tissues (Kalckar, 1939; Lindberg, 1951). Since glyceraldehyde and dihydroxyacetone are also oxidized (Table 1) an obligatory phosphorylation similar to that occurring in kidney (Lindberg, 1951) may initiate the breakdown.

In the above scheme diversion to synthetic processes (assimilation) is likely to take place at the triose phosphate and 'acetate' stages.

SUMMARY

1. Glycerol is oxidized and assimilated by washed suspensions of *Mycobacterium smegmatis* and *Mycobacterium butyricum*.

2. The oxidation is completely inhibited by cyanide and partially inhibited by selective agents such as arsenite, iodoacetate and fluoride. Pyruvate accumulates extracellularly in the presence of arsenite only.

3. Study of dialysed extracts has shown that the primary step in the degradation of glycerol is phosphorylation with adenosine triphosphate. Glycerophosphate is subsequently oxidized to triose phosphate and broken down to pyruvate by reactions of the Embden-Meyerhof scheme.

The author is very grateful to Dr N. L. Edson for his advice and continued interest in this investigation.

REFERENCES

- Baudran, G. (1910). *C.R. Acad. Sci., Paris*, **150**, 1200.
 Bloch, H. (1944). *Schweiz. Z. Path.* **7**, 589.
 Bloch, H., Matter, E. & Suter, E. (1947). *Amer. Rev. Tuberc.* **55**, 540.
 Booth, V. H. & Green, D. E. (1938). *Biochem. J.* **32**, 855.
 Cavallini, D., Frontali, N. & Toschi, G. (1949). *Nature, Lond.*, **163**, 568.
 Dingle, J. H. & Weinzirl, J. (1932). *J. Bact.* **23**, 281.
 Edson, N. L. (1951). *Bact. Rev.* **15**, 147.
 Edson, N. L. & Hunter, G. J. E. (1943). *Biochem. J.* **37**, 563.
 Edson, N. L. & Hunter, G. J. E. (1947). *Biochem. J.* **41**, 139.
 Faine, S., Whiteside, M. & Edson, N. L. (1951). *Proc. Univ. Otago med. Sch.* **29**, 18.
 Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
 Fitzgerald, R. J. & Bernheim, F. (1948). *J. Bact.* **55**, 677.
 Friedemann, T. E. & Haugen, G. E. (1943). *J. biol. Chem.* **147**, 415.
 Friedemann, T. E. & Seibert, F. B. (1939). *Proc. Soc. exp. Biol., N.Y.*, **42**, 41.
 Hammerschlag, A. (1891). *Z. klin. Med.* **12**, 9.
 Hunter, G. J. E. (1952). *Proc. Univ. Otago med. Sch.* **30**, 10.
 Kalckar, H. (1939). *Enzymologia*, **6**, 209.
 Krebs, H. A. (1935). *Biochem. J.* **29**, 1620.
 LePage, G. A. (1947). *J. biol. Chem.* **168**, 623.
 Lindberg, O. (1951). *Biochim. biophys. Acta*, **7**, 349.
 Lindsay, M., O'Donnell, T. V. & Edson, N. L. (1950). *Biochem. J.* **46**, 248.
 Lohmann, K. & Meyerhof, O. (1934). *Biochem. Z.* **273**, 60.
 Merrill, M. H. (1930). *J. Bact.* **20**, 235.
 Merrill, M. H. (1931a). *J. Bact.* **21**, 361.
 Merrill, M. H. (1931b). *J. Bact.* **21**, 375.
 Nakamura, T. (1938). *Tôhoku J. exp. Med.* **34**, 231.
 Nocard, E. & Roux, E. (1887). *Ann. Inst. Pasteur*, **1**, 19.
 Sibley, J. A. & Lehninger, A. L. (1949). *J. biol. Chem.* **177**, 859.
 Voris, L., Ellis, G. & Maynard, L. A. (1940). *J. biol. Chem.* **133**, 491.
 Warburg, O. & Yabusoe, M. (1924). *Biochem. Z.* **146**, 380.
 Wedum, A. G. (1936). *J. Bact.* **32**, 599.

The Action of Trypsin on Polylysine

By S. G. WALEY AND J. WATSON

Courtaulds Ltd., Research Laboratories, Maidenhead, Berks

(Received 4 March 1953)

The way in which enzymes act is an outstanding problem in biochemistry, and of all substrates proteins are of predominating interest. Many investigations into the action of proteolytic enzymes have been carried out (Tiselius & Eriksson-Quensel, 1939; Linderstrøm-Lang & Jacobsen, 1941; Haugaard & Roberts, 1942; Winnick, 1944; Christensen, 1949; Desnuelle, Røvery & Bonjour, 1950; Røvery, Desnuelle & Bonjour, 1950; Smith, 1951), but it is a formidable task to elucidate the nature of all the products formed and even more difficult to measure their individual rates of formation. The kinetics of the hydrolysis of simple substrates catalysed by crystalline enzymes (particularly those from the pancreas, Neurath & Schwert, 1950) have been much studied and are relatively

well understood. In general, such a substrate contains only one bond capable of being hydrolysed in the presence of the enzyme. Thus the problem of the relative rates of hydrolysis of several susceptible bonds in one substrate molecule does not arise. Synthetic polypeptides containing only one amino acid residue may have many susceptible bonds, but their relatively simple constitution restricts the number of products formed. Even here, the question whether all the susceptible bonds are split at the same rate, or some much faster than others, is not readily answered. This paper describes the attempt to answer this question in one case. The customary methods of following the rate of enzymic hydrolyses by measuring the overall rate of peptide-bond splitting are not directly helpful and the newer