3. The base of the pole has a carbohydrate makeup exactly analogous to that of its lower contiguous section, the stem. Moving upwards along the pole, the picture changes sequentially: first fructosans disappear, then their intermediates, leaving only fructose, glucose, and sucrose. On the other hand, in the top-most section of the pole, the seat of flower-head and bulbel formation, the fructosans and their intermediates reappear; the entire picture is in fact similar to that of the carbohydrates in the stem.

We are grateful to Dr V. Subrahmanyam, Director of this Institute, for his helpful interest. Our thanks are due to

Mr M. N. Satyanarayana for carrying out some of the determinations.

REFERENCES

- Bacon, J. S. D. & Bell, D. J. (1948). *Biochem. J.* **42**, 397.
 Bhatia, I. S., Satyanarayana, M. N. & Srinivasan, M. (1953*a*). *Curr. Sci.* **22**, 16.
 Bhatia, I. S., Satyanarayana, M. N. & Srinivasan, M. (1953*b*). *Proc. Soc. biol. Chem. India*, **11**, 26.
 Bhatia, I. S. & Srinivasan, M. (1953). *Nature, Lond.* Submitted for publication.
 Sobotka, H. & Reiner, M. (1930). *Biochem. J.* **24**, 394.
 Srinivasan, M. & Bhatia, I. S. (1953). *Biochem. J.* **55**, 286.

Sedimentation and Diffusion of Human Albumins

3. VARIOUS NORMAL HUMAN ALBUMIN PREPARATIONS

By P. A. CHARLWOOD*

Courtauld Institute of Biochemistry, Middlesex Hospital Medical School, London, W. 1

(Received 22 July 1953)

In the course of a comparative study of normal and pathological albumins (Charlwood, 1952*a, b*), sedimentation and diffusion studies were made on albumins isolated by electrophoresis from normal human sera. The ultracentrifuge results were lower than might have been anticipated, judging from the publications of previous workers. After examination of the factors involved, it was concluded that the discrepancy lay not in the material, but in an error in determinations of temperature in the earlier work. This conclusion is in agreement with recent measurements on other proteins (Cecil & Ogston, 1948; Kegeles & Gutter, 1951; Miller & Golder, 1952; Taylor, 1952).

The diffusion constants, however, having a mean value of 6.32 units at about 0.2% albumin concentration, were higher than most of the few published figures. Other workers usually gave little detail of their experimental methods, and sometimes omitted to specify concentrations. It seemed that some of the differences might be due to the different methods of isolation of the albumins. To investigate this question, a study has now been made of albumins prepared by three different processes, and the effects of variations in buffer composition have been measured simultaneously.

Slight differences of diffusion constant existed among the normal albumins previously examined (Charlwood, 1952*a*). It is known that, under

various conditions, electrophoresis causes a splitting of the albumin peak (Luetscher, 1939; Hoch & Morris, 1945). This occurs during long separations in veronal buffer. It is clear from Fig. 1, which shows the position of an albumin boundary in the cell immediately before sliding the section, that the relative proportions of the two major albumin components will not only differ from the proportions originally present (quite apart from concentration adjustments caused by the fulfilment of electrochemical conditions, as described by Hoch (1948)), but will depend on the position of the boundary relative to the cell end-plate. This position is determined by the conditions and length of run. The relative proportion of faster component is increased by this effect, which could only be overcome by

Fig. 1. Electrophoretic pattern of albumin boundary given by human serum. Ascending (anode) limb. Migration is from left to right. Veronal buffer, pH 8.6, $I=0.1$. Exposure after 22 hr. 20 min. at 4.7 v/cm.

* Present address: Division of Applied Biology, National Research Council Laboratories, Ottawa, Ontario, Canada.

extending the run until the whole boundary was located above the end plate. If the albumin components differed significantly as regards diffusion constant, the presence of different relative amounts would lead to variations from one sample to another, which could account not only for the differences between normal albumins (Charlwood, 1952*a*), but also for the results obtained in some pathological cases (Charlwood, 1952*b*). Some experiments have, therefore, been designed to decide this issue, by actual separation and examination of the faster albumin component observed by Luetscher (1939).

Unfortunately, it has not been possible to obtain any crystalline material for inclusion in these studies. This would have permitted a direct comparison with molecular weights as determined by X-rays (Low, 1952). The latter method gives a molecular weight for crystalline human albumin which is very close to that found for crystalline bovine albumin in recent sedimentation and diffusion measurements (Creeth, 1952).

EXPERIMENTAL

Materials used. Two human albumin preparations, made by the methods of Cohn *et al.* (1944, 1946), were obtained from Harvard. The first, henceforth designated A, contained 0.3% α -globulin. It had been kept at 60° for 10 hr. (in the presence of a small amount of sodium acetyltryptophanate) to sterilize it for injection purposes. The second sample, which will be denoted by B, contained 6% α -globulin. It had not been heated during processing. It was calculated that the small amount of α -globulin in A could be ignored in the sedimentation and diffusion measurements, but that in B was removed by electrophoresis in veronal buffer. The third main albumin preparation used (C) was isolated by electrophoresis (in veronal buffer) of pooled sera from ten normal persons.

All three albumins were studied by sedimentation and diffusion at five different concentrations in the range 0.17–0.8 g./100 ml. The fast component A_1 (see below) was examined at three concentrations in this range.

Electrophoretic separations. These were nearly all carried out in veronal buffer (pH 8.6, $I=0.1$), as described by Charlwood (1952*a*). Owing to the large number of runs required in this work, an automatic control was devised (Warren & Charlwood, 1953) to operate the electrolytic compensator (Johnson & Shooter, 1949). The control, based on a photoelectric principle, switches the compensator on and off in such a manner as to keep the albumin boundary virtually stationary for long periods. This meant that, for the long runs formerly taking up to 10 hr. or more with periodic manual control, automatically controlled runs of 15–20 hr. could be substituted. Whereas in the former the maximum permissible current was used, a lower current (15 mA) could be employed in the longer runs, giving the same separation with less risk of thermal convection. These long runs were carried out overnight, the day being free for recovering the albumin, cleaning and reassembly of the cell and starting a fresh run.

The faster component (A_1) of the albumin A was separated by electrophoresis of a solution containing about 2 g. protein/100 ml., the buffer being sodium acetate (pH 4.0, $I=0.02$) (Luetscher, 1939). Under these conditions a current of only 3 ma was used, because of the lower specific conductance of the buffer.

Sedimentation and diffusion. These were measured as described by Creeth (1952) and Charlwood (1952*a, b*). At concentrations below about 0.4 g./100 ml., two pairs of diffusion runs were carried out on any particular sample, whilst at higher concentrations a single pair sufficed.

All three albumins, A, B and C, were studied in the two following solutions (a) NaCl ($I=0.2$) + sodium acetate buffer ($I=0.02$, pH 4.6) and (b) NaCl ($I=0.2$) + sodium phosphate buffer ($I=0.02$, pH 8.0). In addition, A and C were examined in unbuffered NaCl ($I=0.2$). Measurements on C at the lowest concentration (in NaCl, $I=0.2$) were omitted, the appropriate values obtained in NaCl, $I=0.05$ (Charlwood, 1952*a*), being used in the subsequent analysis. A duplicate set of diffusion measurements was made (in buffer solution (a)) on albumin A which had been dialysed to equilibrium against the veronal buffer used for electrophoresis ($I=0.1$, pH 8.6) and subsequently redialysed against the acetate buffer.

The component A_1 , isolated in acetate buffer, was redialysed against acetate buffer of pH 4.5 ($I=0.05$) before sedimentation and diffusion.

In addition to the above experiments on A and B, sedimentation was observed at a relatively high concentration, but at a low temperature (5–6°). Albumin A was run in sodium phosphate buffer ($I=0.2$, pH 8.0, protein 2.5 g./100 ml.) for nearly 5 hr., and in sodium acetate buffer ($I=0.2$, pH 4.5, protein 2.0 g./100 ml.) for about 4 hr. Albumin B was run in the same acetate buffer at 1.5 g./100 ml. for 4 hr. The purpose of these ultracentrifuging experiments is discussed below.

RESULTS

The results of the main series of experiments are shown in Figs. 2–4 and 6–8. The abscissae denote relative concentrations, expressed as wavelengths of retardation (v) observed in the diffusimeter; conversion into absolute values can be made from the known length of the cell (1.09 cm.) and the specific refractive increment (about 0.00190) of human albumin (Perlmann & Longsworth, 1948). Albumin at 1 g./100 ml. corresponds with $v \sim 38$. Sedimentation constants are all given in Svedberg units and diffusion constants in units of 10^{-7} cm.² sec.⁻¹.

Sedimentation measurements

It has been customary to refer a value of the sedimentation constant to the initial concentration prevailing in the solution, in spite of the known dilution effect due to the sectorial shape of cell (Svedberg & Pedersen, 1940). Values were referred to the mean concentration in the time interval between the first and last exposures in the experiments of Kegeles & Gutter (1951). More recently, Golder (1953) has discussed the general effect of

Fig. 2. Sedimentation of albumin A. Points correspond to the following conditions: $\circ$, NaCl ($I=0.02$); $\bullet$, NaCl ($I=0.02$) + sodium phosphate buffer ($I=0.02$, pH 8.0); $\odot$, NaCl ($I=0.02$) + sodium acetate buffer ($I=0.02$, pH 4.6).

Fig. 3. Sedimentation of albumin B. Points correspond to the following conditions: $\bullet$, NaCl ($I=0.02$) + sodium phosphate buffer ($I=0.02$, pH 8.0); $\odot$, NaCl ($I=0.02$) + sodium acetate buffer ($I=0.02$, pH 4.6). Upper regression line refers to acetate, lower to phosphate solutions.

concentration in relation to measured values of sedimentation constants. In earlier work (Charlwood, 1952*a, b*), consideration of these effects led to the conclusion that it was hardly worth while to depart from the conventional procedure, as all runs were made under closely similar conditions and the dilution effect was not very important. The same holds here, although higher protein concentrations have been used. The sectorial dilution effect would merely require multiplication of the concentrations by a factor virtually the same for all solutions, which would not alter extrapolated $S_{20,w}^0$ values. As S changes only slowly with concentration there is added justification for retaining the usual method.

Fig. 4. Sedimentation of albumin C. Points correspond to the following conditions, except as stated in Experimental section: $\circ$, NaCl ($I=0.02$); $\bullet$, NaCl ($I=0.02$) + sodium phosphate buffer ($I=0.02$, pH 8.0); $\odot$, NaCl ($I=0.02$) + sodium acetate buffer ($I=0.02$, pH 4.6). Upper regression line refers to unbuffered and acetate, lower line to phosphate solutions.

In Figs. 2-4 the results of measurements on A, B and C are summarized. All are corrected to water at 20° , assuming a partial specific volume, $\bar{V}=0.733$ (cf. Charlwood, 1952*b*). The point shown at the lowest concentration for C in unbuffered solution is the mean of the six determinations in sodium chloride ($I=0.05$) given by Charlwood (1952*a*), but all the individual values were used in the statistical analysis. The error variance was estimated from all measurements taken together.

The sedimentation of A (Fig. 2) showed highly significant concentration dependence ($P<0.001$), but no detectable differences in the effects of the various supporting media. The line shown is the regression line for the whole set of points.

Since, at finite concentrations, B sedimented more slowly in phosphate buffer than in acetate, two regression lines are given in Fig. 3. Similarly, in the case of C, lower results were obtained in phosphate solution than in acetate, or unbuffered sodium chloride. Thus, in Fig. 4, two regression lines are shown.

In no case do the values of $S_{20,w}$ for a particular preparation, obtained in different solutions, give significantly different estimates when extrapolated to zero concentration. Consequently, for each albumin a weighted mean value of $S_{20,w}^0$ has been calculated (Table 1), together with the estimated error. Further calculation has shown that albumin B possesses a significantly lower $S_{20,w}^0$ than A or C, which cannot be considered as differing from each other.

In Fig. 5 are shown patterns given by albumin A in the low-temperature experiments. There is no sign of any hump developing on the peak, or any tendency for it to split. A similar result was

Table 1. *Molecular characteristics of three different albumin preparations*

Albumin	$S_{20,w}^0$	$D_{20,w}^0$	Molecular weight	
			$\bar{V}=0.733$	$\bar{V}=0.75$
A	4.31 ± 0.04	6.07 ± 0.07	64 500	68 900
B	4.20 ± 0.06	6.14 ± 0.04	62 100	66 300
C	4.33 ± 0.04	6.43 ± 0.06	61 000	65 100

The ranges shown are 95% fiducial limits.

Fig. 5. Ultracentrifuge patterns of albumin A at about 5°. Sedimentation is from right to left. Left-hand photograph at 2.5 g. protein/100 ml. after 285 min. at top speed (phosphate buffer, $I=0.2$, pH 8.0). Right-hand photograph at 2.0 g. protein/100 ml. after 124 min. at top speed (acetate buffer, $I=0.2$, pH 4.5).

Table 2. *Sedimentation and diffusion of albumin A₁*

(Measurements in 0.05 I sodium acetate buffer, pH 4.5.)

Fig. 6. Diffusion of albumin A. Points correspond to the following conditions: O, NaCl ($I=0.2$); ●, NaCl ($I=0.2$) + sodium phosphate buffer ($I=0.02$, pH 8.0); ⊙, NaCl ($I=0.2$) + sodium acetate buffer ($I=0.02$, pH 4.6). Upper regression line refers to unbuffered and phosphate, lower line to acetate solutions.

Fig. 7. Diffusion of albumin B. Points correspond to the following conditions: ●, NaCl ($I=0.2$) + sodium phosphate buffer ($I=0.02$, pH 8.0); ⊙, NaCl ($I=0.2$) + sodium acetate buffer ($I=0.02$, pH 4.6).

Fig. 8. Diffusion of albumin C. Points correspond to the following conditions, except as stated in Experimental section: O, NaCl ($I=0.2$); ●, NaCl ($I=0.2$) + sodium phosphate buffer ($I=0.02$, pH 8.0); ⊙, NaCl ($I=0.2$) + sodium acetate buffer ($I=0.02$, pH 4.6). Upper regression line refers to unbuffered and phosphate, lower line to acetate solutions.

Table 3. *Diffusion of albumin A after contact with veronal buffer*

(Measurements in 0.2 I sodium chloride + 0.02 I sodium acetate buffer, pH 4.6.)

$D_{20,w}$	v				
	6.05	12.28	18.22	24.04	30.18
	6.08	5.94	6.05	5.80	5.89

obtained with albumin B. It is interesting to compare the sedimentation constants measured under these conditions with values extrapolated to the same concentrations calculated from regression lines in Figs. 2 and 3. For albumin A, in the acetate buffer, the value of $S_{20,w}^0$ was found to be 3.74 , the extrapolated value and 95% fiducial limits being 3.73 ± 0.12 . Agreement is considerably better than might be reasonably expected. For albumin B, $S_{20,w}^0 = 3.90$, the corresponding extrapolated figures being 4.09 ± 0.11 , based on the regression line for acetate buffer (Fig. 3). Considering the uncertainties of temperature measurement in these experiments (the final values being about $4-5^\circ$ higher than the initial, compared with the rise of $1-1.5^\circ$ associated with normal, shorter runs) there seems little justification for attempting to attach any particular significance to the discrepancy of about 2% in this case.

The sedimentation results on A_1 are shown in Table 2. They may be compared with those for the whole albumin illustrated in Fig. 2 (see also Discussion).

Diffusion measurements

Diffusion results, corrected to water at 20° , are shown in Figs. 6-8. In general, each point represents the mean of one pair of determinations (above about 0.4 g./100 ml.) or two pairs (below 0.4 g./100 ml.). An exception is again the point corresponding to the lowest concentration in sodium chloride (albumin C), which was derived by taking the mean of the values for the six albumin preparations examined in $0.05 \text{ } I$ sodium chloride (Charlwood, 1952a). Individual values were used in calculating means, regressions, and limits of error.

Albumins A and C (Figs. 6, 8) gave lower values of D in acetate buffer than in the other solutions. Of the two regression lines shown in each figure, one corresponds to acetate buffer, the other to the remaining measurements. As albumin B (Fig. 7) did not show any significant dependence on concentration or buffer, the best estimate of $D_{20,w}^0$ in this case was calculated as the mean of all results.

Only in one case (C) do the extrapolated $D_{20,w}^0$ values for a particular albumin show any sign of appreciable difference. The difference is exactly on the probability limit ($P \sim 0.05$) and, as discussed below, has been ignored in calculating the weighted mean shown in Table 1. From the figures in Table 1 it appears that albumin C has a $D_{20,w}^0$ highly significantly greater than A or B, which do not differ between themselves.

The diffusion results obtained with A_1 and those given by A after preliminary treatment with veronal buffer, are shown in Tables 2 and 3 respectively. They may be compared with those for albumin A in Fig. 6.

DISCUSSION

In the first paragraph of this paper reference has been made to the accuracy of sedimentation measurements. According to Waugh & Yphantis (1952) the temperature of the rotor in the Spinco ultracentrifuge drops about 1° on accelerating to maximum speed, and rises by a corresponding amount on deceleration. This would mean that sedimentation constants calculated (as in this series of papers) from the mean of initial and final rotor temperatures would be about 2.5% too low. Shulman (1953) found his values in the Spinco machine to be generally 2-3% lower than those obtained with the oil turbine run under the standard conditions of Cecil & Ogston (1948). However, melting-point experiments by Shulman (1953) did not appear to support the findings of Waugh & Yphantis. It was pointed out by Shulman (1953) that this might well be due to the length of the runs used for the melting-point experiments. In long runs the precise manner of variation of rotor temperature during the period at top speed may well be such as to disguise the temperature alterations which occur on acceleration or deceleration. The experiments in the present paper being of relatively short duration, the correction would be expected to apply. It must be mentioned that Baldwin & Ogston (quoted by Shulman, 1953) have obtained results with bovine serum albumin in the oil turbine ultracentrifuge which agree with measurements made in the Spinco, rather than with the slightly higher values Shulman found in his oil turbine experiments. Since the whole matter could still bear further investigation, no correction has been made here. All measurements in the present series of papers thereby remain directly comparable; and, if at a later date it is decided that the correction ought, without any doubt, to be made, it will be simple to apply the appropriate factor to the whole set of sedimentation values.

The regression of sedimentation and diffusion values on protein concentration would have been defined more accurately if determinations could have been made over a wider range of protein concentration. It would not have been easy to obtain sufficient albumin by electrophoretic separation to work at much higher concentrations. Moreover, a number of alterations would have had to be made in the construction of the diffusimeter, to extend the present upper limit of protein concentration of about 1 g./100 ml. which did not seem to be justified.

The dependence of diffusion constant on the source of albumin, protein concentration, and buffer conditions suggests that the variation shown in the results of other workers (quoted by Charlwood 1952a) may be ascribable to these factors. It has

been shown, however, that results at different concentrations and in different electrolyte solutions all lead to the same value of $D_{20,w}^0$ for any particular sample. The possible exception to this statement is albumin C in acetate buffer, which tends to give a higher value of $D_{20,w}^0$ (Fig. 8) than under other conditions. It must, however, be pointed out that this sample was separated electrophoretically from pooled sera from a group of persons quite distinct from those who provided the source of the first batch of material (on which measurements in phosphate and unbuffered solutions were made). Since small individual variations exist (Charlwood, 1952*a*), it would be unwise to assume that the acetate buffer is responsible for the suggested effect.

It is interesting to note how closely a previous assumption (Charlwood, 1952*a, b*), that sedimentation and diffusion measurements on albumin at $\sim 0.2\%$ concentration may be used in the calculation of molecular weights without appreciable error, is borne out by the present experiments. The $S_{20,w}^0/D_{20,w}^0$ ratio for albumin C differs by only 0.8% from the $S_{20,w}/D_{20,w}$ ratio found at low concentration (Charlwood, 1952*a*).

In experiments carried out by Dr J. M. Creeth for the Department of Biochemistry, Oxford (Tilley, 1953), it was observed that β -lactoglobulin, although giving a single peak in the ultracentrifuge at ordinary room temperatures, shows a distinct splitting into two components of considerably different molecular weight when sedimenting at lower temperatures ($\sim 5^\circ$). This has been ascribed to association of molecules (Tilley, 1953). Albumins A and B (Fig. 5) showed no sign of this behaviour, but gave single peaks. As the sedimentation constants calculated in these experiments agreed with those observed at room temperature (see above), it can be assumed that these albumins are not associated at all under these conditions. Sedimentation experiments are normally carried out at room temperature and the results are corrected to 20° . If there had been any evidence of temperature-dependent association, the validity of this correction would have had to be examined, because runs have been made at temperatures as low as 15° , and as high as 26° .

Molecular weights. The molecular weights shown in the fourth column of Table 1 were calculated on the basis of a partial specific volume, $\bar{V}$, of 0.733 (see Charlwood, 1952*a*). If a value of 0.75 were used they would all be increased by the same factor (1.07) as in the fifth column. In any case they decrease relatively from A to C, but the differences indicated may be underestimations or overestimations on account of the possible errors indicated by the fiducial limits.

The results in Table 1 might be taken to indicate

that treatment with veronal buffer causes irreversible changes in size and shape of albumin molecules, leading to differences in observed molecular weights and frictional ratios. The evidence presented in Table 3 is, so far as it goes, against this hypothesis, as the duplicate measurements of the diffusion behaviour of A after treatment with veronal buffer do not differ, within experimental error, from those in Fig. 6 obtained without such treatment.

An alternative interpretation of the figures in Table 1 is that, at certain stages in the ethanol fractionation, there occurs an elimination of some species of lower molecular weight and different frictional ratio which normally migrate in the albumin boundary. In this connexion, as in earlier work (Charlwood, 1952*a, b*), it would be interesting to see whether or not the molecular weights of the faster and slower albumin components are distinguishable from each other. There is no *a priori* reason for favouring either alternative, as it is well-known that proteins of widely separated molecular weights may possess virtually identical electrophoretic mobilities (e.g. the β_1 -globulins studied by Oncley, Scatchard & Brown, 1947); and, conversely, proteins of the same molecular weight may exhibit very different electrophoretic mobilities (e.g. the α_2 - and γ -globulins of Oncley *et al.* 1947).

The figures in Table 2 indicate that the faster albumin component isolated in acetate buffer is of almost, if not exactly, the same molecular weight as the whole albumin. Unfortunately, there is no evidence to show whether this faster component corresponds with that which separates in veronal buffer. It cannot, therefore, be stated categorically that the idea discussed above (in relation to Fig. 1) has been proved false. The accuracy of the results in Table 2 is not so great as to preclude a value for the diffusion constant which could account for the small differences observed among normal human albumins at a low concentration (Charlwood, 1952*a*), but is not very likely to explain the present results, or those reported in some pathological cases (Charlwood, 1952*b*).

From the evidence given here it is not possible to conclude that each electrophoretic component of the albumin is comprised of groups of molecules differing in both molecular weight and solubility in ethanol. But even if this likely hypothesis were true, it would not contribute towards a solution of the problem of the pathological albumins. Whether the abnormal molecular weights found for albumins in nephrosis are due to unusual relative proportions of normal molecules, or to the presence of entirely different molecules, must remain an open question, soluble only by finer methods of analysis.

SUMMARY

1. Sedimentation and diffusion measurements have been made in the concentration range 0.17–0.8 g./100 ml. on human serum albumins obtained by ethanol fractionation and by electrophoretic separation.

2. The effect of variation in composition of the buffer has been examined with the result that, for a given preparation, sedimentation or diffusion may be affected at finite concentrations, but values extrapolated to zero concentration do not differ significantly.

3. Values of $S_{20,w}^0$ and $D_{20,w}^0$ vary slightly from one preparation to another, indicating differences in average molecular weight. Evidence is presented which suggests that this is due to the separation of groups of molecules of different molecular characteristics rather than to irreversible changes produced by buffer conditions.

4. The faster of the two components which appear on electrophoresis of albumin at pH 4.0 has been isolated. Its sedimentation and diffusion properties were indistinguishable from those of whole albumin.

The author wishes to thank Prof. E. C. Dodds, F.R.S., for his interest and help. He is very grateful to the Department of Physical Chemistry, Harvard Medical School, for gifts of two of the albumin preparations, and to Dr D. N. Baron for providing the normal sera. This work was carried out whilst the author was an Imperial Chemical Industries Research Fellow in the University of London.

REFERENCES

- Cecil, R. & Ogston, A. G. (1948). *Biochem. J.* **43**, 592.
 Charlwood, P. A. (1952a). *Biochem. J.* **51**, 113.
 Charlwood, P. A. (1952b). *Biochem. J.* **52**, 279.
 Cohn, E. J., Oncley, J. L., Strong, L. E., Hughes, W. L. jun. & Armstrong, S. H. jun. (1944). *J. clin. Invest.* **23**, 417.
 Cohn, E. J., Strong, L. E., Hughes, W. L. jun., Mulford, D. J. Ashworth, J. N., Melin, M. & Taylor, H. L. (1946). *J. Amer. chem. Soc.* **68**, 459.
 Creeth, J. M. (1952). *Biochem. J.* **51**, 10.
 Golder, R. H. (1953). *J. Amer. chem. Soc.* **75**, 1739.
 Hoch, H. (1948). *Biochem. J.* **42**, 181.
 Hoch, H. & Morris, C. J. O. R. (1945). *Nature, Lond.*, **156**, 234.
 Johnson, P. & Shooter, E. M. (1949). *Science*, **109**, 39.
 Kegeles, G. & Gutter, F. J. (1951). *J. Amer. chem. Soc.* **73**, 3770.
 Luetscher, J. A. jun. (1939). *J. Amer. chem. Soc.* **61**, 2888.
 Low, B. W. (1952). *J. Amer. chem. Soc.* **74**, 4830.
 Miller, G. L. & Golder, R. H. (1952). *Arch. Biochem. Biophys.* **36**, 249.
 Oncley, J. L., Scatchard, G. & Brown, A. (1947). *J. phys. Chem.* **51**, 184.
 Perlmann, G. E. & Longworth, L. G. (1948). *J. Amer. chem. Soc.* **70**, 2719.
 Shulman, S. (1953). *Arch. Biochem. Biophys.* **44**, 230.
 Svedberg, T. & Pedersen, K. O. (1940). *The Ultracentrifuge*. Oxford University Press.
 Taylor, J. F. (1952). *Arch. Biochem. Biophys.* **36**, 357.
 Tilley, J. M. A. (1953). Thesis, University of Oxford.
 Warren, R. L. & Charlwood, P. A. (1953). *Nature, Lond.*, **171**, 353.
 Waugh, D. F. & Yphantis, D. A. (1952). *Rev. sci. Instrum.* **23**, 609.

The Protective Action of Neutral Solutes against Haemolysis by Freezing and Thawing

By J. E. LOVELOCK

National Institute for Medical Research, Mill Hill, London, N.W. 7

(Received 22 July 1953)

It has been known for many years that sugar solutions have a slight effect in protecting living cells against damage during freezing and thawing (Smith, 1953). Polge, Smith & Parkes (1949) discovered that spermatozoa could be frozen and thawed without loss of motility if glycerol or some similar polyhydric alcohol was included in their suspending medium. Investigation of the protective action of glycerol was facilitated by the observation that red blood cells also were protected by glycerol against the adverse effects of freezing and thawing (Smith, 1950). The nature of the destructive processes which take place when red blood cells are frozen and

thawed, and of the protection given by glycerol was then described (Lovelock, 1953a, b). Briefly, damage on freezing is a consequence of the increase in concentration of electrolytes within and without the cell. In the presence of glycerol the electrolyte concentration at temperatures below the freezing-point is greatly diminished, by an amount sufficient to explain the protection afforded by glycerol.

The effect of glycerol in lowering the concentration of electrolytes in equilibrium with ice at temperatures below 0° is a simple consequence of the colligative properties of solutions. It follows that the protective action of glycerol is unlikely to be