Table 4. *Content of ^{35}S of hair from fed animals*

Animal no.	...	...	11	12	15
Units fed/day	...	...	15	15	15
Total [^{35}S]methionine fed in 4 weeks ($\mu\text{g.}$)			441	441	441
Weeks	[^{35}S]Methionine equiv. of weekly hair growth ($\mu\text{g./g.}$)				
1	0.141	0.282	0.706		
2	0.886	0.581	1.85		
3	1.48	2.13	2.25		
4	Died	3.04	2.85		
5	—	2.30	2.42		
6	—	1.85	1.29		
7	—	1.052	0.820		
8	—	0.958	0.760		
9	—	0.460	0.346		
10	—	0.394	0.425		
11	—	0.389	0.292		
12	—	0.302	0.220		
13	—	0.235	0.185		
14	—	0.263	0.111		
15	—	0.172	0.075		
16	—	0.150	0.027		
17	—	Died	0.007		
Total recovery as [^{35}S]methionine in hair ($\mu\text{g./animal}$)	2.00	11.64	11.71		
Percentage recovery	0.5	2.6	2.7		

the ^{35}S was distributed throughout the system in the same manner as if it had been fed or injected (cf. Tarver & Schmidt, 1939). It seems likely that once the methionine reaches the blood stream the demethylation proceeds in the usual manner irrespective of the method of introduction. This does not preclude the possibility that a small proportion of the topically applied methionine might be converted into cystine locally in the root cells, but to demonstrate such a mechanism a more detailed investigation, possibly involving tissue-culture techniques, would be necessary.

The Effect of Ammonia on the Oxidation of Glucose by *Zygorhyncus moelleri*

By V. MOSES

Department of Botany, University College, London

(Received 14 July 1953)

During an investigation of the respiration of filamentous fungi it was observed that when glucose was given to carbohydrate-starved suspensions of several species 2–3 hr. elapsed before the rate of oxygen consumption became constant. This behaviour is markedly different from glucose oxidation by many other micro-organisms under similar conditions, e.g. yeast and the green alga *Chlorella*,

in which the addition of glucose to starved cells results in a constant respiratory rate being attained within 10–15 min.

Induction periods have been noticed by other workers in the oxidation of acetate by both yeast and by bacteria. Winzler (1940), investigating acetate oxidation by suspensions of baker's yeast, found that preparations starved for 2 hr. required

SUMMARY

1. [^{35}S]Methionine was applied to the skin of the guinea pig; this resulted in incorporation of ^{35}S in the protein of the growing hair, largely as cystine. This was shown by the distribution of the radioactivity in a paper chromatogram of the hair hydrolysate. The ^{35}S content of the hair was similar whether the specimen was from the area of application of methionine or from a remote area. About 1 % of the applied ^{35}S was incorporated in the hair.

2. The amount of ^{35}S absorbed was proportional to the amount rubbed into the skin over the concentration range investigated.

3. If the same amount of [^{35}S]methionine was fed to the guinea pig, about 2.5 times as much ^{35}S was incorporated into the hair.

4. Injection of [^{35}S]methionine intramuscularly was relatively more efficient, as much as 5 % of ^{35}S being incorporated in hair protein; larger doses were utilized to a lesser degree.

The author wishes to thank Dr R. H. Marriott for his valuable advice and suggestions; Mr D. M. Brown and Mr J. A. Dimsdale for their care of the animals and practical help; Mr T. L. Hardy for his advice and assistance with the paper chromatography; and Mr E. W. Powell for much useful discussion in the planning period.

REFERENCES

- Block, R. J. & Bolling, D. (1951). *The Amino Acid Composition of Proteins and Foods*, 2nd ed. Springfield: Thomas.
- Comar, C. L., Hansard, S. L., Hood, S. L., Plumlee, M. P. & Barrentine, B. F. (1951). *Nucleonics*, **8**, 19, fig. 5.
- Du Vigneaud, V., Kilmer, G. W., Rachele, J. R. & Cohn, M. (1944). *J. biol. Chem.* **155**, 645.
- Henriques, F. C., Kistiakowsky, G. B., Margnetti, C. & Schneider, W. G. (1946). *Ind. Engng Chem. (Anal.)*, **18**, 349.
- Tarver, H. & Schmidt, C. L. A. (1939). *J. biol. Chem.* **130**, 67.
- Tarver, H. & Schmidt, C. L. A. (1942). *J. biol. Chem.* **146**, 69.
- Williamson, M. B. & Fromm, H. J. (1953). *Proc. Soc. exp. Biol., N.Y.*, **83**, 329.

7 min. for the oxidation rate to become constant, whereas when starved for 70 hr. the suspension required 70 min., although the final rate of oxygen consumption reached in each case was the same. Barron, Ardau & Hearon (1950) found that *Corynebacterium creatinovorans* grown on acetate and subsequently starved by suspending it in water and aerating with oxygen for 24 hr. showed an induction period on the addition of acetate, and at least 90 min. elapsed before the rate of oxygen uptake became constant. No explanation of these findings were given in either of these two instances.

Lynen (1942) also found an induction period when acetate was added to yeast which had been starved for 15–20 hr. He attributed this to the disappearance during starvation of metabolites necessary for acetate oxidation, and which had to be resynthesized before acetate could be oxidized at a maximal rate.

It was decided to use one of the fungi for an investigation of the factors involved in the appearance of the induction period. The species chosen was *Zygorhynchus moelleri*, a member of the Mucorales. No reference has been found in the literature to previous metabolic studies performed with this organism.

EXPERIMENTAL

Growth. The organism used was *Z. moelleri* (Commonwealth Mycological Institute no. 21994). Considerable difficulty was experienced at first in preparing a suspension of the organism sufficiently homogeneous and finely divided to enable replicate samples to be pipetted into Warburg micro-respirometer flasks. Various methods have been recommended for the preparation of homogeneous fungal suspensions: growth in a Warburg flask itself (Mann, 1944); inoculation of fragmented mycelium into the medium (Dorrell & Page, 1947); inoculation of the medium with a very heavy spore suspension, with shaking during growth (Darby & Goddard, 1950); and aeration of the culture with a forced airstream (Foster, 1949). A discussion of the various methods is also given by Kluver & Perquin (1933). The following technique was finally adopted: the organism was maintained on potato-glucose-agar slopes, which induced heavy sporulation. Cultures on agar 1–2 weeks old were washed with about 10 ml. of sterile distilled water, accompanied by scraping with a wire loop, and the spore suspension obtained, containing about 4×10^7 spores/ml., was filtered through glass wool in the simple apparatus shown in Fig. 1. The 10 ml. of spore suspension was then inoculated into a 100 ml. Erlenmeyer flask containing 22 ml. of the following medium: glucose, 50 g.; NH_4Cl , 2 g.; KH_2PO_4 , 5.4438 g.; N-NaOH , 4.56 ml.; MgSO_4 , 7H₂O, 0.1 g.; NaCl, 0.1 g.; Lemco (liver extract) or Marmite (yeast extract), 2 g.; trace element solution, 1 ml.; made up to 1 l. with distilled water. The trace element solution (Horowitz & Beadle, 1943) contained the following trace elements as salts (mg./l.): Boron, 0.01 (as $\text{Na}_2\text{B}_4\text{O}_7$, 10H₂O); Mo, 0.02 (as $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$, 4H₂O); Fe, 0.2 (as FeSO_4 , 7H₂O); Cu, 0.1 (as CuSO_4 , 5H₂O); Mn, 0.02 (as MnSO_4 , 4H₂O); Zn, 2.0 (as ZnSO_4 , 7H₂O). The medium was autoclaved at 15 lb./sq.in. for 15 min.; the pH after autoclaving was 5.2–5.3.

The inoculated medium was shaken for about 18 hr. in a Warburg water bath at 25° (150 oscillations/min.), by which time a good, and very finely divided growth was obtained. The hyphae were centrifuged, washed with 0.066M phosphate buffer, pH 6.8, and resuspended either in 25 ml. of the growth medium without glucose (if carbohydrate-starved cells were required) or in 25 ml. of 0.066M phosphate buffer, pH 6.8 (if cells starved of both carbohydrate and nitrogen were required). They were shaken on the Warburg apparatus for a further 24 hr. to complete the starvation process and then centrifuged, washed with buffer, and resuspended in 35 ml. of 0.1M phosphate buffer, pH 6.8. This suspension contained 1.5–2.0 mg. dry weight of cells/ml. Except where stated to the contrary, the phosphate buffers used in this work were prepared from KH_2PO_4 and KOH. Aseptic precautions were observed during the starvation period but not during the measurement of respiration. Periodic bacteriological tests performed after an experiment have never revealed more than very slight bacterial contamination, even in the presence of both added glucose and NH_4Cl . Even when preparations which had been starved in unsterilized buffer, were incubated with glucose and NH_4Cl it was found that only small numbers of bacteria developed.

Although cell suspensions produced in this way on different occasions were probably not identical with regard to density, homogeneity, or even physiological state, the differences found were apparently quite small.

Manometric methods. Oxygen uptake was measured with Warburg manometers using cups with double side bulbs, and centre wells to which was added 0.2 ml. 15% (w/v) KOH as CO_2 absorbent.

Substrates were added from the side bulb and the main compartment of the cup contained 2.0 ml. of the suspension of hyphae plus sufficient water to bring the final volume of the reaction mixture to 3.0 ml. Organic acids were added as solutions of their sodium salts, and the pH of each solution was adjusted to the required value before use. The gas phase was air save in anaerobic experiments when the manometers were flushed for 10 min. with O_2 -free N_2 .

Fig. 1. Apparatus for filtering spore suspension.

Under anaerobic conditions *Z. moelleri* produces CO_2 from glucose and at pH 6.8 a considerable amount of this gas is retained by the buffer. The amount so retained was corrected for by the procedure outlined by Umbreit, Burris & Stauffer (1949), in which α , the solubility coefficient, is replaced by α' in the calculation of the vessel constants. The value of α' was calculated on the basis of the equation:

$$\frac{\alpha'}{\alpha} = \frac{(\text{HCO}_3^-) + (\text{CO}_2)}{(\text{CO}_2)} = [\text{antilog}(\text{pH} - 6.382)] + 1,$$

where 6.382 is the first dissociation constant for carbonic acid under the conditions of the experiment (Umbreit *et al.* 1949). It should be noted that α' increases rapidly with the pH, and consequently the method is only valid if the pH remains constant throughout the experiment. Further, owing to the rapid increase in the value of α' with pH, it is inadvisable to use this method in buffers more alkaline than pH 7.0.

The respiratory quotient (R.Q.) was measured by the direct method (Dixon, 1943) and the amount of CO_2 formed was corrected for retention by the method outlined above. In four experiments the pH of the medium was measured both at the beginning and at the end of the experiment using a glass electrode and pH meter. The initial values were 6.81, 6.81, 6.80, and 6.82 respectively, and the final values 6.83, 6.79, 6.81, and 6.81 respectively. When the effect of cyanide on the rate of oxygen uptake was investigated, the KOH-KCN mixtures recommended by Robbie (1946) were used for the absorption of CO_2 .

Nitrogen analyses. Determinations of the N fractions were carried out as described by Syrett (1953*b*). Briefly, this consisted of centrifuging a sample of the cell suspension and determining the residual free extracellular NH_4^+ in the supernatant in Conway units; the NH_3 released was trapped in 0.05N- H_2SO_4 in the central compartment, and was subsequently estimated by nesslerization. The centrifuged cells were washed three times with water, and the final centrifugate, still in the centrifuge tubes, immersed in boiling water for 2 min. to kill the cells and rupture the cell membranes. The cell residues were extracted 3 times with water at 80° for 15 min., and the extracts pooled. Total N determinations were performed on the cell residues (insoluble-N fraction) and on the extract (total soluble-N fraction) by the micro-Kjeldahl technique. The NH_3 produced was determined in Conway units followed by nesslerization. The free NH_4^+ in the extract before digestion (free intracellular NH_4^+) was determined directly as described above. All the Conway distillations were allowed to proceed overnight, and all determinations were performed in duplicate.

Dry-weight determinations. The dry weights of mycelial suspensions were measured by drying samples of the suspensions overnight to constant weight at 105°.

RESULTS

Oxidation of glucose. Freshly prepared suspensions of *Z. moelleri* consumed oxygen rapidly, and addition of glucose to such suspensions only resulted in a relatively small increase in the rate of oxygen uptake about double that in the absence of glucose. If the hyphae were starved for 24 hr. by incubating them in a medium containing no carbohydrate but otherwise similar to that used for

growth, the endogenous oxygen uptake was reduced. Addition of glucose to suspensions of hyphae which had been so treated resulted in a slow but steady increase in the rate of oxygen uptake which only became constant 2-3 hr. after addition of the substrate. By contrast, suspensions of baker's yeast which had been similarly treated had a lag of some 10-15 min. Fig. 2 shows the results of such an experiment. The final Q_{O_2} ($\mu\text{l. O}_2$ consumed at s.t.p./mg. dry wt./hr.) was 54. The Q_{O_2} (glucose) varied from preparation to preparation but was usually in the range 50-58. No oxidation products of glucose other than carbon dioxide have as yet been found.

Effect of sodium azide on the rate of oxygen uptake. When a small amount of glucose (6.67 μmoles) was added to suspensions of starved hyphae the extra oxygen consumed was much less than that required for complete oxidation and corresponded to between 16 and 25 % of the theoretical value. This suggested that a considerable amount of the substrate was assimilated. Sodium azide is known to inhibit

Fig. 2. Comparison of time taken for the rate of O_2 consumption to become constant on the addition of 66.7 μmoles of glucose to carbohydrate-starved suspensions of baker's yeast and *Z. moelleri*. Arrow indicates addition of glucose. Flasks contained 3.1 mg. dry wt. of *Z. moelleri* or approx. 14 mg. wet wt. of yeast. 0.066M phosphate buffer, pH 6.8. A, yeast; B, *Z. moelleri*; C, *Z. moelleri* with no added glucose.

oxidative assimilation in other types of cell—see Pickett & Clifton (1943); Winzler (1940); Syrett (1951), and Reiner & Spiegelman (1947)—and Fig. 3 shows the effect of 10^{-3} M sodium azide on the oxidation of glucose by starved suspensions of *Z. moelleri*; it will be noted that the total amount of oxygen consumed was markedly increased but that the inhibitor had no effect on the lag.

Azide usually slightly inhibited the rate of oxygen uptake in the presence of glucose by carbohydrate-starved cells though this did not occur in the experiment recorded in Fig. 3. On the other hand, 10^{-3} M sodium azide increased the rate of endogenous oxygen uptake. If the hyphae were starved by incubation for 24 hr. in phosphate buffer pH 6.8, i.e. in the absence of both carbohydrate and a nitrogen source, then the maximum rate of oxygen uptake in the presence of glucose was unaffected by this concentration of azide. Except where otherwise stated, cells starved in phosphate buffer were used in all subsequent experiments, and such preparations will be referred to as starved cells or starved hyphae.

Effect of ammonium chloride on the oxidation of glucose by starved hyphae. Ammonium chloride increased the rate of oxidation of glucose by suspensions of starved hyphae but had no effect on the endogenous respiration (Fig. 4). The final rate of oxygen uptake attained was the same whether the ammonium chloride was added before, together

Fig. 3. Effect of NaN_3 on the total O_2 uptake when glucose is oxidized by carbohydrate-starved suspensions of *Z. moelleri*. 6.67 μmoles of glucose added to 6.0 mg. dry wt. of cells/flask as indicated by the arrow. 0.066M phosphate buffer, pH 6.8. A, in the presence of 10^{-3} M- NaN_3 ; B, no NaN_3 ; C and D, no glucose, with and without NaN_3 respectively.

with, or after the substrate. Ammonium sulphate had the same effect as an equivalent amount of ammonium chloride, indicating that the effect was due to the cation. This stimulatory effect of ammonium salts was inhibited by 10^{-3} M sodium azide (Fig. 4) and by 4.6×10^{-4} M cyanide, but not by 1.67×10^{-4} M 2:4-dinitrophenol at pH 6.8.

Fig. 5 shows that the increase in the rate of oxygen uptake is related to the amount of ammonium chloride added over the range 0–5 $\mu\text{moles/ml}$. In Fig. 5 the maximum rate of oxygen uptake in the presence of glucose and ammonium chloride, expressed as a percentage of the maximum rate attained in the presence of glucose alone, is plotted against μmoles ammonium chloride added.

If glucose was added to suspensions of starved hyphae in two separate quantities, each of 6.67 μmoles , and the second addition made when the rate of oxygen uptake had fallen to that of the endogenous, there was little or no lag period after the second addition. This was the case whether or not

Fig. 4. Effect of NH_4Cl on the rate of oxidation of glucose by carbohydrate- and nitrogen-starved cells in the presence and absence of NaN_3 . Glucose concn. 0.022M in all cases except E. Each flask contained 3.2 mg. dry wt. of cells in 0.066M phosphate buffer, pH 6.8. Where used 9.3 μmoles of NH_4Cl were added/flask. A, NH_4Cl present from start; B, NH_4Cl added at arrow 2; C, no NH_4Cl added. A, B, and C without NaN_3 . D, as A, B, and C all in the presence of 10^{-3} M- NaN_3 ; all these curves coincided. E, no glucose added, with and without NH_4Cl but without NaN_3 . Glucose added to A–D at arrow 1.

ammonium chloride was present—Fig. 6. It will be noted that, even with these small amounts of glucose, ammonium chloride had a slight stimulatory effect, and this is further borne out by the data presented in Table 1. The rates of oxygen uptake and the total oxygen uptake after several hours in replicate samples of the same batch of cells under the same conditions differed by not more than 2% from the mean.

That the break in the oxygen uptake curve coincided with the disappearance of glucose from the medium was shown by a semi-quantitative determination of the glucose remaining at the time when the rate of oxygen uptake had returned to the endogenous value. The Ihl-Dische reagent was used

Fig. 5. The effect of the amount of ammonium chloride on the rate of O_2 uptake by carbohydrate- and nitrogen-starved cells oxidizing glucose. Results expressed as percentage of control with no NH_4Cl . Two experiments shown. Glucose concn. $0.022M$ in all cases. Each flask contained 3–4 mg. dry wt. of cells in $0.066M$ phosphate buffer, pH 6.8.

and the blue colours produced by samples of the medium were compared by eye with a series of standards of suitable concentration. It was found that when the respiratory rate had fallen to that of the endogenous the glucose concentration in the medium was not more than $10 \mu g./ml.$, as compared with an initial concentration of $400 \mu g./ml.$

Fig. 6. Addition of two quantities of glucose, each of $6.67 \mu moles$, to fully starved cells in the presence and absence of added NH_4Cl . Each flask contained 3.7 mg. dry wt. of cells in $0.066M$ phosphate buffer, pH 6.8. Glucose added as indicated by the arrows. A, in the presence of $9.3 \mu moles$ of NH_4Cl present from the start; B, no NH_4Cl ; C, no glucose added, with and without NH_4Cl .

Table 1. *The reproducibility of the small stimulation by ammonium chloride of the rate of oxygen uptake when the glucose concentration was low*

Three experiments with different cultures were carried out. Each flask contained 3–4 mg. dry wt. of cells; on each separate occasion the dry wt./flask was constant. Where indicated $9.3 \mu moles$ of NH_4Cl were added to the flask; two consecutive quantities, each of $6.67 \mu moles$ of glucose were added to each flask. The medium contained $0.066M$ phosphate buffer, pH 6.8. Fig. 6 shows the progress curves of Expt. 1.

Expt.	Rate of O_2 uptake with 1st quantity of glucose ($\mu l./hr./flask$)	Ratio: B/A	Rate of O_2 uptake with 2nd quantity of glucose ($\mu l./hr./flask$)	Ratio: B/A
1. A No NH_4Cl B With NH_4Cl	116.0 } 136.6 }	1.18	114.3 } 156.6 }	1.37
2. A No NH_4Cl B With NH_4Cl	108.7 } 148.0 }	1.36	110.1 } 190.0 }	1.73
3. A No NH_4Cl B With NH_4Cl	120.0 } 158.0 }	1.32	113.2 } 165.9 }	1.47

Effect of ammonium chloride on the fermentation of glucose. Fig. 7 shows that ammonium chloride has no effect on the rate of carbon dioxide formation from glucose under anaerobic conditions; for comparison, the effect of ammonium chloride on the oxidation of glucose by a sample of the same suspension of starved hyphae is shown. It will be noted that a significant period of time elapsed before the rates of both carbon dioxide formation and of oxygen uptake became constant. Thus, despite the fact that under both aerobic and anaerobic conditions there was a lag in the metabolism of glucose, ammonium chloride exerts its stimulating effect only in the presence of oxygen.

Effect of pH on the rates of oxygen uptake. The effect of pH on the rates of endogenous oxygen uptake of starved cells and on the rates of oxygen

Fig. 7. Effect of NH_4Cl on the rates of O_2 uptake, and of CO_2 evolution under anaerobic conditions, of fully starved cells. A, D, and E, O_2 uptake. B, C, and F, CO_2 evolution. A and B, NH_4Cl present from start; C and D, no NH_4Cl ; E and F, no glucose or NH_4Cl . Curves B and C coincided. $9.3 \mu\text{moles}$ of NH_4Cl present in flasks where indicated. Each flask contained 2.8 mg. dry wt. of cells in 0.066 M phosphate buffer, pH 6.8. Glucose added to flasks A–D at arrow to give a concn. of 0.022 M.

uptake with glucose, both in the presence and in the absence of ammonium chloride, was examined. Phosphate buffers (0.2 M) at a range of pH values were prepared by adjusting solutions of potassium dihydrogen phosphate with hydrochloric acid or sodium hydroxide to the desired pH as measured with a glass electrode. The Warburg flasks were made up as follows: 1.0 ml. buffer, 1.0 ml. of a suspension of starved cells in 0.9% (w/v) NaCl and glucose and ammonium chloride solutions as required. The maximum constant rates of oxygen uptake were recorded and at the end of the experiment the pH of the medium in each flask was measured using a miniature glass electrode assembly capable of measuring pH on about 0.15 ml. of liquid only (Hanes, 1951). The results are given in Table 2.

Effect of other nitrogenous compounds on the oxidation of glucose. The effect of the addition of nitrate, nitrite, hydroxylamine, urea, L-aspartic acid, L-asparagine, and L-glutamic acid on the rate of glucose oxidation by suspensions of starved hyphae was next examined, and the results are given in Table 3. L-Glutamic acid and L-asparagine were about as effective as an equivalent amount of ammonia; urea and L-aspartic acid were slightly less effective, and nitrite, nitrate, and hydroxylamine did not increase significantly the rate of oxygen uptake.

Effect of ammonium chloride on the rate of oxidation of other substrates. Table 4 shows the effect of ammonium chloride on the rate of oxygen uptake by suspensions of starved hyphae in the presence of glucose, sucrose, fructose, acetate, and succinate. Since the suspensions did not oxidize succinate at pH 6.8 but did so at pH 3.5, the experiment with succinate recorded in this table was carried out at pH 3.5 (potassium dihydrogen phosphate adjusted to pH 3.5 with hydrochloric acid). Duplicate manometers were set up for each substrate and when the rate of oxygen uptake in the presence of substrate had become constant (120–135 min.) ammonium chloride was added to one of each pair of manometers, the other serving as a control. The experiment was continued until the rates of oxygen

Table 2. *Effect of pH on the rate of oxygen uptake under various conditions*

Each flask contained 7.2 mg. dry wt. of cells and 0.066 M phosphate buffer of the pH indicated. Glucose, where present, was at a concn. of 0.022 M. Where indicated $9.3 \mu\text{moles}$ of NH_4Cl were added to the flask. The maximum constant rate of O_2 uptake is recorded together with the initial and final pH of the medium in each flask.

Initial pH	Endogenous		With glucose		With glucose plus NH_4Cl	
	Q_{O_2}	Final pH	Q_{O_2}	Final pH	Q_{O_2}	Final pH
3.73	6.3	5.14	33.6	4.93	51.9	3.40
4.98	8.6	5.73	40.0	5.69	56.9	4.76
6.00	9.4	6.24	41.5	6.11	61.7	5.90
7.12	10.3	7.15	31.5	7.15	48.2	7.12
7.98	—	—	14.9	8.17	27.6	7.77
8.81	10.3	8.52	13.8	8.57	23.2	8.02

Table 3. *Stimulation of glucose oxidation by various nitrogenous compounds*

Each flask contained 6.52 mg. dry wt. of cells. The medium contained 0.066M phosphate buffer, pH 6.8, and 0.022M glucose in all cases. When the rates of O₂ uptake became constant the nitrogenous substances were added as indicated, the increased constant oxidation rate measured and the time interval between adding the nitrogen and the attainment of the increased oxidation rate noted.

Nitrogenous compound added	Nitrogenous compound added (μ moles)	Nitrogen added (μ atoms)	Q_{O_2}	Increase over control (%)	Time interval (min.)
None (no glucose)	0	0	6.4	—	—
None	0	0	26.4	—	—
NH ₄ Cl	9.3	9.3	53.1	101.1	144
NH ₄ Cl	4.65	4.65	40.6	54.1	92
KNO ₃	9.3	9.3	28.1	6.4	160
NaNO ₂	9.3	9.3	29.0	9.9	129
NH ₂ OH, HCl	9.3	9.3	23.2	-12.2	105*
Urea	4.65	9.3	44.6	69.2	120
L-Aspartic acid	9.3	9.3	40.6	54.1	180
L-Glutamic acid	9.3	9.3	49.9†	87.2	—†
L-Asparagine	4.65	9.3	47.2	81.4	123

* Initial inhibition of oxygen uptake followed by subsequent recovery.

† Rate of O₂ uptake still increasing very slowly 285 min. after adding L-glutamic acid.

Table 4. *Effect of ammonium chloride on the oxidation of several substrates*

Rates of oxidation of glucose, sucrose, fructose, acetate, and succinate in the presence and absence of NH₄Cl. Each flask contained 4.65 mg. dry wt. of cells. The medium contained 0.066M phosphate buffer, pH 6.8 (for succinate oxidation the pH was 3.5). Substrate concn. glucose, fructose, and acetate, 0.02M; sucrose and succinate, 0.01M. 9.3 μ moles of NH₄Cl added where indicated.

Substrate	Q_{O_2}		Increase of Q_{O_2} in presence of NH ₄ Cl	Percentage increase of Q_{O_2} in presence of NH ₄ Cl
	No NH ₄ Cl	With NH ₄ Cl		
None	5.0	5.0	0.0	0.0
Glucose	37.0	66.9	29.9	80.8
Sucrose	35.5	60.4	24.9	70.3
Fructose	11.0	12.7	1.7	15.7
Acetate	54.4	66.7	12.3	22.5
Succinate	29.8	31.2	1.4	4.6

uptake had again become constant (a further 90–120 min.). Fructose was oxidized very slowly and ammonium chloride had no effect on the rate of oxygen uptake. The rate of oxygen uptake in the presence of succinate was greater than that in the presence of fructose but it too was unaffected by the addition of ammonium chloride; however, since this experiment was carried out at pH 3.5, this observation may not be significant, although it was noted that ammonium chloride stimulated glucose oxidation at this pH—see Table 2. Glucose and sucrose were oxidized at approximately the same rates, and ammonium chloride had a marked stimulatory effect. Acetate was oxidized the most rapidly of all substrates tested (Q_{O_2} 54.4 as compared with 37.0 for glucose), but even so the rate of oxygen uptake was stimulated to some extent by ammonium chloride.

The fate of added ammonia nitrogen. The fate of ammonia nitrogen added to suspensions of starved hyphae was next examined and the effect of 10⁻³M sodium azide was also investigated. The suspensions plus additions were shaken in air for 3 hr., at the end of which time the distribution of nitrogen in both cells and suspending fluid was determined. The oxygen uptake was measured manometrically in parallel experiments. The results are given in Table 5. In the absence of azide, 78.8 % of the added ammonia nitrogen disappeared and 54.5 % was accounted for in the insoluble fraction. In the presence of azide assimilation of ammonia nitrogen was almost completely inhibited and 95.4 % of the ammonia nitrogen added was recovered as such. Azide had no effect on the distribution of nitrogen in the cells to which no ammonium chloride had been added. Since azide inhibits both the stimulatory action of ammonium chloride and the assimilation of ammonia nitrogen, it is possible that the stimulatory effect and the assimilatory process are related.

Effect of ammonium chloride on the respiratory quotient. Fig. 8 shows the effect of ammonium chloride on the R.Q. It will be noted that, on addition of ammonium chloride to suspensions of starved hyphae oxidizing glucose, the R.Q. increased from 1.14 to 1.29 and subsequently fell to a value slightly greater than the initial value. This effect is in complete contrast to that observed by Syrett (1953a) working with *Chlorella*. In this organism addition of ammonia resulted in a fall in the R.Q. from 1.29 to 0.749 followed by a rise to a value of 1.17 some 90–100 min. later.

Effect of pretreatment with glucose and ammonium chloride. A suspension of starved hyphae was divided into two equal parts, to one of which was added glucose and to the other glucose plus

Table 5. *Distribution of nitrogen in starved cells with and without the addition of an external source of ammonium chloride, in the presence of glucose, and in the presence and absence of sodium azide*

The cells were divided into four equal samples, each containing 24.0 mg. dry wt. of cells (N 8.24 % dry wt.). The cells were incubated in 15 ml. 0.066M phosphate buffer, pH 6.8, containing 0.022M glucose together with the following substances: sample 1: none; sample 2: 2.5 mg. of NH_4Cl (654 μg . N); sample 3: 0.001M- NaN_3 ; sample 4: 2.5 mg. of NH_4Cl + 0.001M- NaN_3 . The suspensions were shaken in air at 25°. 3 hr. after the addition of the NH_4Cl , samples of each suspension were removed for analysis and the amount of N in each fraction was determined. The O_2 consumption of a sample of each suspension was simultaneously followed.

Fraction	Sample 1 (μg . N)	Sample 2 (μg . N)	Difference between samples 1 and 2 (μg . N)	Difference expressed as % of N added (as NH_4Cl) to sample 2	Sample 3 (μg . N)	Sample 4 (μg . N)	Difference between samples 3 and 4 (μg . N)	Difference expressed as % of N added (as NH_4Cl) to sample 4
Free NH_4^+ in external medium	0.0	75.0	75.0	11.5	0.0	549.8	549.8	84.0
Free NH_4^+ in cells	12.0	19.5	7.5	1.1	11.3	85.5	74.2	11.3
Insoluble N in cells	1545.0	1901.3	356.3	54.5	1477.5	1455.0	-22.5	-3.4
Soluble N in cells	420.0	579.0	159.0	24.3	420.6	447.0	26.4	4.0
Total N	1977.0	2574.8	597.8	91.4	1909.4	2537.3	627.9	95.9
Final constant rate of O_2 uptake ($\mu\text{l.}/\text{hr.}/\text{flask}$)	187	330	—	—	191	207	—	—

Rate of O_2 uptake with no added NH_4Cl , glucose or NaN_3 : 54 $\mu\text{l.}/\text{hr.}/\text{flask}$.

Fig. 8. Change of R.Q. following addition of NH_4Cl to fully starved cells oxidizing glucose. 9.3 μmoles of NH_4Cl added to 3.9 mg. dry wt. of cells in 0.066M phosphate buffer: original pH 5.26; final pH 5.17. NH_4Cl added at arrow; 0.022M glucose present from start.

ammonium chloride. Both samples were shaken in air for 3 hr. at 25°, at the end of which time the hyphae were centrifuged off, washed and suspended in water. The ability of both suspensions to oxidize glucose, acetate, ethanol and succinate was then examined. In the case of succinate the pH of the medium was buffered at pH 3.5 as described above; the other three substrates were tested at pH 6.8 in the usual way. The results are given in Table 6. It will be seen that when glucose and acetate were the substrates, the rates of oxygen uptake were greater after pretreatment with glucose and ammonium chloride than after pretreatment with glucose alone. Ethanol oxidation showed a qualitatively similar response, though the effect was smaller; the rate of oxidation of succinate was similar after both types of treatment. It will also be seen that the dry weight of the hyphae was increased by both treatments.

Table 6. *Effect of treating cells with ammonium chloride and glucose on the dry weight and the rates of oxygen uptake with several substrates*

96.8 mg. dry wt. of cells were divided into two samples. Each sample was suspended in 15 ml. of 0.066M phosphate buffer, pH 6.8, and received 333.3 μmoles of glucose; sample 2 also received 77.5 μmoles of NH_4Cl . Both samples were shaken in air at 25° for 3 hr. The cells were then centrifuged, washed, and each sample suspended in 15 ml. of water. 1 ml. of cell suspension was used in each Warburg flask, which also contained 0.066M phosphate buffer, pH 6.8 (for succinate oxidation the pH was 3.5) and the substrate at a final concn. of 0.022M. Dry weight determinations were performed on the original cells before treatment, and on each of the two samples after treatment with glucose with or without NH_4Cl .

Substrate	$Q_{O_2}^*$		Percentage increase in sample 2 over sample 1
	Sample 1. Cells treated with glucose only	Sample 2. Cells treated with glucose + NH_4Cl	
Glucose	25.7	36.5	42.2
Acetate	42.4	50.5	19.1
Ethanol	16.7	20.4	22.3
Succinate	12.4	12.7	2.5

Dry weight of cells in equivalent volumes of suspension (mg.):

Before treatment	16.1
After glucose alone (sample 1)	18.5
After glucose + NH_4Cl (sample 2)	19.4

* Calculated on dry-weight basis of original untreated cells.

DISCUSSION

Washed suspensions of freshly harvested *Z. moelleri* have a high endogenous respiration and addition of glucose to such suspensions only caused a relatively small increase in the rate of oxygen uptake. On the other hand, if the suspensions were starved for 24 hr. in phosphate buffer at pH 6.8, the endogenous respiration was reduced and addition of glucose resulted in a steadily increasing rate of oxygen uptake which became constant after a period of 2–3 hr. This phenomenon is not peculiar to the aerobic metabolism of glucose by suspensions of starved hyphae, since a similar lag was observed when such suspensions metabolized glucose anaerobically. The fact that the lags observed under both aerobic and anaerobic conditions were of similar duration suggests that a common factor, or factors, are involved.

Sodium azide is known to inhibit oxidative assimilation, oxidative phosphorylation and adaptive enzyme formation (Clifton, 1946; Spiegelman, 1947; Spiegelman, Kamen & Sussman, 1948; Winzler, 1940, 1944; Winzler, Burk & du Vigneaud, 1944). This inhibitor at a concentration of $10^{-3}M$ had no effect on the time course of oxygen uptake in the presence of glucose. Consequently, this increase in the rate of oxygen uptake is not due to adaptive enzyme formation, and the cause of it remains unknown. But azide was not without action on the metabolism of glucose. It was found that it increased the total amount of oxygen consumed. In the absence of azide the total oxygen uptake was 16–25 % of that required for the complete oxidation of the glucose added to carbon dioxide and water, whereas in the presence of azide almost the theoretical uptake was observed, suggesting that this mould has very great assimilatory powers.

When ammonium chloride was added to a suspension of starved hyphae oxidizing glucose the final rate of oxygen uptake was somewhat greater than that obtained in the presence of glucose alone. But ammonium chloride had no effect on the endogenous rate of oxygen uptake; nor was the rate of anaerobic metabolism of glucose influenced by this compound. L-Glutamic acid, L-asparagine, and to a lesser extent, L-aspartic acid and urea showed effects similar to that of ammonium chloride on the rate of glucose oxidation; but nitrate, nitrite, and hydroxylamine had no action on this process. The effect was not specific for glucose; the rates of oxidation of sucrose, and, to a lesser extent, acetate were also increased by ammonium chloride. On the other hand, the rates of oxygen uptake in the presence of fructose and succinate were unaffected by this treatment.

Sodium azide at a final concentration of $10^{-3}M$ completely abolished the effect of ammonium ions

on the oxidation of glucose. Azide is known to inhibit synthetic processes and it would seem reasonable to suggest that the effect of ammonium ions is associated with the synthesis of some nitrogenous compound or compounds, possibly protein in nature, which are essential for the oxidation of the substrate. Analysis showed that the ammonium chloride added to suspensions oxidizing glucose almost completely disappeared, and that its disappearance was accounted for by an increase in the nitrogenous constituents of the hyphae. More than half of the ammonia nitrogen disappearing was accounted for by an increase in the nitrogen content of the insoluble fraction, an observation in keeping with the view that the action of ammonium chloride is associated with protein synthesis. It was particularly significant, therefore, that azide was found to inhibit not only the effect of ammonium chloride on respiration but also the assimilation of ammonia nitrogen. It is noteworthy in this connexion that Yamaguchi & Matsuzaki (1952), working on the adaptation of *Pseudomonas aeruginosa* to propionate, found that the developments of enzyme activity was stimulated by the addition of ammonium sulphate; below $4 \times 10^{-4}M$ there was practically a linear relationship between ammonium sulphate added and the maximum level of enzyme activity attained. Endogenous respiration was practically unaffected by ammonium sulphate. Spiegelman & Dunn (1947) found that the presence of an external nitrogen source stimulated the rate of adaptation and considerably raised the attainable level of enzyme activity in the adaptation of yeast to galactose and maltose fermentation. This stimulation was absent if the cells were unable to assimilate the added nitrogen, e.g. in the presence of azide.

It is not possible as yet to say whether general protein synthesis occurs in the presence of glucose and ammonium ions, or whether the effect is specific for those enzymes concerned with the oxidation of glucose. The results given in Table 6 show that pre-incubation of hyphae with glucose and ammonium chloride results in an increased rate of oxidation of both glucose and acetate, suggesting that the amounts of the enzymes concerned with the oxidation of these two compounds are increased. This must remain a suggestion until such time as methods have been perfected for the quantitative estimation of the amounts of these enzymes within the hyphae.

SUMMARY

1. *Zygorhyncus moelleri* was grown in a glucose-ammonium chloride medium and then starved for 24 hr. in a carbohydrate-free medium. When glucose was added to the starved cells an induction period of 2–3 hr. was observed before the rate of oxygen uptake rose to a constant maximum value.

2. In the presence of azide the final rate of oxygen uptake was somewhat lower than in its absence, although the length of the induction period was not greatly affected. These cells were not nitrogen-starved.

3. The respiration of suspensions starved of both carbohydrate and nitrogen was not inhibited by azide, but was greatly stimulated by small quantities of ammonium chloride. This stimulation was abolished by suitable concentrations of azide and cyanide.

4. Neither the endogenous respiration nor the fermentation of glucose under anaerobic conditions was affected by added ammonium chloride.

5. L-Glutamic acid, L-aspartic acid, L-asparagine, and urea stimulated glucose oxidation, though none so effectively as ammonia; nitrate, nitrite, and hydroxylamine were inactive.

6. The oxidation rates of sucrose, ethanol, acetate and succinate were also stimulated to varying degrees by ammonia, but the stimulation was much weaker in most cases than with glucose; only with sucrose did the stimulation approach that with glucose.

7. In the presence of glucose the ammonia was bound, and after 3 hr. appeared mainly in the insoluble nitrogen fraction; azide inhibited this binding of ammonia.

8. These results are discussed and it is suggested that enzyme adaptation is a possible interpretation of these phenomena consistent with the observed facts.

I wish to express my thanks to Prof. W. H. Pearsall, F.R.S., Mr P. J. Syrett, and Mr H. Tristram for their valuable help and criticism, and to the Department of Scientific and Industrial Research for a maintenance grant.

REFERENCES

- Barron, E. S. G., Ardao, M. I. & Hearon, M. (1950). *Arch. Biochem.* **29**, 130.
 Clifton, C. E. (1946). *Advanc. Enzymol.* **6**, 269.
 Darby, R. T. & Goddard, D. R. (1950). *Amer. J. Bot.* **37**, 379.
 Dixon, M. (1943). *Manometric Methods*, 2nd ed. Cambridge University Press.
 Dorrell, W. W. & Page, R. M. (1947). *J. Bact.* **53**, 360.
 Foster, J. W. (1949). *Chemical Activities of Fungi*, 1st ed. New York: Academic Press Inc.
 Hanes, C. S. (1951). *Nature, Lond.*, **168**, 566.
 Horowitz, N. H. & Beadle, G. W. (1943). *J. biol. Chem.* **150**, 325.
 Kluyver, A. J. & Perquin, L. H. C. (1933). *Biochem. Z.* **266**, 68.
 Lynen, F. (1942). *Liebigs Ann.* **552**, 270.
 Mann, T. (1944). *Biochem. J.* **38**, 339.
 Pickett, M. J. & Clifton, C. E. (1943). *J. cell. comp. Physiol.* **21**, 77.
 Reiner, J. M. & Spiegelman, S. (1947). *J. cell. comp. Physiol.* **30**, 347.
 Robbie, W. A. (1946). *J. cell. comp. Physiol.* **27**, 181.
 Spiegelman, S. (1947). *J. cell. comp. Physiol.* **30**, 315.
 Spiegelman, S. & Dunn, R. (1947). *J. gen. Physiol.* **31**, 153.
 Spiegelman, S., Kamen, M. D. & Sussman, M. (1948). *Arch. Biochem.* **18**, 409.
 Syrett, P. J. (1951). *Ann. Bot., Lond., N.S.*, **15**, 473.
 Syrett, P. J. (1953a). *Ann. Bot., Lond., N.S.* **17**, 1.
 Syrett, P. J. (1953b). *Ann. Bot., Lond., N.S.*, **17**, 21.
 Umbreit, W. W., Burris, R. H. & Stauffer, J. F. (1949). *Manometric Techniques and related Methods for the Study of Tissue Metabolism*, 2nd ed. Minneapolis: Burgess Publishing Co.
 Winzler, R. J. (1940). *J. cell. comp. Physiol.* **15**, 343.
 Winzler, R. J. (1944). *Science*, **99**, 327.
 Winzler, R. J., Burk, D. & du Vigneaud, V. (1944). *Arch. Biochem.* **5**, 25.
 Yamaguchi, S. & Matsuzaki, E. (1952). *Misc. Rep. Res. Inst. nat. Resour. Tokyo*, no. 26, 47.

Studies on Yeast Metabolism

3. THE INTRACELLULAR LEVEL OF PYRUVATE DURING YEAST FERMENTATION

By W. E. TREVELYAN AND J. S. HARRISON

Research and Development Department, Distillers Co. Ltd., Epsom, Surrey

(Received 30 November 1953)

The rate of carbon dioxide production during alcoholic fermentation is a direct measure of the velocity of the yeast carboxylase system, i.e. of the reaction:

This is irreversible (Vennesland, 1951); thus, in the steady state during fermentation, the rate of pyruvate decarboxylation is likely to be determined by the substrate concentration at the enzyme

surface. Unfortunately, this quantity cannot be directly determined. The amount of pyruvate in unit weight of yeast at any stage during fermentation is, in principle, determinable. This intracellular level of pyruvate may or may not be simply related to the kinetically important concentration at the site of carboxylase action. If it is, then its measurement at different fermentation rates should give information about the kinetics of yeast carboxylase *in vivo*.