acetyl-o-aminobenzophenone (IV) as colourless long needles, m.p. 107–108°. (Found: C, 65.8; H, 4.3; N, 5.8; Cl, 13.2. $C_{15}H_{11}O_2NCl$ requires C, 65.8; H, 4.4; N, 5.1; Cl, 13.0%.)

Cyclization of the chloroacetyl derivative was effected by dissolving it (5.36 g.) in hot ethanol (50 ml.) and diluting with water (140 ml.). When the solution became clear again, NaOH (1.5 g.) in water (10 ml.) was added, and a yellow colour developed owing to partial hydrolysis. The solution was refluxed for 30 min. and acidified with HCl while still hot. On cooling, colourless, chlorine-free, prismatic needles separated and were collected (4.3 g., 72% overall yield, m.p. 240–242°). Crystallization from ethanol gave synthetic viridicatin, m.p. 264–265°, not depressed on admixture with natural viridicatin from *P. cyclopium*. (Found: C, 75.7; H, 4.7; N, 6.1. Calc. for $C_{15}H_{11}O_2N$: C, 75.9; H, 4.7; N, 5.9%.) The two specimens also gave identical intense emerald-green ferric reactions. The ON-dimethyl derivative of synthetic viridicatin, prepared as described above, melted at 197–198° and gave no depression in m.p. on admixture with the ON-dimethyl derivative of natural viridicatin of the same m.p. (Found: C, 76.8; H, 5.7; N, 5.9. Calc. for $C_{17}H_{15}O_2N$: C, 76.95; H, 5.7; N, 5.3%.)

SUMMARY

1. A hitherto undescribed mould metabolic product, cyclophenin, has been isolated from culture filtrates of laboratory cultures of a strain of *Penicillium cyclopium* Westling.

2. Cyclophenin, $C_{17}H_{15}O_2N$, forms colourless tablets, m.p. 207°, $[\alpha]_{D}^{20} - 306^\circ$ in ethanol.

3. Cyclophenin is readily decomposed by dilute mineral acids giving one mole each of carbon dioxide, methylamine and viridicatin, $C_{15}H_{11}O_2N$, which is optically inactive.

4. Viridicatin, previously isolated from *P. viridicatum* Westling, by Cunningham & Freeman (1953), has now been prepared from five different strains of *P. cyclopium* but appears to be absent from the strain of this species which gives cyclophenin.

5. Viridicatin is oxidized in alkaline solution by gaseous oxygen to one mole each of o-aminobenzophenone and oxalic acid.

6. Viridicatin has been synthesized and has been shown to be 2:3-dihydroxy-4-phenylquinoline or its keto tautomer.

7. The five strains of *P. cyclopium* which gave viridicatin also gave good yields of palitantin and two of them gave the anti-fungal substance, frequentin.

8. Two possible, closely related, structures are suggested for cyclophenin.

We wish to thank Sir Robert Robinson, O.M., F.R.S., for his friendly advice on the structure of cyclophenin.

REFERENCES

- Alsberg, C. L. & Black, O. F. (1913). *Bull. U.S. Bur. Pl. Ind.* no. 270, 1.
 Anslow, W. K., Breen, J. & Raistrick, H. (1940). *Biochem. J.* **34**, 159.
 Birkinshaw, J. H. (1952). *Biochem. J.* **51**, 271.
 Birkinshaw, J. H., Callow, R. K. & Fischmann, Catherine F. (1931). *Biochem. J.* **25**, 1977.
 Birkinshaw, J. H., Oxford, A. E. & Raistrick, H. (1936). *Biochem. J.* **30**, 394.
 Birkinshaw, J. H. & Raistrick, H. (1932). *Biochem. J.* **26**, 441.
 Birkinshaw, J. H. & Raistrick, H. (1936). *Biochem. J.* **30**, 801.
 Birkinshaw, J. H., Raistrick, H., Ross, D. J. & Stickings, C. E. (1952). *Biochem. J.* **50**, 610.
 Brian, P. W. & Hemming, H. G. (1945). *Ann. appl. Biol.* **32**, 214.
 Campbell, A. H., Foss, M. E., Hirst, E. L. & Jones, J. K. N. (1945). *Nature, Lond.*, **155**, 141.
 Camps, R. (1899). *Arch. Pharm., Berl.*, **237**, 683.
 Camps, R. (1902). *Arch. Pharm., Berl.*, **240**, 145.
 Cunningham, K. G. & Freeman, G. G. (1953). *Biochem. J.* **53**, 328.
 Curtis, P. J. & Duncanson, L. A. (1952). *Biochem. J.* **51**, 276.
 Curtis, P. J., Hemming, H. G. & Smith, W. K. (1951). *Nature, Lond.*, **167**, 557.
 Fournneau, E. & Sabetay, S. (1928). *Bull. Soc. chim. Fr.* **43**, 859.
 Oxford, A. E. & Raistrick, H. (1935). *Biochem. J.* **29**, 1599.
 Oxford, A. E., Raistrick, H. & Smith, G. (1942). *Chem. & Ind.* **61**, 485.
 Scheifele, H. J. & DeTar, D. F. (1952). *Org. Synth.* **32**, 8.

The Purification of Alkaline Phosphatases of Animal Tissues

By R. K. MORTON*

Department of Biochemistry, University of Cambridge

(Received 5 October 1953)

Despite the fact that much of the considerable literature concerning alkaline phosphatases is devoted to methods for purification, no procedure for obtaining a homogeneous preparation from animal tissues has so far been described. This may possibly be due to the fact that the cytoplasmic

alkaline phosphatases are intimately associated with insoluble cellular granules. Special procedures must therefore be used to obtain these enzymes in true solution in order to permit of purification.

The difficulty of extracting alkaline phosphatases from tissue 'homogenates' was recognized by early investigators (see Ehrensward, 1933). It was found, however, that various forms of proteolysis, such as controlled autolysis (Albers & Albers, 1935) and

* Present address: Department of Biochemistry, University of Melbourne, Australia.

digestion with trypsin (Ehrensward, 1933; Schmidt & Thannhauser, 1943) enabled release of some of the enzyme into true solution. This was confirmed by Kabat (1941) who showed that alkaline phosphatase of mouse kidney is sedimented by centrifuging at 27 000 rev./min. for 1 hr. while, after prolonged autolysis, most of the enzyme remained in the supernatant.

As will be indicated in the Discussion, however, there are numerous objections to the use of proteolytic procedures for separating enzymes from particulate material in order to permit of their purification. Therefore, when a purified preparation of alkaline phosphatase of calf intestinal mucosa was required for studies of the relationship between the hydrolytic and transferase activities (see Morton, 1953*a*), other methods of separating the enzyme from the insoluble particles were sought. It was confirmed that phosphatase of calf intestinal mucosa (as well as of other tissues) was associated with insoluble cellular particles. By treatment with butanol the enzyme was quantitatively released into true solution. Subsequent fractionation yielded a preparation which was more active than any previously described and the evidence available suggested that it was essentially homogeneous. The principles of the purification procedure have been briefly reported (Morton, 1950).

MATERIALS AND METHODS

Enzymic activities

Alkaline phosphatase. Both the unit of activity and the specific activity have been defined in a previous paper (Morton, 1953*b*). Activities were determined essentially as therein described, using 4.8 ml. of buffered substrate and 0.2 ml. of suitably diluted enzyme. Final concentrations of the reactants were: sodium β -glycerophosphate (0.02*M*), either veronal-HCl (0.04*M*) or ethanolamine-HCl buffer (0.02*M*), pH 9.5 at 38°, and magnesium acetate (0.001 or 0.01*M*). The lower concentration of Mg^{2+} was used in testing the crude enzyme preparation, since higher concentrations are inhibitory to these (Abul-Fadl & King, 1949*a*; Morton, 1952). The amount of enzyme was adjusted so that hydrolysis was never more than 5% of the total substrate, and was usually less than 1%. Under these conditions, the rate of hydrolysis is proportional to enzyme concentration.

Other enzymes. Diphosphopyridine nucleotide-cytochrome *c* reductase, diaphorase, and succinic dehydrogenase activities were determined essentially as previously described (Morton, 1954), except that reduction of dyes was estimated visually, the tests being qualitative only.

Chemical estimations

Inorganic phosphate was determined by the method of Fiske & Subbarow (1925).

Nitrogen was estimated by a micro-Kjeldahl procedure or by a modification of the method described by Johnson (1941) using Nessler reagent. For rapidly following purification procedures, the maximum absorption between 270 and 280 $m\mu$. (usually 278 $m\mu$.), as determined in a Beckman

spectrophotometer, was used as a measure of protein concentration. The absorption values were converted to protein nitrogen by frequent calibration using the procedures previously mentioned.

Dry weights were determined after freezing and drying *in vacuo* over conc. H_2SO_4 for 48 hr. at room temp. (18°).

Lipid was estimated by the loss in dry weight after repeated extraction of the trichloroacetic acid precipitate with ether-ethanol (1:3) at room temp.

EXPERIMENTAL AND RESULTS

Localization of alkaline phosphatase of calf intestinal mucosa

The intestine of a 3-day-old calf was collected as soon as possible after killing. The convolutions of the intestine were removed by cutting away the mesenteric tissue with scissors and the small intestine was thoroughly rinsed by forcing a strong jet of running tap water through it. Portions of the intestine, about 2 ft. in length, were laid flat on an enamelled tray and the mucosa forced out by scraping the intestine with the edge of a plastic spatula. The mucosa was transferred to a beaker and cooled to 0° in an ice bath.

About 20 g. of mucosa were suspended in 200 ml. of 0.25*M* sucrose (at 0°) and gently stirred with a mechanical stirrer for 15 min. The sucrose suspension was then fractionated by differential centrifugation as shown in Fig. 1.

Mucosa (20 g.) suspended in 0.25 *M* sucrose (200 ml.).
Stirred 15 min. with mechanical stirrer at slow speed

Fig. 1. Fractionation of intestinal mucosa.
All procedures carried out at 0–2°.

Because of the limited centrifugal force (maximum about 14 000 *g*) which was available, many of the smaller particles could not be completely sedimented, and the final supernatants were either cloudy or opalescent. Therefore, a study of the quantitative distribution of alkaline phosphatase did not seem justified, and instead the specific activities of the various fractions were determined. Qualitative tests were carried out for diaphorase, diphosphopyridine nucleotide-cytochrome *c* reductase and succinic dehydrogenase activities. The results are shown in Table 1.

A portion of each of the fractions *A*, *B* and *C* was fixed by addition of neutral formaldehyde (40%) to give a final concentration of 2% in the sucrose suspension. The particles were then sedimented by centrifuging (14 000 *g*, 1 hr., 0–2°), washed twice with 40 ml. water and finally suspended in water (5 ml.). Various dilutions of each fraction were then transferred to cellulose nitrate films, the specimens dried *in vacuo*, and shadowed with gold–palladium at an angle of approx. 20°. They were examined in a Siemens electron-microscope.

The particles of fraction *A* were of irregular size and shape and were largely aggregated. Among the fields examined there appeared to be few particles less than 200 *mμ*. in diameter. The various dilutions of fraction *B* showed particles which were mostly of about 100–300 *mμ*. in

diameter, although some were less than 100 *mμ*. There was some aggregation. Fraction *C* contained many particles which appeared to be spherical. They ranged from about 200 *mμ*. to less than 20 *mμ*. in diameter with a mode of about 80 *mμ*.

Localization of alkaline phosphatase of cow mammary gland

A mammary gland from a lactating cow, obtained from the slaughter house within 2 hr. of death, was cooled in ice. Two sections of tissue from different quarters were minced and washed twice in 0.25 *M* sucrose (at 0°) to remove milk. About 100 g. of mince was then disintegrated for 1 min. in a Waring Blender in 500 ml. of sucrose (0.25 *M*). The 'homogenate' was centrifuged at 2500 *g* for 20 min., at 0° and the precipitate discarded. The supernatant was centrifuged at 14 000 *g* for 15 min. and the precipitate (fraction *M*₁) resuspended in 5 ml. sucrose (0.25 *M*). The supernatant was then centrifuged at 20 000 *g* for 1 hr. in the high-speed attachment of an International centrifuge and the precipitate (fraction *M*₂) resuspended in 5 ml. sucrose. The final supernatant (fraction *M*₃) was still slightly turbid.

The specific alkaline phosphatase activities of the fractions are shown in Table 2.

Table 1. *Properties of fractions from calf intestinal mucosa*

Cytochromes observed in a thick smear of mucosa or precipitate, reduced with Na₂S₂O₄, using a low-dispersion micro-spectroscope. Enzymes estimated as described under Methods. + Indicates qualitative test positive; – indicates negative.

	Whole 'homogenate' in 0.25 <i>M</i> sucrose	Fractions (see Fig. 1)		
		<i>A</i> (mitochondria)	<i>B</i> (mixed particles)	<i>C</i> (microsomes)
Approximate size of particles	Not determined	300 <i>mμ</i> .	300–100 <i>mμ</i> .	200–20 <i>mμ</i> .
Appearance of particles	Not examined	Irregular clumps	Some large particles and aggregates	Spheroidal isolated particles
Lipid (%)	Not determined	Not determined	Not determined	29
Cytochrome components	<i>a</i> , <i>c</i>	Not determined	Not detected	Not detected
Alkaline phosphatase (units/mg. N)	72	50	240	640
Diphosphopyridine nucleotide–cytochrome <i>c</i> reductase	+	Not determined	+	+
Diaphorase	+	Not determined	+	+
Succinic dehydrogenase	+	+	+	–

Table 2. *Properties of fractions from cow mammary gland*

For explanation of symbols, see Table 1. Xanthine oxidase tested as a dehydrogenase (see Methods).

	Whole 'homogenate' in 0.25 <i>M</i> sucrose	Fraction, and conditions of sedimentation		
		<i>M</i> ₁ 14 000 <i>g</i> , 15 min. (mitochondria)	<i>M</i> ₂ 20 000 <i>g</i> , 1 hr. (microsomes)	<i>M</i> ₃ Remaining supernatant
Appearance of precipitates	Salmon-pink in colour	Tan	Brown	Opalescent
Cytochrome components	<i>a</i> , <i>c</i>	<i>a</i> , <i>b</i> , <i>c</i>	Not detectable	Not detectable
Alkaline phosphatase (units/mg. N)	11	20	65	4
Xanthine oxidase	+	+	+	+
Diaphorase	+	+	+	–
Succinic dehydrogenase	+	+	+	–

Separation of alkaline phosphatase activity from particulate components of animal tissues

In a previous paper (Morton, 1953b) it was shown that treatment of milk or milk products with *n*- or *iso*-butanol released the alkaline phosphatase into true solution from the lipoprotein particles. The same method was therefore examined with dispersions of several different tissues from the rat and rabbit. The tissue was ground in a mortar with an equal weight of fine acid-washed sand, and with twice its weight of 0.05M veronal-HCl buffer at pH 8.0. A small sample of this dispersion was removed. A volume of *n*-butanol equal to the weight of tissue was then added, the material stirred at 35° for 15 min. and separated into distinct phases by centrifuging (2500 g, 45 min., at room temp.). The volume of the clear aqueous phase was measured, and a sample removed for alkaline phosphatase determination. The aqueous extracts contained over 80% of the total alkaline phosphatase of the untreated dispersions, while complete recovery of the enzyme could be obtained by re-extraction with butanol saturated 0.1M-NaHCO₃. Moreover, the enzyme was in true solution, since neither centrifuging in the high-speed attachment of the International centrifuge (20 000 g, 1 hr.), nor slow filtration through a thick bed of Hyflo Supercel (Johns Mansville and Co.) reduced the enzymic activity. In some cases, especially when intestinal dispersions were used, the aqueous layer remaining after butanol treatment of the tissue homogenate was slightly turbid even after prolonged centrifuging at 2500 g. This turbidity appeared to be due to mucous material which, however, was not associated with the enzyme, since slow filtration through Hyflo Supercel yielded a clear aqueous extract containing the total enzyme. The results of this survey of the effectiveness of butanol for release of the enzyme from various tissues are summarized in Table 3.

The aqueous extracts obtained after butanol treatment of animal tissues may be dialysed free of butanol and then frozen and dried *in vacuo* with very little loss of phosphatase activity. Moreover, the butanol-saturated aqueous material may be stored at 0–2° for considerable periods (experimental observations continued for 8 months in one case) without visible bacterial contamination and with only slight decline in activity.

Purification of alkaline phosphatase of calf intestinal mucosa

Calf intestines were collected from the slaughter house as soon as possible after killing, and the mesenteric membranes removed. The mucosa was then obtained as described

Table 3. *Alkaline phosphatase activities of aqueous extracts of animal tissues treated with butanol*

The tissue 'homogenates' (at pH 7.4) were treated with excess *n*-butanol at 35° for 15 min., centrifuged, and the aqueous layer was removed and tested (see above).

Tissue	Activity of extract		Recovery of enzyme (%)
	(Units/ml.)	(Units/mg. N)	
Rat kidney	48	35	87
Rabbit kidney	118	60	—
Rabbit liver	59	26	82
Rat intestine	95	30	94
Rabbit heart	17	11	—
Rabbit skeletal muscle	5	—	—

previously, transferred to a stainless-steel beaker and immediately cooled to 0–2° in a refrigerated bath. About 3 l. of mucosa were obtained from the intestines of forty calves.

The 'homogenate' of mucosa rapidly undergoes autolysis, which releases some of the phosphatase from the particulate material. This is to be avoided, not only because it results in loss of enzyme, but also because it is desirable to prevent contamination of the enzyme by products of proteolysis. It is therefore necessary to proceed with the preparation as quickly as possible after collection of the mucosa.

Stage 1. The mucosa was dispersed in 3 vol. of water (at 0°) and adjusted to pH 7.5 by slow addition of *N*-NaOH. After vigorous mechanical stirring for about 30 min. at 0°, the 'homogenate' was centrifuged (1800 g, 40 min.) in aluminium cups (1300 ml. capacity) in an International Serum Centrifuge (total capacity 13 l.). The supernatant was poured off and filtered through a layer of well-washed cotton wool on a Büchner funnel in order to remove loose fat and other insoluble aggregates. The filtrate was immediately cooled to 0° in a stainless-steel container held in a cold bath at –10°. The precipitate, which contained no more than 10% of the phosphatase of the 'homogenate', was discarded.

The initial disintegration of the mucosa was obtained using a 4-blade stainless-steel stirrer, driven at high speed by a 0.05 h.p. motor. Alternatively, the mucosa may be disrupted in a Waring-type blender for about 2 min., using 3 vol. distilled water.

Stage 2. The filtrate at 0–2° was adjusted to pH 5.0 by addition of 2M acetate buffer, pH 4.0, held for 45 min. at 0° and then centrifuged (1800 g, 40 min. at room temp.). The white precipitate was then thoroughly dispersed in 8 l. of 0.15M-NaCl. The suspension was adjusted to pH 7.5 with 0.5M-Na₂CO₃, cooled to 0° and stirred gently at 0–2° overnight. The enzyme was then reprecipitated at pH 5.0 as before, and similarly collected by centrifuging. At this stage the supernatant was clear and almost free of soluble protein. The precipitate was then washed in the cups by rubbing up into 8 l. of distilled water (at 0°) and immediately recentrifuged as previously. By this means the ionic strength of the material was lowered, and the last traces of soluble protein were removed.

This precipitate was again dispersed in distilled water at 0° and made up to 5 l. The white dispersion (now at pH 5.5) was at this stage free of mucous material and of low viscosity, while all traces of blood, pigments and soluble proteins had been removed.

As indicated in the Discussion it is preferable to sediment the particles carrying alkaline phosphatase by high-speed centrifuging at neutral pH, rather than by precipitating as described above. Therefore, where suitable equipment is available for centrifuging relatively large volumes at high speeds, the following modification (stage 2a) is recommended.

Stage 2a. The filtrate from stage 1 is centrifuged at 10 000 g for 30 min. at 0–4°. The supernatant is then centrifuged at 40 000–60 000 g for 1–2 hr. The precipitate is resuspended in 5 vol. of 0.15M-NaCl using a Waring-type blender (1 min.), the suspension readjusted to pH 7.2 with 0.2M-NaHCO₃, and gently stirred overnight at 0–2°. The particles are again sedimented (40 000–60 000 g, 1 hr.). The precipitate is suspended in 5 vol. of distilled water, the suspension stirred for 1 hr. at 0–2°, and finally recentrifuged as before. The precipitate is suspended in 3 vol. of distilled water and then treated as described under stage 3.

Stage 3. About 2 l. of *n*-butanol were slowly added over a period of 15 min. with vigorous mechanical stirring. The material was then heated to 38° for 5 min. and immediately centrifuged in 1 l. glass cups (1800 g, 30 min.). The clear, colourless aqueous layer was removed by suction and filtered through a thick layer of Hyflo Supercel over no. 1 Whatman paper on a 1 l. Büchner funnel. The filtrate was then adjusted to pH 8.5, held at 0° overnight and again filtered through Hyflo Supercel to remove a small precipitate.

Stage 4. The filtrate was adjusted to pH 6.4, cooled to -5° and the enzyme directly precipitated at -5° by addition of ether (10%, v/v) and acetone (60%, v/v). After settling overnight at -7°, the supernatant was removed by suction, and the precipitate collected by centrifuging at -5°.

It is often convenient to dry the precipitate at this stage and to accumulate material from several batches of mucosa before proceeding further with purification. For this purpose, the precipitate is washed twice with dry acetone at -15° followed with dry anaesthetic ether at -15°, and then dried, initially in a stream of N₂ and finally *in vacuo* over H₂SO₄ and CaCl₂. The dry powder may be stored over NaOH and CaCl₂ *in vacuo* at 0°. It retains activity for at least 12 months. For further purification it is treated exactly as the undried precipitate.

Usually, drying was omitted and the 60% (v/v) acetone precipitate was dissolved in a minimum of 0.05 M veronal-HCl buffer at pH 6.4 and dialysed against 0.015 M magnesium acetate, at pH 6.4. Insoluble material was removed by centrifuging (20 000 g, 1 hr., at 2°).

Stage 5. Following addition of ether (10%, v/v) the enzyme was precipitated at -5° with acetone between 35 and 48% (v/v). This fraction was dissolved in 0.05 M veronal-HCl buffer at pH 6.4 and dialysed against glass-distilled water (at 0°) until free of organic solvent.

Stage 6. The solution was adjusted to pH 4.9 with 0.05 N acetic acid, and heated rapidly to 48° for 2 min. A slight opalescence appeared. Magnesium acetate (0.5 M) was added to give a final concentration of 0.015 M, the solution adjusted to pH 6.4 and the enzyme precipitated at -5° with acetone between 40 and 50% (v/v). The slight precipitate at 40% (v/v) acetone was discarded. The precipitate between

40 and 50% (v/v) acetone was dissolved in buffer and dialysed free of organic solvent as for stage 5.

Stage 7. The solution was adjusted to pH 5.5 with 0.5 N acetic acid, and successive 20 mg. lots of washed charcoal, prepared as previously described (Morton, 1953c) were added slowly after mixing to a thick slurry in distilled water. Each precipitate was removed by centrifuging (5000 g, 15 min., at 2°) and the procedure repeated until about 30% of the enzymic activity was adsorbed. The solution was then centrifuged (14 000 g, 30 min.), adjusted to pH 8.0 with 0.1 N-NaOH, and filtered on a Büchner funnel through a thin layer of magnesium carbonate over a no. 1 Whatman filter paper in order to remove colloidal charcoal.

Stage 8. The enzyme solution was adjusted to pH 6.4 with 0.1 N acetic acid and dialysed at 2° against 0.015 M magnesium acetate for 4 hr. with continuous stirring. The solution was cooled to 0°, ether added to 10% (v/v) and the enzyme precipitated at -5° with acetone between 40 and 48% (v/v). The precipitate was dissolved in glass-distilled water, and dialysed against glass-distilled water (at 0°) for 4 hr. with frequent changes of water.

A summary of the procedure and the results for preparation no. 6 are reported in Table 4. Preparation no. 6 gave a final product of specific activity of 83 500 units/mg. N (*Q_p* 579 000) compared with the initial supernatant of activity 97 units/mg. N, representing a purification on this basis of 860 times. Since much inert material was removed during the first centrifuging, the total purification compared to the activity of the original mucosa was well over 1000 times. On the basis of non-dialysable dry weight, the total purification was estimated to be about 1700 times. The final yield was 6% of the activity of the first supernatant.

The dialysed solution from stage 8 (Table 4) was frozen and dried *in vacuo* over H₂SO₄. The dried material was stored *in vacuo* over CaCl₂ at 0°. This preparation lost about 40% of its activity after storage for 12 months in the dry state. It is noteworthy that dilute solutions of pure alkaline phosphatase may decline considerably in activity when held at -15°, apparently due to irreversible protein denaturation.

During investigation of various purification procedures, alkaline phosphatase was prepared by adding *n*-butanol

Table 4. *Purification of alkaline phosphatase from calf intestinal mucosa*

Stage	Vol. (ml.)	Activity (units/ml.)	Total units (× 10 ³)	Total N (mg./ml.)	Specific activity (units/mg. N)	Yield (%)	Purification
1	12 000	205	2 460	2.12*	97	100	1
2	5 000	216	1 080	0.82	263	44	3
3	4 450	166	739	0.03	5 533	30	57
4	55	12 360	680	2.05	6 029	28	62
5	20	26 250	525	1.96	13 393	21	138
6	20	18 900	378	0.94	20 106	15	208
7	32	7 340	235	0.11	66 727	10	688
8	10	14 200	142	0.17	83 529	6	860

* Nitrogen estimated after dialysis, 24 hr. at 0°, against distilled water.

Stage

Summary of procedures

- Supernatant from 3 l. of mucosa in about 9 l. distilled water, pH 7.5, centrifuged 1800 g, 40 min.
- Precipitate at pH 5.0, repeatedly washed and suspended in distilled water.
- Filtered aqueous layer following butanol treatment, held pH 8.5, 14 hr. at 0° and refiltered.
- Dialysed solution of precipitate obtained with ether (10%, v/v) and acetone (60%, v/v).
- Dialysed solution of fraction obtained with ether (10%, v/v) and acetone (35–48%, v/v).
- Dialysed solution after heating to 48°, 2 min. at pH 4.9 and fractionation with acetone (40–50%, v/v).
- Filtrate following adsorption of inert protein on charcoal at pH 5.5.
- Dialysed solution of fraction obtained with ether (10%, v/v) and acetone (40–48%, v/v).

Table 5. *Partial purification of alkaline phosphatase from calf intestinal mucosa*

Stage	Vol. (ml.)	Activity (units/ml.)	Total units ($\times 10^3$)	Total N (mg./ml.)	Specific activity (units/mg. N)	Yield (%)	Purification
1	2 000	600	1 200	7.70	78	100	1
2	1 750	555	971	0.77	721	81	9
3	1 900	465	885	0.28	1 661	74	21
4	105	7 850	824	2.92	2 688	69	35
5	30	17 200	516	2.70	6 370	43	82
6	20	22 000	440	1.90	11 579	37	149
7	18	16 900	304	0.47	35 957	25	463
8	10	26 500	265	0.69	38 406	22	493

Summary of procedures

Stage	
1	Mucosa (1 l.) disintegrated in 1 l. distilled water and adjusted to pH 6.5.
2	<i>n</i> -Butanol (1.5 l.) added, vigorously stirred, and heated to 38° for 5 min., centrifuged at 1800 g, 40 min. Aqueous layer filtered through Hyflo Supercel.
3	Filtrate adjusted to pH 4.9 and filtered through Hyflo Supercel; filtrate adjusted to pH 8.6, held 12 hr. at 0°, refiltered as previously.
4	Enzyme (at pH 6.4) precipitated with ether (15%, v/v) and acetone (52.5%, v/v) at -5°. Precipitate dissolved in and dialysed against distilled water.
5	Solution at pH 4.9 heated to 48° for 2 min., cooled to -5°, adjusted to pH 6.4 and fractionated with ether (10%, v/v) and acetone (35-40%, v/v).
6	Dialysed enzyme solution fractionated at -5° with ether (10%, v/v) and acetone (40-45%, v/v). Precipitate dissolved and dialysed against distilled water.
7	Solution adjusted to pH 5.5 and treated with washed charcoal until about 30% enzymic activity adsorbed.
8	Solution adjusted to pH 6.4 and fractionated with acetone (40-53%, v/v) at -5°. Precipitate dissolved and dialysed against distilled water.

Table 6. *Summary of partial purification of alkaline phosphatases from ox kidney, liver and mammary gland*

Procedure	Specific activities of fractions (units/mg. non-dialysable N)		
	Kidney	Liver	Mammary gland
1 Aqueous suspension of washed particles	129	54	Not determined
2 Clear aqueous solution following <i>n</i> -butanol treatment	925	260	312
3 Solution heated, 42°, 5 min. at pH 6.8. Held pH 8.6, 8 hr., 2°	Not applied	Not applied	633
4 Precipitate of total phosphatase with ether and acetone, -7°	1300	Not determined	1780
5 Treatment at pH 4.9, 30 min., 18°, followed by pH 8.5, 12 hr., 0°	2800	930	Not applied
6 Fractionation with acetone at -5°	6900	2800	4400

directly to the 'homogenate' of intestinal mucosa in distilled water. Further fractionation gave preparations which were more active than those previously obtained by other workers. However, it was not possible to isolate the purified enzyme by this procedure, the maximum activity obtained in this way being 38 400 units/mg. N. This seems to be due to the failure to remove the soluble protein before release of the particle-bound enzyme. Apparently the soluble material contains a number of proteins which behave similarly to alkaline phosphatase during fractionation. Preparations by this method are nevertheless suitable for very many purposes and the enzyme is recovered in high yield (about 22%). Moreover, the preparative procedures do not necessitate the use of large-capacity, or high-speed centrifuges. Therefore, an outline of the procedures used and the results obtained is given in Table 5.

Partial purification of the alkaline phosphatases of ox kidney, liver and mammary gland

Using procedures essentially similar to those described above, and on the basis of experience gained in purification of milk phosphatase (Morton, 1953*d*), alkaline phosphatases were partially purified from kidney, liver and mammary gland. In each case the phosphatase was obtained in solution by treatment of the aqueous suspension of precipitated

particles with *n*-butanol. Further purification involved: (1) precipitation with ether and acetone at -7°; (2) exposure to pH 4.9 at room temperature (about 18°) for 30 min., and subsequently to pH 8.5 for 12 hr. at 0°; (3) fractionation with acetone at pH 6.4 and at -5°, retaining the most active fractions.

In this way preparations of high activity were obtained, as follows: kidney, 6900 units/mg. N (Q_p 47 800); liver, 2800 units/mg. N (Q_p 19 400) and mammary gland, 4400 units/mg. N (Q_p 30 500). However, the yields were rather low, viz. between 2 and 5% of the initial material. The results are based on a single experiment in each case. The phosphatases from these tissues, like the similar enzyme from milk (Morton, 1953*c*), were found to be somewhat more labile than intestinal phosphatase. A summary of the results is given in Table 6.

DISCUSSION

Association of alkaline phosphatase with microsomal particles

While detailed distribution studies have not been carried out, it has been shown in this investigation that the specific activity of the microsome fraction,

both from calf intestine (Table 1) and cow mammary gland (Table 2) is very much greater than that of the mitochondrial fraction as well as of the whole tissue dispersion. This indicates that the alkaline phosphatase of these tissues is concentrated in the microsomes, and supports the earlier findings of Kabat (1941) with mouse kidney, Chantrenne (1947) with mouse liver, and Hers, Berthet, Berthet & Duve (1951) with both kidney and intestine of rabbit and guinea pig.

It has repeatedly been found that all the alkaline phosphatase of calf intestinal mucosa is removed by slow filtration of a tissue dispersion through a thick bed of Hyflo Supercel. This is not due to adsorption of the protein since the phosphatase, when in true solution following butanol treatment, is no longer retained by Hyflo Supercel. Thus it is concluded that all the alkaline phosphatase of intestinal mucosa is bound to particles. It is noteworthy that the recent studies of Novikoff, Podber, Ryan & Noe (1953) indicate that alkaline phosphatase of rat liver occurs partly in true solution and hence, in this respect, differs from the enzyme of kidney, mammary gland, milk and intestine.

The occurrence of microsomes in cow's milk suggests that some of these particles, like mitochondria, occur freely in cells (Morton, 1953*d*). However, it is likely that most of the heterogeneous microsome fraction isolated from tissue dispersions is derived by breakdown of other cellular structures. The phosphatase-rich brush border, intracellular membranes, and the so-called 'Golgi apparatus' probably contribute to the microsomes from intestinal mucosa. These particles, in contrast to microsomes from rat liver (Strittmatter & Ball, 1952), lack any red haemoproteins.

The effect of butanol on microsomal particles

During studies of the alkaline phosphatase in milk, it was found that this enzyme was firmly bound to lipoprotein particles ('milk microsomes') from which it could be quantitatively released into true solution by treatment with butanol (Morton, 1953*b*). The alkaline phosphatases of animal tissues may similarly be quantitatively released by treatment with butanol (Table 3). It is possible to use this treatment for assay of alkaline phosphatase in tissues, and the procedure has several advantages over use of whole tissue dispersions or autolytic extracts. The disadvantages of assaying alkaline phosphatase in tissue suspensions has been frequently noted (see Kay, 1932; Mathies, 1951). Moreover, as found by Mathies (1951) in the case of hog kidney, autolysis may not always liberate the enzyme from the particulate components. Further advantages of the use of butanol extracts for enzyme assay are: (1) Clear extracts are obtained, which retain their activity for long periods, and

therefore may be stored for assay at a convenient time. (2) Natural inhibitors, if present, may usually be removed by dialysis. (3) Manipulations are simplified, and truly representative samples may be accurately prepared. (4) The butanol extract may be used for estimation of the total lipid and phospholipid of the material, if this information is required. Nucleic acid phosphorus and protein phosphorus may then be readily estimated without interference from lipid phosphorus.

The release of the alkaline phosphatase appears to be dependent on the disruption of lipid-protein bonds. The resistance of this enzyme to tryptic digestion (Schmidt & Thannhauser, 1943; Abul-Fadl & King, 1949*b*) permits the use of proteolytic methods for this purpose. However, it is most undesirable to use such procedures for purifying proteins, since (1) further protein (or proteins) are added, which must later be removed; (2) protein degradation products are formed, ranging from large polypeptides to peptides and amino acids, thus considerably increasing fractionation difficulties; (3) the enzyme to be purified may react with these products in a specific manner; and (4) the enzyme itself may be modified by proteolysis.

In contrast to butanol, other organic solvents such as chloroform, toluene, ether and acetone do not release alkaline phosphatase into true solution from aqueous suspensions of microsomal particles. The excess butanol recovered after treatment of microsomes as previously described has been found to contain predominantly phospholipids, mostly of lecithin and cephalin types with some acetal phospholipid, as well as a non-phosphorylated compound which induces ketonuria in rats (Stewart & Young, 1952). Much of the butanol-soluble material is precipitated, partly in crystalline form, when the butanol extract is held at -15° for 24 hr.

Purification of alkaline phosphatases

Although the purification procedure has been described in detail only for the alkaline phosphatase of intestinal mucosa, experience with liver, kidney and mammary gland has shown that it may be used for purification of alkaline phosphatases of other animal tissues without major modification. The essential features of the purification method are: (1) Separation of the enzymically active particulate component as far as possible from soluble proteins and other material. (2) Release of the bound enzyme by treatment with butanol. (3) Fractionation with acetone (or acetone plus ether). (4) Selective denaturation of inert protein by treatment at pH 4.9 followed by treatment at pH 8.5. In the case of intestinal phosphatase, treatment at low pH should be combined with selective heat denaturation, viz. 48° , 2 min. (Table 4, stage 6). (5) Adsorption of inert protein on animal charcoal. In order to

obtain homogeneous preparations from tissues other than intestinal mucosa, it would be necessary to determine the most profitable sequence of these procedures.

The value of the early removal of soluble protein, followed by release of the enzyme by butanol treatment, is indicated by the sixty-fold purification gained at stage 3 (Table 4). However, the particles were obtained by repeated precipitation at pH 5.0. Undoubtedly the precipitate would contain protein, insoluble at this pH but soluble at pH 7.5, and mitochondria as well as microsomes. It is not surprising, therefore, that it was found in a small-scale trial that butanol treatment of washed microsomes, prepared by differential centrifuging at pH 7.2, gave a clear extract containing the alkaline phosphatase purified about 300-fold as compared with the whole 'homogenate'. The advent of centrifuges capable of handling relatively large volumes at high centrifugal forces has now made it possible to use such a procedure in preparative work, whereas this was not possible in 1949-50, when the work described here was completed. It is therefore strongly recommended that washed microsomes should be prepared by differential sedimentation as previously suggested (stage 2a), rather than by precipitation at pH 5.0. The remaining purification should then be carried forward as described.

During development of the purification method, a number of procedures were examined at each stage. In addition to fractionation with acetone and ether, low-temperature fractionation was examined using ethanol, *n*- and *iso*-propanol and dioxan, with and without addition of ether. None were as satisfactory as acetone, and some, such as dioxan, were found to be very toxic to the enzyme. Other adsorbents, such as calcium phosphate and alumina (C γ) gels, were found to be less effective than charcoal in removing inert protein. Adsorption of inert protein was found of much less value if used prior to stage 7 (Table 4). The enzyme could be readily adsorbed on calcium phosphate and alumina (C γ) gels, as well as on charcoal, and eluted with dilute salt solutions. However, if the enzyme were so treated after stage 7 (Table 4) the specific activity always declined. This suggests that there is some denaturation of the purified protein by spreading on the surface of the adsorbent. Hence adsorption and elution, such as used by Roche & Bouchilloux (1950), has been avoided in this purification procedure.

The concentrations of acetone given in the purification procedure (Table 4) should be used as a guide rather than as invariable limits, and it is recommended that small-scale trials should be carried out before treatment of the main bulk, in order to determine the most profitable fractionation

limits. This also applies to stage 7 (Table 4), since the amount of charcoal to use is somewhat dependent on the impurities present after stage 6. Should the enzyme be adsorbed by using excess charcoal, it may be largely recovered by eluting 3 times with 0.25 M sodium carbonate containing 0.5 M ammonium sulphate and adjusted to pH 8.0.

Schmidt & Thannhauser (1943) obtained from calf intestinal mucosa a final preparation of specific activity 26 000 units/mg. N, and Roche & Bouchilloux (1950) modified the procedure of Thoai, Roche & Roger (1947) to obtain from dog intestine a preparation of specific activity 28 000 units/mg. N. From electrophoretic data, they considered this to be 51.7% pure and so calculated that pure intestinal phosphatase should have a specific activity of 54 300 units/mg. N. It should be stressed that the high activities reported by Thoai *et al.* (1947) and Roche & Bouchilloux (1950) were obtained only after prolonged dialysis and reactivation of the enzyme with alanine. The best preparation obtained by Abul-Fadl, King, Roche & Thoai (1949), who also used dog intestinal mucosa but did not employ alanine re-activation, had a specific activity of 15 700 units/mg. N.

The preparation of specific activity 83 500 units/mg. N (Q_0 , 579 000), isolated from calf intestinal mucosa, is thus more active than any of the preparations described by other workers, and meets the criterion of purity, based on electrophoretic analyses, established by Roche & Bouchilloux (1950). Despite numerous attempts it has not been possible to obtain a more active enzyme. Homogeneity is also indicated by the absence of phosphodiesterase and pyrophosphatase activities, and of carbohydrate (Morton, 1952; cf. Schmidt & Thannhauser, 1943). Moreover, the nitrogen content (15.2%) and the ultraviolet absorption spectrum (Morton, 1952) are also consistent with the preparation being a purified protein. Because of the limited amount of the purified material finally obtained, no attempt was made to crystallize the intestinal alkaline phosphatase. However, in view of the fact that microcrystalline material was obtained by Schmidt & Thannhauser (1943) with a preparation of considerably lower activity, it seems probable that conditions for crystallization could now be developed.

The preparations obtained from ox kidney, liver and mammary gland cannot be considered as homogeneous enzymes, although they are much more active than those hitherto obtained from these tissues. The enzyme from ox kidney, for example, has a specific activity of 6900 units/mg. N, and thus is about twice as active as the most active material previously prepared from the same source (Abul-Fadl *et al.* 1949). Similarly, the enzyme of specific activity 4400 units/mg. N obtained from mammary

gland was somewhat more active than the preparation of specific activity 3800 units/mg. N obtained by Caputto & Marsal (1941). However, further purification of the enzymes from kidney, liver and mammary gland was not pursued. An essentially homogeneous preparation of this class of alkaline phosphatases, which are distinct from intestinal alkaline phosphatases (Morton, 1952), had already been obtained from milk (Morton, 1953c). It has a specific activity of 15 300 units/mg. N.

SUMMARY

1. The alkaline phosphatases of calf intestinal mucosa and of cow mammary gland are associated with microsomal particles.

2. By treatment with *n*-butanol at 35° the enzyme is readily obtained in true solution. On the basis of this effect a convenient method for assay of alkaline phosphatase of animal tissues is proposed.

3. The purification of alkaline phosphatase from calf intestinal mucosa is described. It involves butanol treatment of washed microsomes, followed by acetone precipitation, heating and charcoal adsorption.

4. The most active preparation had an activity (with β -glycerophosphate) of 83 500 units/mg. N (Q_p 579 000). This represents a purification of over 1000 times and thus is the most active phosphatase preparation as yet obtained.

5. The alkaline phosphatases of kidney, liver and mammary gland have been partially purified by similar methods.

I wish to thank Dr M. Dixon, F.R.S., for suggesting this problem, and for his valuable advice and encouragement throughout the investigation. I also wish to thank Dr V. Cosslett for granting me the facilities of the electron microscope at the Cavendish Laboratory, University of Cambridge, and Mr R. Horne for taking the electron micrographs. This work was carried out while holding a Travelling Scholarship

awarded by the Gowrie Scholarship Trust of Australia. My thanks are also due to the N.S.W. Milk Board for some financial assistance.

REFERENCES

- Abul-Fadl, M. A. M. & King, E. J. (1949*a*). *Biochem. J.* **44**, 435.
 Abul-Fadl, M. A. M. & King, E. J. (1949*b*). *Biochem. J.* **44**, 434.
 Abul-Fadl, M. A. M., King, E. J., Roche, J. & Thoai, N.-v. (1949). *Biochem. J.* **44**, 428.
 Albers, H. & Albers, E. (1935). *Hoppe-Seyl. Z.* **232**, 165, 189.
 Caputto, R. & Marsal, A. (1941). *Rev. Soc. argent. Biol.* **17**, 139.
 Chantrenne, H. (1947). *Biochim. biophys. Acta*, **1**, 437.
 Ehrensward, G. (1933). *Hoppe-Seyl. Z.* **217**, 274.
 Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
 Hers, H. G., Berthet, J., Berthet, L. & Duve, C. de (1951). *Bull. Soc. Chim. biol., Paris*, **33**, 21.
 Johnson, M. J. (1941). *J. biol. Chem.* **137**, 575.
 Kabat, E. A. (1941). *Science*, **93**, 43.
 Kay, H. D. (1932). *Physiol. Rev.* **12**, 384.
 Mathies, J. C. (1951). *Biochim. biophys. Acta*, **7**, 387.
 Morton, R. K. (1950). *Nature, Lond.*, **166**, 1092.
 Morton, R. K. (1952). Thesis submitted for Ph.D. degree, University of Cambridge.
 Morton, R. K. (1953*a*). *Nature, Lond.*, **172**, 65.
 Morton, R. K. (1953*b*). *Biochem. J.* **55**, 786.
 Morton, R. K. (1953*c*). *Biochem. J.* **55**, 795.
 Morton, R. K. (1953*d*). *Nature, Lond.*, **171**, 734.
 Morton, R. K. (1954). *Biochem. J.* **57**, 231.
 Novikoff, A. B., Podber, E., Ryan, J. & Noe, E. (1953). *J. Histochem. Cytochem.* **1**, 27.
 Roche, J. & Bouchilloux, S. (1950). *Bull. Soc. Chim. biol., Paris*, **32**, 732.
 Schmidt, G. & Thannhauser, S. J. (1943). *J. biol. Chem.* **149**, 369.
 Stewart, H. B. & Young, F. G. (1952). *Nature, Lond.*, **170**, 976.
 Strittmatter, C. F. & Ball, E. G. (1952). *Proc. nat. Acad. Sci., Wash.*, **38**, 19.
 Thoai, N.-v., Roche, J. & Roger, M. (1947). *Biochim. biophys. Acta*, **1**, 61.

Copper Complexes in Grassland Herbage

By C. F. MILLS*

Department of Physical Sciences, Wye College (University of London), Ashford, Kent

(Received 20 February 1954)

Copper deficiency disorders of cattle and sheep in Britain frequently exist in areas where the dietary copper intake of the grazing animal appears to be adequate. In areas where neonatal ataxia of lambs (Swayback) is found (a condition which can be prevented by copper therapy) herbage copper values

are appreciably higher than those of areas in Australasia where the apparently identical syndrome, enzootic ataxia, is common. Thus Bennetts & Beck (1942) reported outbreaks of enzootic ataxia in Western Australia on pastures ranging in copper content from 1.1 to 6.1 p.p.m. (on dry matter) with a mean of 3.3 p.p.m. Cunningham (1944) described a copper deficiency in New Zealand where sheep

* Present address: Rowett Research Institute, Bucksburn, Aberdeenshire.