- Jermyn, M. A. (1952). *Aust. J. Sci. Res. B*, **5**, 433.
 Karunairatnam, M. C. & Levvy, G. A. (1951). *Biochem. J.* **49**, 210.
 Lederberg, J. (1950). *J. Bact.* **60**, 381.
 Levvy, G. A. (1952). *Biochem. J.* **52**, 464.
 Levvy, G. A. & Marsh, C. A. (1954). *Science*, **119**, 337.
 Lineweaver, H. & Burk, D. (1934). *J. Amer. chem. Soc.* **56**, 658.
 McIlvaine, T. C. (1921). *J. biol. Chem.* **49**, 183.
 Markham, R. (1942). *Biochem. J.* **36**, 790.
 Montgomery, E. M., Richtmyer, N. K. & Hudson, C. S. (1942). *J. Amer. chem. Soc.* **64**, 190.
 Nef, J. U. (1914). *Liebigs Ann.* **403**, 329.
 Niwa, K. (1943). *Science (Japan)*, **13**, 171, 289. Cited in *Chem. Abstr.* (1951), **45**, 10271.
 Rehorst, K. (1928). *Ber. dtsch. chem. Ges.* **61 B**, 163.
 Seidman, M. & Link, K. P. (1950). *J. Amer. chem. Soc.* **72**, 4324.

The Action Pattern of Potato Phosphorylase

BY W. J. WHELAN AND J. M. BAILEY*

Department of Chemistry, University College of North Wales, Bangor

(Received 15 April 1954)

The action of phosphorylase is expressed by the equation

The enzyme belongs to the group of carbohydrate-transferring enzymes, the function of which is the transfer of sugar residues from the donor substrate (in this case, glucose 1-phosphate or amylose), either singly or severally to an acceptor substrate containing a hydroxyl group (primer or orthophosphate).

The present investigation is concerned with two problems which are common to these transferring enzymes. The main problem is the elucidation of the mode of action of the enzyme, but this is bound up with the problem of the function of the maltodextrins (maltosaccharides) in the forward (synthetic) reaction. Confining the attention to this synthetic reaction two patterns of action may be envisaged. These are described as 'single-chain' and 'multi-chain'. In the former, a molecule of phosphorylase continuously increases the length of a single primer chain until the molecule becomes immune to further enzyme action (e.g. by precipitating from solution), before transferring its activity to a second molecule of primer. In the multi-chain type of action random synthesis occurs and all the primer chains grow at approximately equal rates. This seems the more probable mechanism and has generally been assumed in the past (e.g. see Swanson, 1948), Bernfeld & Meutémédian (1948), however, favoured single-chain action. If it could be shown that multi-chain synthesis occurred, then by the use of primer molecules of known size, the synthesis of linear chain molecules of any predetermined average length could be accomplished.

* Present address: Department of Chemistry, Iowa State College, Ames, Iowa, U.S.A.

These substances would provide the material for studies of the variation in physical and chemical properties in a homologous series of polymers and of the action patterns of other enzymes. In this paper it is established that synthesis by phosphorylase follows the multi-chain pattern, and synthetic amyloses have been prepared for use in studies of variation in iodine stain with chain length (Whelan & Bailey, to be published) and of the action patterns of Q-enzyme (Peat, Whelan & Bailey, 1953) and β -amylase (Whelan & Bailey, to be published). Through the initiative of Mould & Synge (1954) they have also been used in developing a new method of fractionating large molecules.

The second problem required a study of the behaviour of maltodextrins as primers in the synthetic reaction. Hanes (1940) first showed that the synthetic reaction did not proceed in the absence of starch, dextrins or maltose. Green & Stumpf (1942) failed to detect priming action with maltose but confirmed Hanes's other observations. Weibull & Tiselius (1945) hydrolysed a starch dextrin (produced amylolytically) with acid and fractionated the products on charcoal using aqueous ephedrine as eluting agent. They found that the product of lowest molecular weight to exhibit priming activity was the trisaccharide, maltotriose. We have re-investigated the priming activity of the maltodextrins using material prepared as described by Whelan, Bailey & Roberts (1953). It will be shown that the choice of primer can exercise a profound influence on the polydispersity of the synthetic amylose; if maltotriose had been the only primer used in the study of action pattern the conclusion might well have been that single-chain synthesis was taking place. Maltotriose is unique among the maltodextrins in respect of the action on it of starch-metabolizing enzymes. It is very slowly attacked by β -amylase (Whelan *et al.* 1953)

and by crude salivary α -amylase (Whelan & Roberts, 1953) and was found to be a relatively very inefficient primer in amylose synthesis (Bailey, Whelan & Peat, 1950). This last property has been the subject of a detailed study which is also reported below.

Some aspects of this work have been summarized by Bailey *et al.* (1950) and Bailey & Whelan (1952).

MATERIALS AND METHODS

α -D-Glucose 1-phosphate. The primer-free dipotassium salt was prepared as by McCready & Hassid (1944), and used as a 0.1 M aqueous solution, adjusted to pH 7.0 with H_2SO_4 .

Maltodextrins. These were prepared from potato amylose and the concentrations of their aqueous solutions determined as described by Whelan *et al.* (1953).

Achroic primer. Potato amylose (100 mg.) was dissolved in N-NaOH (10 ml.) and acidified with $\text{N-H}_2\text{SO}_4$ (12 ml.) giving a solution 0.1 N with respect to acid. This was heated for 2 hr. at 100° , when the iodine stain was found to have disappeared. The solution was then cooled, neutralized (NaOH), diluted to 100 ml. and stored under toluene.

Buffers. Unless otherwise mentioned, all buffers were 0.2 M in concentration and were constituted as follows: citrate (citric acid, NaOH ; pH 7.0), phosphate (KH_2PO_4 - Na_2HPO_4 ; pH 6.7) and acetate (acetic acid- NaOH ; pH 4.8 and 7.0).

Iodine solution. All iodine staining was carried out with 0.2% (w/v) I_2 in 2% (w/v) KI .

α -Amylase. This was prepared from human saliva as by Whelan & Roberts (1953). The method is essentially that of Meyer, Fischer, Staub & Bernfeld (1948).

β -Amylase. A preparation was obtained from soya beans as described by Peat, Pirt & Whelan (1952*b*) and a specimen of the crystalline enzyme from sweet potato was kindly provided by Dr A. K. Balls. Both preparations exhibited the same properties (Peat, Pirt & Whelan, 1952*a, b*) and were used interchangeably. The enzyme activity was determined as by Hobson, Whelan & Peat (1950).

Phosphorylase. The enzyme was obtained from potatoes (3.5 kg.) following the method of Barker, Bourne, Wilkinson & Peat (1950) as far as the elution of the lead complex. The protein was then fractionated with 50% (w/v) $(\text{NH}_4)_2\text{SO}_4$ solution (pH 7.0; 0°) by rejecting the fraction precipitating at 19% (w/v) $(\text{NH}_4)_2\text{SO}_4$ and precipitating the phosphorylase at 35% (w/v). This was redissolved in water and the procedure repeated by rejecting the further precipitates obtained at 21, 23 and 25% (w/v). The enzyme was precipitated with 35% (w/v) $(\text{NH}_4)_2\text{SO}_4$ after each of these steps. Centrifuging was carried out at room temperature. The final precipitate was dissolved in 0.5 M sodium citrate buffer (pH 6.0; 75 ml.) and freeze-dried. In a typical preparation the resulting powder (10 g.) had an activity of 7.5 units/g. (see below).

Determination of inorganic phosphate. The method of Allen (1940) was used. In this method the molybdenum blue colour is developed with a mixture of 60% (w/v) HClO_4 , 1% (w/v) amidol (2,4-diaminophenol hydrochloride) in 20% (w/v) NaHSO_4 and 8.3% (w/v) ammonium molybdate. Glucose 1-phosphate was frequently present in solutions under determination. An appreciable error resulted from hydrolysis of the phosphate ester when in contact with these reagents. An attempt was made to eliminate this

error by making the measurements at exactly the same time after mixing the reagents, but failed when it was shown that the rate of hydrolysis of the ester was influenced by the amount of inorganic phosphate present. The problem was resolved by the accidental discovery that hydrolysis was very much reduced by halving the specified quantity of amidol reagent. The quantities of reagents now used were, HClO_4 (2 ml.), amidol (1 ml.) and molybdate (1 ml.) in a total volume of 25 ml. By this means the average increase in intensity of the blue colour between 9 and 30 min. after mixing the reagents was reduced from 4.0 to 0.6 scale units on the 'EEL' colorimeter (Evans Electroselenium Ltd.) using 1.3 cm. diam. cells and Ilford filter no. 608 (max. transmission, 680 m μ). The average equivalent of 0.1 mg. of phosphorus in colorimeter scale units is 65.

Determination of glucose 1-phosphate. A modification of the method of Hanes (1940) was used. This employs acidic hydrolysis of the ester by heating its solution for 7 min. at 100° with the 60% (w/v) HClO_4 (2 ml.) in a total volume of 10 ml., cooling and adding amidol and molybdate as above and finally diluting to 25 ml. with water. Inorganic phosphate present before hydrolysis is also determined and the difference between the determinations is the measure of ester phosphate. When the amount of inorganic phosphate is so large as to make the difference determination inaccurate this ion is first removed by precipitation with magnesia mixture (MgCl_2 , 110 g. + NH_4Cl , 140 g. + 1.65 N- NH_4OH , 2 l.; stand for 3 days and filter). Control experiments with solutions varying from 0.5 to 3.5 mm with respect to glucose 1-phosphate and up to 50 mm in inorganic phosphate showed that losses as high as 24% of ester phosphate occurred by co-precipitation. No loss occurred if the inorganic phosphate content was kept below 5 mm, by dilution if necessary, before adding magnesia reagent (0.25 vol.). Even after precipitation with magnesia reagent it was still necessary to determine inorganic phosphate present before hydrolysis, since traces of inorganic phosphate remained unprecipitated.

Determination of phosphorylase activity. Two digests containing citrate buffer (1 ml.), 1% soluble starch (0.5 ml.), enzyme solution (0.5 ml.) and water (0.5 ml.) were prepared and brought to 35° in a thermostat. To one digest was added 0.1 M glucose 1-phosphate (1 ml.), preheated to 35° . After 12 min. a portion (1 ml.) was removed and added to 60% (w/v) HClO_4 (2 ml.) in water (approx. 8 ml.) for determination of inorganic phosphate. At the same time the ester phosphate solution (1 ml.) was added to the second digest with shaking and a portion (1 ml.) immediately withdrawn for determination of the initial inorganic phosphate content of the digest. The unit of activity of the enzyme is defined as that amount which will cause the liberation of 0.1 mg. P in the above digest during 3 min.

Measurement of light absorption. Light-absorption curves of iodine-stained polysaccharides were measured in the Unicam SP. 500 spectrophotometer, using 1 cm. cells.

Inhibition of enzymic impurities. In some experiments traces of amylolytic enzymes, Q-enzyme and phosphoglucosmutase present in the phosphorylase preparation were inhibited by the inclusion in the digest of HgCl_2 . Phosphatase, attacking glucose 1-phosphate, was inhibited by ammonium molybdate (Bailey, Thomas & Whelan, 1951). Control experiments showed that these inhibitors did not interfere with phosphate determination nor did they influence the iodine-staining properties of synthetic amyloses.

Reducing power. This was determined with the Somogyi (1945) copper reagent.

Preparation of synthetic amyloses. Details are given of a typical preparation.

The digest contained maltohexaose (3.2×10^{-5} moles), glucose 1-phosphate (60 ml.), citrate buffer (4 ml.), 8.3% (w/v) ammonium molybdate (2 ml.), 0.153 mm-HgCl₂ (0.6 ml.) and phosphorylase (200 mg.) in a total volume of 100 ml. The enzyme solution was added last and the solution was immediately diluted to the mark, mixed and a 0.3 ml. portion removed for inorganic phosphate determination, the enzyme being inactivated at once by running the solution into the perchloric acid. During incubation at 35°, further portions were removed for measurement of inorganic phosphate and the phosphate/time curve was extrapolated to the point at which calculation had shown that the average chain length of the synthetic amylose would be 60 glucose units. At this time (2 hr. 35 min. incubation) a 55 ml. portion was removed and immediately heated to 100° in a boiling-water bath, heating being continued for 3 min. The inorganic phosphate content of the cooled solution was determined and used to calculate the chain length of the amylose. The quantity of phosphate liberated gives the number of glucose units added to the primer molecules, initially six units in length. Division of the molar amount of added glucose by the known molar concentration of primer gives the average number of glucose units added to each primer molecule. In this particular experiment 55 units were added. The average chain length of the amylose was therefore 61. The remainder of the digest was incubated for a further 2 hr. and, after inactivating the enzyme, the average chain length was found to be 81.5. Both solutions were dialysed for 7 days against running tap water, centrifuged, and the supernatant solutions freeze-dried.

RESULTS AND DISCUSSION

The priming action of maltodextrins

Products of the partial acidic hydrolysis of amylose, i.e. glucose, maltose, maltotriose, etc., were separated by charcoal chromatography and tested, in equimolar concentration, for priming activity in the synthetic reaction. The results shown in Fig. 1 confirm previous observations (Green & Stumpf, 1942; Weibull & Tiselius, 1945) that glucose and maltose are devoid of priming activity. Maltotriose and the higher dextrins exhibit activity. Experiments were next performed to examine the relationship between the rate of phosphate liberation and the concentration of the maltodextrin primers. It became evident that the concentration of primer used in the above experiment was too high to indicate the activity of maltotriose relative to the activities of the higher dextrins. It was found that a linear relationship between these two functions existed only up to a concentration of 2×10^{-5} M for the tetra-, penta- and hexa-saccharides and up to 6×10^{-4} M for maltotriose under the experimental conditions given in Table 1 which shows the molar equivalent priming activities and demonstrates that under these conditions maltotriose has only

9% of the activity of the tetraose. Malto-pentaose and -hexaose show successively higher activities.

It was of interest to compare the activities of the maltodextrins after degradation by β -amylase in order to determine whether inferences drawn from a study of their behaviour on β -amylolysis (Whelan *et al.* 1953) could be substantiated. These earlier experiments had shown that a small amount of β -amylase, while without action on maltotriose, rapidly converted malto-tetraose, -pentaose and -hexaose into, respectively, maltose (2 mol. prop.),

Fig. 1. Priming activity of glucose and maltodextrins obtained by charcoal chromatography. Each digest except the control contained glucose or maltodextrin (1 μ mole), citrate buffer (1 ml.), glucose 1-phosphate (1 ml.), phosphorylase (50 mg.) and water to 3.2 ml. During incubation at 35°, portions (0.5 ml.) were withdrawn for determination of liberated phosphate.

Table 1. *Priming activities of equimolar amounts of maltodextrins before and after β -amylolysis*

Maltodextrin (0.62 μ mole maltotriose, 0.06 μ mole malto-tetraose, -pentaose and -hexaose) was incorporated in digests containing glucose 1-phosphate (0.2 ml.), phosphorylase (20 mg.), citrate buffer (1 ml.) and water to 3 ml. After incubation at 35° for 75 min. a portion (1 ml.) was withdrawn for determination of liberated phosphate. The same quantities of dextrin, mixed with acetate buffer (pH 4.8; 0.1 ml.), were treated with crystalline β -amylase (2.6 units) in a total volume of 1.3 ml. After 2.5 hr. at 35° the enzyme was inactivated by being heated at 100° and the above quantities of ester phosphate, buffer and enzyme were added for determination of residual priming activity. Control experiments were performed to determine phosphate liberation in the absence of added primer.

Primer	Phosphate liberated (as μ g. P/ μ mole primer)	
	Before β -amylase action	After β -amylase action
Maltotriose	35	34
Maltotetraose	390	0
Maltopentaose	425	34
Maltohexaose	497	14

maltose and maltotriose (1 mol. prop. of each) and maltose (3 mol. prop.). High concentrations of β -amylase slowly convert maltotriose into maltose and glucose (Whelan *et al.* 1953; Whelan & Roberts, 1953). The priming activities of the products of the action of a small amount of β -amylase on the maltodextrins also confirm these conclusions (Table 1). Maltotriose activity was unaffected by β -amylolysis, that of the tetraose disappeared, pentaoase activity fell to the expected triose level, and hexaoase activity to 2.8% of its original level, due either to incomplete degradation to maltose or to the presence of impurities of maltopentaose and/or maltoheptaose. A further argument favouring this interpretation of β -amylase action is provided in the accompanying paper (Whelan & Roberts, 1954).

Maltotriose. Because of the relatively very low priming activity of this preparation a number of experiments were performed to test the possibility that this activity was not an intrinsic property of the triose but was due to an impurity either in the sugar or in the phosphorylase preparation. The need for such further tests was emphasized by the reasoning that the true priming activity of the triose relative to the higher dextrins is certainly much lower than that depicted in Table 1. Ideally, priming activity should be measured in terms of the rate of addition of a single glucose unit to the primer molecule. In practice, this would require greater amounts of primer than were available. However, where the rates of successive addition of several glucose units do not differ appreciably, the measurement of the overall rate of addition of many glucose units, as was being done here, provides a reasonably accurate measure of the true priming activity. In the case of an inefficient primer this measure is not accurate since the liberation of inorganic phosphate is greatly accelerated after the first addition of a glucose unit to form an efficient primer. This point is illustrated by the curve depicting maltotriose priming activity in Fig. 1. These results show that rate of reaction expressed in terms of a single measurement (the method used in Table 1) is dependent on the time of measurement and increases with time. This does not happen in the tests with the higher dextrins (Fig. 1). From a knowledge of the variation in iodine-staining properties of synthetic amyloses with chain length (Whelan & Bailey, to be published) it was found that when using maltotriose as a primer the products of synthesis accumulated in sufficient concentration to become detectable by iodine staining only when their average chain length was at least 100 units. A rough calculation based on this fact suggested that the true relative priming activities of the triose and tetraose might be in the ratio of only 1:10⁴.

The following experiments, of which only the last is reported in detail (Table 2), led us to conclude

that an impurity in the maltotriose preparation, the most probable being maltotetraose, was not responsible for the priming activity.

The first series of experiments involved the action of the α - and β -amylases on the maltotriose. It had already been shown that after treatment with a small amount of β -amylase maltopentaose possessed a residual priming activity, equal on a molar basis to that of the maltotriose, while the triose itself showed undiminished activity (Table 1). These results were suggestive of the absence of a maltotetraose impurity, since this substance loses all activity when subjected to the same treatment (Table 1). Nevertheless, it could not be overlooked that maltotriose, itself very slowly attacked by β -amylase, might act as a competitive inhibitor of β -amylase and largely prevent the hydrolysis of a trace impurity. When, however, the rates of β -amylolysis of malto-tetraose or -pentaose in the presence and absence of an equimolar amount of maltotriose were compared, no effect of maltotriose on the rates of β -amylolysis of the higher dextrins was observed.

Since a small amount of β -amylase had failed to diminish the priming activity of the maltotriose preparation, an amount of enzyme sufficient to bring about the complete hydrolysis of the triose itself was next used and priming activity was measured at stages during the reaction. No initial rapid fall in activity, which would have indicated the removal of an impurity by the β -amylase, was observed; at all stages during the hydrolysis the loss of priming activity was proportional to the amount of maltotriose hydrolysed. In view of the demonstration that the addition of maltotriose had no influence on the rapid β -amylolysis of the tetraose (see above), this new result was a clear indication of the intrinsic priming activity of maltotriose. Furthermore, prolonged treatment of the maltotriose specimen with salivary α -amylase caused no detectable diminution in its priming activity. This α -amylase differs from β -amylase in having no action on maltotriose but, like the latter enzyme, converts maltotetraose into maltose (Whelan & Roberts, 1953).

When it was found that refractionation of the triose specimen on charcoal did not diminish its priming activity, a final test for the presence of a priming impurity was applied. Two digests were prepared containing phosphorylase, glucose 1-phosphate together with, in one case, maltotriose and in the other, the equivalent amount of maltotetraose in terms of priming activity. Since the 'priming equivalent' of maltotetraose was only 9% (molar basis) of that of maltotriose (when measured as in Table 1), it is to be expected that a greater weight of amylose would be synthesized with the maltotriose primer than with the tetraose, assuming

that the chains do not grow to infinite length but that synthesis is terminated by retrogradation. The results in Table 2 show that this in fact occurred. It is further to be expected that if maltotriose is a primer, albeit a poor one, $\lambda_{\max}$ of the iodine-stained products would change very little throughout the reaction. This follows because the product of addition of one glucose residue, i.e. maltotetraose, is so much more efficient as a primer that, once formed, the tetrasaccharide primer becomes considerably extended in length during the time interval before another maltotriose molecule is involved. Experiment confirms this view; the products are blue staining throughout the reaction and $\lambda_{\max}$ changes very little, viz. from an initial value of 620 m μ . to a final value of 640 m μ . (It has been established that maltodextrin chains become blue-staining when longer than about 45 units; Whelan & Bailey, to be published).

In contrast, when maltotetraose was used as primer there occurred the normal sequence of changes in iodine stain which, as will be shown in a later section, is a criterion of multi-chain synthesis, and this despite the fact that the much smaller amount of tetraose used made difficult the recognition of the early products of synthesis.

Bernfeld & Meutémédian (1948) claimed that only blue-staining products could be detected during the phosphorylase synthesis of amylose. It can, however, be inferred that these authors were using either a preparation in which maltotriose was the only primer or, more probably, one so deficient in primer that only the blue-staining products, which also have the greatest intensity of stain per unit weight, could be detected.

The work now reported was carried out in 1950-1, at which time the existence of D-enzyme in the potato was not suspected. This enzyme is able to

effect a disproportionation of maltodextrins, e.g. maltotriose into a mixture containing glucose, unchanged maltotriose, maltotetraose, -pentaose, etc. (Peat, Whelan & Rees, 1953). Clearly, the presence of this enzyme could seriously affect the course of amylose synthesis from glucose 1-phosphate and maltodextrin primer and, in particular, contamination of the phosphorylase preparation might have been responsible for the apparent priming activity of the maltotriose. None of the experiments mentioned above could have revealed such action since they were tests for the purity of the substrate and not of the enzyme preparation. D-Enzyme appears not to function by a phosphorylating mechanism so that a test for its presence could be made by incubation of the phosphorylase with maltotriose before the addition of glucose 1-phosphate. It was thus shown that synthesis proceeded at exactly the same initial slow rate as when there was no pre-incubation (Table 3).

All the above experiments lead to the conclusion that maltotriose is a primer in the synthetic reaction and confirm our original statement (Bailey *et al.* 1950). Its low activity relative to that of the higher dextrins has since been confirmed by French & Wild (1953) using an ingenious paper-chromatographic method. Weibull & Tiselius (1945) reported, on the other hand, that maltotriose, -tetraose, -pentaose and -hexaose exhibited approximately equal priming activity, but it seems possible either that the maltotriose was contaminated with higher dextrins or that the activity was not being measured with the correctly proportioned amounts of dextrin (cf. Fig. 1 and Table 1).

Action pattern of phosphorylase

Three independent methods were used in the examination of the action pattern of the enzyme.

Table 2. *Amylose synthesis using equal priming activities of maltotriose and maltotetraose*

Each digest contained glucose 1-phosphate (10 ml.), acetate buffer (pH 7.0; 5 ml.) 0.153 mm-HgCl₂ (1 ml.), 8% ammonium molybdate (0.55 ml.), phosphorylase (300 mg.); one digest contained maltotriose (0.93 mg.) and the other, maltotetraose (0.126 mg.), the total volume being 25 ml. During incubation at 35° portions (0.5 ml. each) were withdrawn for measurement of phosphate liberation and $\lambda_{\max}$ of the iodine-stained solution as in Fig. 3. Chain length was calculated as in the Methods section.

Maltotriose primer			Maltotetraose primer		
Chain length	Iodine stain		Chain length	Iodine stain	
	$\lambda_{\max}$	Colour		$\lambda_{\max}$	Colour
8.3	620	Green	6.2	—	Yellow
12	620	Green	10	—	Yellow
19	620	Blue-green	15	500	Red
25	—		21	520	Red
36	—		30	535	Red-purple
50	—		36	555	Purple
97	635		44	570	Blue
195	640		55	585	Blue-green
			101	618	
			164	620	

First, the kinetics of the reaction were studied in order to determine the relation between rate of amylose synthesis and primer concentration; secondly, the values of $\lambda_{\max}$ of iodine-stained amyloses were measured at stages during polysaccharide synthesis from maltotetraose primer; and thirdly, synthetic amyloses of calculated average chain length were prepared from maltohexaose primer and examined by Mould & Synge (1954) using the method of electrokinetic ultrafiltration.

Effects of variation in primer and enzyme concentrations. The rate of synthesis of amylose was measured in digests in which only the primer concentration was varied. This is the procedure followed in determining the Michaelis constant for the phosphorylase-primer complex. A linear relationship is obtained when the reciprocals of reaction velocity and primer concentration are plotted against each other (Fig. 2a). The same effect is shown in Fig. 2b where, at constant concentration of glucose 1-phosphate, the amounts of phosphorylase and primer were kept constant but the concentrations were varied by dilution. The linear relationship between the reciprocal of the reaction velocity and the square of the digest volume (Fig. 2b) indicates that the reaction is bimolecular with respect to enzyme and primer. Both results indicate that the primer behaves as a normal substrate in the reaction, and are consistent with a multi-chain pattern of enzyme action.

Light absorption of iodine-stained synthetic amyloses. The course of amylose synthesis from maltotetraose by the agency of phosphorylase was followed by plotting the light-absorption curves of

the products stained with iodine at intervals during digestion (Fig. 3). The visual changes in colour are from brown, through red and purple to a pure blue (cf. Swanson, 1948). Since the reverse changes in colour occur when amylose is randomly fragmented by acid or by α -amylase it is clear that this pattern of absorption curves is to be associated with the regular change in length of all primer molecules. The spectral pattern of single-chain synthesis would be very different since only the final product would contribute to the light absorption. In consequence $\lambda_{\max}$ should remain constant and independent of the amount of glucose 1-phosphate consumed. The single-chain pattern was simulated, for reasons already discussed, when maltotriose was used as primer (Table 2).

It was important to determine whether, as in synthesis, phosphorolytic degradation of amylose also proceeded as a multi-chain action. This was examined by reversing the course of the synthetic reaction shown in Fig. 3 by adding an excess of inorganic phosphate to the digest. The pattern of absorption curves obtained was qualitatively the same as for synthesis but at corresponding stages in

Table 3. *Priming activity of maltotriose after pre-incubation with phosphorylase*

Three digests were prepared containing citrate buffer (1 ml.), 8.3% ammonium molybdate (0.1 ml.), 0.153 M-HgCl₂ (0.1 ml.), phosphorylase (30 mg.) and maltotriose (1 mg.) in 2.8 ml. After 0, 2 and 4 hr. respectively, at 35°, glucose 1-phosphate (1.2 ml.) was added to each digest and portions (0.5 ml.) were removed for determination of liberated phosphate. Incubation was at 35°. A control digest was used to determine phosphate liberation in the absence of added primer, and further control digestions showed that phosphorylase activity did not begin to decay appreciably until after 8 hr. at 35°.

Time of incubation (hr.)	Time of pre-incubation (hr.)		
	0	2	4
	Phosphate liberated (colorimeter units)		
1	2.7	3.3	2.8
2	8.6	8.7	7.8
4	15.9	14.9	12.0
6	25.4	23.5	15.6
8	34.8	27.5	22.0

Fig. 2. (a) Michaelis-Menten relationship for the primer-phosphorylase system. Digests contained achroic primer (0, 0.1, 0.2, 0.5, 1.0 and 2.0 mg.), phosphorylase (100 mg.), glucose 1-phosphate (2 ml.), acetate buffer (pH 7.0; 1 ml.) and water to 6 ml. After incubation for 3 hr. at 20° portions (1 ml.) were withdrawn for measurement of phosphate liberation. Rate is expressed as mg. phosphorus liberated/min. in the whole digest and concentration as mg. primer in the digest. (b) Showing that at constant initial glucose 1-phosphate concentration, synthesis is bimolecular with respect to enzyme and primer. Digests contained achroic primer (1 mg.), phosphorylase (50 mg.), acetate buffer (pH 7.0; 2, 3, 4 and 6 ml.), glucose 1-phosphate (2, 3, 4 and 6 ml.) and water to 10, 15, 20 and 30 ml. respectively. During incubation at 35°, portions (0.2, 0.3, 0.4 and 0.6 ml. respectively) were withdrawn for measurement of liberated phosphate. The rate of liberation was uniform in each digest during the first 60 min. Rate is expressed as mg. phosphorus liberated/min. in the whole digest.

synthesis and degradation $\lambda_{\max}$ was higher in the latter reaction (Fig. 4). Phosphorolysis therefore differs from synthesis in giving an appearance of some degree of single-chain action. This behaviour could be shown however to be in accordance with the predictable characteristics of completely multi-chain action. During multi-chain synthesis, even from a homogeneous primer, the synthetic amylose becomes polydisperse, and increasingly so. It can be adduced on theoretical grounds that polydispersity of the product increases also when the

Fig. 3. Light-absorption curves of iodine-stained products during amylose synthesis from maltotetraose primer. The digest contained maltotetraose (10 mg.), glucose 1-phosphate (10 ml.), acetate buffer (pH 7.0; 10 ml.), phosphorylase (300 mg.) and water to 100 ml. At intervals during incubation at 35° portions (0.5 ml.) were stained with iodine (0.5 ml.) in a total volume of 25 ml. and their light absorption determined; further portions (0.5 ml. each) were withdrawn for measurement of liberated phosphate.

Fig. 4. Variation in $\lambda_{\max}$ of iodine-stained amylose during synthesis and degradation. Values for synthesis (●) obtained from the experiment of Fig. 3. After incubation of this digest for 250 min. a portion (88.5 ml.) was removed and mixed with phosphate buffer (20 ml.). During further incubation at 35° portions (0.61 ml.) were removed for iodine staining as in Fig. 3 and further portions (2 ml.) for measurement of glucose 1-phosphate as in Methods section. Results represented as ○.

reaction is reversed by the addition of inorganic phosphate to the digest. It will diminish as the minimum length of priming chain is approached. Only when this minimum is reached are all the chain molecules of the same length, a length (three glucose units, see above) which is well below the chain length requisite for iodine staining, estimated to be *ca.* 12 units. From a study, to be published later, of the relationship between chain length and $\lambda_{\max}$ of the iodine stain it has emerged that, provided the average chain length is less than 80 units, the $\lambda_{\max}$ values of amyloses of the same average length increase with increasing degree of polydispersity. It is known that the average chain lengths in the experiments of Fig. 4 did not exceed 80 units. It therefore follows that at corresponding values of chain length in the synthesis and phosphorolysis of amylose, the $\lambda_{\max}$ value for phosphorolysis will be the greater, because of the higher degree of polydispersity. This is precisely the effect which was observed.

A direct implication of the single-chain hypothesis is that at any time during amylose synthesis the iodine-staining material will consist of the final product, together with the relatively negligible amount of material undergoing extension of chain. In multi-chain action the properties of the product will depend on the extent to which synthesis has proceeded. Thus it follows that if in a series of syntheses the primer concentration is varied, but the reaction is arrested at stages when the amount of glucose 1-phosphate consumed in each case is the same, then the iodine-stained products will have the same or different values of $\lambda_{\max}$, according to whether single- or multi-chain action is operative. To this statement must be added the proviso that synthesis must not be allowed to proceed to the point at which the value of $\lambda_{\max}$ has become independent of chain length (Whelan & Bailey, forthcoming publication). The results of an experiment of this type (Fig. 5) are clearly in favour of multi-chain synthesis and the relative values of $\lambda_{\max}$ accord with the expectation that the greater the amount of primer present, the shorter is the average chain length of the product at stages where an equal number of glucose units has been distributed among the primer molecules. This effect has previously been noticed by Sumner, Somers & Sisler (1944).

Electrokinetic ultrafiltration of synthetic amyloses. The foregoing experiments provide strong circumstantial evidence that the enzyme action pattern is multi-chain. It was realized, however, that none of this evidence excluded the possibility that the action might not be entirely multi-chain but might also show single-chain characteristics. Such behaviour has been demonstrated in the case of β -amylase and was detected by the finding that the proportion

of each type of action could be varied by changing the temperature of incubation (French, Knapp & Pazur, 1950; Whelan & Bailey, to be published). Two series of experiments showed that similar considerations did not apply to the action of potato phosphorylase. First, the action pattern was shown not to vary with temperature (Whelan & Bailey, to be published). Secondly, a direct examination of the range of DP (degree of polymerization) in the synthetic amyloses was made by Mould & Synge (1954). (In the case of linear molecules DP is, of course, synonymous with chain length when both are expressed in terms of polymer units.) Before the latter results are considered it is profitable to consider in some detail the factors which may influence the course of multi-chain synthesis.

It is obvious that in any polymer synthesized by the endwise apposition of polymer units to priming molecules the range of DP of the chain molecules must increase as the average molecular weight is increased. Reference is being made here and throughout the paper to the absolute range of DP. As pointed out by Flory (1940) the range decreases with increase in DP if expressed as a percentage of mean DP of the preparation (see Table 4). The increase in absolute range of DP occurs even if the

addition of the units is completely random and statistical calculation of this range can be made at any stage during such synthesis (Table 4). However, a number of factors may operate to increase the range. These are as follows:

(1) Table 1 shows that the priming efficiency of maltodextrins increases with molecular size in the series from maltotriose to maltohexaose and it cannot be assumed that the activity of maltohexaose is the maximum possible. The effect on the range of DP of the synthetic amylose of using a primer of submaximal activity has been mentioned above in comparing the products synthesized when malto-triose and -tetraose are used as primers. This same effect, although much less serious than in the case of maltotriose, must influence the range of DP in the synthetic amyloses which we prepared from maltohexaose for examination by Mould & Synge (1954). In the absence of knowledge of the priming activity of maltodextrins of greater molecular weight than the hexaose, no estimate of the magnitude of this effect can be made. It is possible, however, that priming activity passes through a maximum and begins to diminish with subsequent increase in chain length. Such an effect would operate to decrease the range of DP.

(2) The turnover number of potato phosphorylase has not been determined; that of muscle phosphorylase is 4×10^4 linkages formed per minute in the synthetic reaction at pH 6.7 and 30° (Cori, Cori & Green, 1943). If the potato enzyme has the same high rate of action the possibility must be considered that once contact has been established between enzyme and primer the latter may not be able to diffuse away after the apposition of one glucose unit but may undergo further chain lengthening. Such multiple attack (a term introduced by French & Wild, 1953) must occur even though synthesis may be completely random and would be favoured by a high rate of enzyme action. Again, no information on this aspect is available except that, as mentioned above, the effect is certainly operative during β -amylolysis and varies with temperature (the turnover number of β -amylase is 2.5×10^5 ; England & Singer, 1950; England, Sorof & Singer, 1951), but the pattern of phosphorylase action on amylose molecules up to an average chain length of 49 units does not vary with temperature (Whelan & Bailey, to be published).

(3) Naturally occurring amyloses show a marked tendency to retrograde from solution and this also begins to occur during the synthesis of amylose when the average chain length becomes greater than about 200 units. The process is gradual and does not appear to terminate the length of every chain at the same limit. The extent of retrogradation is to some degree controlled by the amylose concentration and by the other solutes present. It may not always be

Fig. 5. Light-absorption curves of iodine stains of equal weights of amylose, synthesized from varying amounts of primer. The digests, corresponding to curves A, B, C and D, contained glucose 1-phosphate (1 ml.), phosphorylase (15 mg.) and achroic primer (2, 1, 0.5 and 0.2 mg. respectively) in 10 ml. Liberation of phosphate was followed during incubation at 35° and when, in each digest, 0.46 mg. of phosphate had been liberated, portions (2 ml.) were withdrawn and stained with iodine (0.5 ml.) in a total volume of 25 ml.

Table 4. Comparison of experimentally determined and calculated ranges of DP in synthetic amyloses

Calculated values were obtained by Boyne (see text) and the experimental values were determined by Mould & Syngé (1954). In each case the central range of chain length (DP) estimated to contain 95% of the polysaccharide molecules is given.

	Degree of polymerization (glucose units)						
	34	47.5	77	103	160	205	250
Average chain length	23-46	35-62	64-89	80-124	150-170	140-250	170-320
Experimental range	23-45	34-61	60-94	83-123	135-185	177-233	219-281
Calculated range							

evident in the form of a visible precipitate but it may be assumed that a retrograded molecule is no longer a primer. When retrogradation begins to occur, therefore, the range of DP of the amylose molecules will begin increasingly to exceed the calculated range.

These three effects would tend to increase the range of DP in the synthetic amylose as compared with the range expected from normal kinetic considerations. The opportunity of testing their influence was provided by Mould & Syngé (1954), who examined a series of synthetic amyloses by electrokinetic ultrafiltration and provided estimates of their DP ranges for comparison with the calculated values. The latter values were calculated by Mr A. W. Boyne of the Rowett Research Institute assuming completely random multi-chain action and no variation in priming activity with increase in chain length. These assumptions lead to a Poisson distribution of chain lengths, readily permitting the determination of the central range of DP containing 95% of the polysaccharide molecules (cf. Flory, 1940). The results for a series of synthetic amyloses, ranging in average chain length from 34 to 250 glucose units, are given in Table 4.

The finding of Mould & Syngé (1954) that the mobility of a synthetic amylose in collodion is a function of its calculated average chain length provides a direct confirmation of the multi-chain nature of phosphorylase action. The results of their experiments with a series of synthetic amyloses are given in Table 4 together with the corresponding calculated ranges of DP. There is a striking agreement between calculation and observation of range of DP in the region between average DP 34 and average DP 160. Clearly none of the factors discussed above has any noticeable effect on the composition of an amylose preparation having an average DP within this range. Above DP 160 such effects are beginning to operate; the action pattern is beginning to change, not because of a change in the behaviour of the enzyme, but because of the changing properties of the primer substrate.

It is of interest to compare the results of Table 4 with the electrokinetic ultrafiltration analysis of a polysaccharide of calculated average DP 75.5 synthesized from a heterogeneous primer. This

analysis and the preparation of the primer is reported in an accompanying paper (Mould & Syngé, 1954) where it is shown that the actual DP range of the polysaccharide was far removed from the calculated DP, being 110-200 with a predominance of the higher polymers. The explanation of the difference between the two values for DP is similar to that given above in connexion with the properties of the amyloses synthesized from maltotriose and will not be enlarged upon here except to point out that the use of an inhomogeneous primer in preparing synthetic amyloses obviously leads to erroneous estimates of DP when calculation is made in terms of phosphate liberation. For this reason the results of Swanson (1948), where the iodine stain/chain length relationship of synthetic amyloses is reported, are in error.

SUMMARY

1. The action pattern of potato phosphorylase has been shown to be completely multi-chain both in the synthesis and the phosphorolysis of amylose.

2. Synthetic amyloses of known average chain length have been prepared and up to 160 their degrees of polydispersity (determined by Mould & Syngé, 1954) are shown to be entirely in agreement with values calculated on the assumption of multi-chain action.

3. The priming activities of the maltodextrins have been compared. Maltotriose is the lowest member of the series to exhibit priming activity. This activity is very weak and the higher members, malto-tetraose, -pentaose and -hexaose are much more efficient.

We gratefully acknowledge the advice and encouragement given by Professor Stanley Peat, F.R.S. One of us (J. M. B.) thanks the Department of Scientific and Industrial Research for the award of a maintenance grant. We also thank Drs D. L. Mould and R. L. M. Syngé and Mr A. W. Boyne for providing the data in Table 4.

REFERENCES

- Allen, R. J. L. (1940). *Biochem. J.* **34**, 858.
- Bailey, J. M., Thomas, G. J. & Whelan, W. J. (1951). *Biochem. J.* **49**, liv.
- Bailey, J. M. & Whelan, W. J. (1952). *Biochem. J.* **51**, xxxiii.

- Bailey, J. M., Whelan, W. J. & Peat, S. (1950). *J. chem. Soc.* p. 3692.
- Barker, S. A., Bourne, E. J., Wilkinson, I. A. & Peat, S. (1950). *J. chem. Soc.* p. 84.
- Bernfeld, P. & Meutémédian, A. (1948). *Helv. chim. acta*, **31**, 1735.
- Cori, C. F., Cori, G. T. & Green, A. A. (1943). *J. biol. Chem.* **151**, 39.
- Englard, S. & Singer, T. P. (1950). *J. biol. Chem.* **187**, 213.
- Englard, S., Sorof, S. & Singer, T. P. (1951). *J. biol. Chem.* **189**, 217.
- Flory, P. J. (1940). *J. Amer. chem. Soc.* **62**, 1561.
- French, D., Knapp, D. W. & Pazur, J. H. (1950). *J. Amer. chem. Soc.* **72**, 1866.
- French, D. & Wild, G. M. (1953). *J. Amer. chem. Soc.* **75**, 4490.
- Green, D. E. & Stumpf, P. K. (1942). *J. biol. Chem.* **142**, 355.
- Hanes, C. S. (1940). *Proc. Roy. Soc. B*, **129**, 174.
- Hobson, P. N., Whelan, W. J. & Peat, S. (1950). *J. chem. Soc.* p. 3566.
- McCready, R. M. & Hassid, W. Z. (1944). *J. Amer. chem. Soc.* **66**, 560.
- Meyer, K. H., Fischer, E. H., Staub, A. & Bernfeld, P. (1948). *Helv. chim. acta*, **31**, 2158.
- Mould, D. L. & Synge, R. L. M. (1954). *Biochem. J.* **58**, 571.
- Peat, S., Pirt, S. J. & Whelan, W. J. (1952*a*). *J. chem. Soc.* p. 705.
- Peat, S., Pirt, S. J. & Whelan, W. J. (1952*b*). *J. chem. Soc.* p. 714.
- Peat, S., Whelan, W. J. & Bailey, J. M. (1953). *J. chem. Soc.* p. 1422.
- Peat, S., Whelan, W. J. & Rees, W. R. (1953). *Nature, Lond.*, **172**, 158.
- Somogyi, M. (1945). *J. biol. Chem.* **160**, 61.
- Sumner, J. B., Somers, G. F. & Sisler, E. (1944). *J. biol. Chem.* **152**, 479.
- Swanson, M. A. (1948). *J. biol. Chem.* **172**, 825.
- Weibull, C. & Tiselius, A. (1945). *Ark. Kemi Min. Geol.* **19A**, no. 19.
- Whelan, W. J., Bailey, J. M. & Roberts, P. J. P. (1953). *J. chem. Soc.* p. 1293.
- Whelan, W. J. & Roberts, J. G. (1954). *Biochem. J.* **58**, 569.
- Whelan, W. J. & Roberts, P. J. P. (1953). *J. chem. Soc.* p. 1298.

Products of the β -Amylolytic of Maltodextrins

BY W. J. WHELAN AND J. G. ROBERTS

Department of Chemistry, University College of North Wales, Bangor

(Received 15 April 1954)

β -Amylase degrades α -1:4-linked polymers of D-glucopyranose by an end-wise action, commencing at the non-reducing chain end, in which alternate linkages are hydrolysed and maltose is liberated in the β -form (Kuhn, 1925; Hanes, 1937; Hestrin, 1949; Bird & Hopkins, 1954). In the linear series of polymers based on maltose (maltodextrins) the smallest molecule which undergoes rapid hydrolysis is maltotetraose (Whelan, Bailey & Roberts, 1953). Maltotriose is attacked very slowly with the production of glucose and maltose. From a study of changes in reducing power during hydrolysis (Whelan *et al.* 1953), and of the priming action in the enzymic synthesis of amylose after treatment of the maltodextrins with β -amylase (Whelan & Bailey, 1954), it was inferred that the rapid β -amylolysis proceeded no further than the conversion of maltotetraose and -hexaose into maltose, and of maltopentaose into equimolar amounts of maltose and maltotriose. The present work affords conclusive proof of the correctness of these deductions by the isolation and characterization of these products.

EXPERIMENTAL AND RESULTS

Maltodextrins. These were prepared from potato amylose as by Whelan *et al.* (1953).

β -Amylase. The crystalline enzyme from sweet potato was used. This was kindly provided by Dr A. K. Balls. Its activity was determined as by Hobson, Whelan & Peat (1950).

Sugar acetates. These were prepared and the maltotriose acetate was purified on Magnesol-Celite as described by Whelan & Roberts (1953).

β -Amylolytic of maltodextrins. The maltodextrin (ca. 100 mg.) was dissolved in water (10 ml.) and Somogyi's (1945*b*) deproteinizing reagents, 5% (w/v) ZnSO_4 (2 ml.) + equiv. amount of 0.4N-Ba(OH)₂, added (cf. Whelan *et al.* 1953). The supernatant solution was divided into a 1 ml. portion used to determine the maltodextrin concentration (Pirt & Whelan, 1951) and a 10 ml. portion to which was added 0.2M sodium acetate-acetic acid buffer (pH 4.8; 6 ml.), β -amylase (60 units; 1 ml.) and water to 50 ml. Measurements of reducing power were made on 1 ml. portions at intervals during incubation at 35° using the Somogyi (1945*a*) copper reagent. The hydrolyses were virtually complete after 30 min. but digestion was allowed