and its impurities to be analyzed and related to the ionic strength of the precipitating salt.

Comparison of technical difficulties involved in performing each test. In performing a constant solvent solubility test the composition of the solvent must be chosen with great care. For a satisfactory result can be attained only if the most soluble of all the proteins in the mixture saturates the solvent while at the same time the solvent must permit an analyzable quantity of the least soluble protein to dissolve. Now in order to determine the true composition of a protein solution it is necessary to proceed from the supersaturated side so that the alteration in composition caused by solvent washing of the solid protein mixture can be avoided. Working from the supersaturated side in this case is usually more difficult and less accurate.

In the case of the variable solvent and specific property solubility tests these difficulties do not occur since all that has to be done is to add increasing quantities of salt until all the protein present has been precipitated.

SUMMARY AND CONCLUSIONS

1. A new type of phase rule solubility test for protein purity is described, and quantitative graphical and mathematical methods are provided for its analysis.

2. A method by which the test described could be extended to form the basis of a general method of purification of any protein with a specific accurately measurable quality is indicated.

3. The relationship of the new test to the two previously described solubility tests is indicated and the relative merits of the three tests discussed.

4. The new test has been applied to the analysis of a liver extract relative to its content of liver esterase.

5. The preparation of pure pig liver esterase is described.

It is a pleasure to express our indebtedness to both Mr G. S. Adair, F.R.S. and Dr D. W. G. Style for their invaluable advice on many points.

REFERENCES

- Butler, A. M., Blatt, H. & Southgate, H. (1935). *J. biol. Chem.* **109**, 755.
 Cohn, E. J. (1925). *Physiol. Rev.* **5**, 349.
 Cohn, E. J. (1941). *Chem. Rev.* **28**, 395.
 Derrien, Y. (1944). *Bull. Soc. Chim. biol.* **26**, 1091.
 Falconer, J. S. & Taylor, D. B. (1945). *Nature, Lond.*, **155**, 303.
 Falconer, J. S. & Taylor, D. B. (1946). *Biochem. J.* **40**, 831.
 Herriott, R. M., Desreux, V. & Northrop, J. H. (1940). *J. gen. Physiol.* **24**, 213.
 Jameson, E. & Roberts, D. B. (1937). *J. gen. Physiol.* **20**, 475.
 Kuhn, R. & Desnuelle, P. (1937). *Ber. dtsch. chem. Ges.* **70**, 1907.
 Northrop, J. H. (1941). *J. gen. Physiol.* **25**, 465.
 Northrop, J. H. & Kunitz, M. (1930). *J. gen. Physiol.* **13**, 781.
 Roche, A., Dorier, M. & Samuel, L. (1936a). *C.R. Soc. Biol., Paris*, **121**, 1019.
 Roche, A., Dorier, M. & Samuel, L. (1936b). *C.R. Soc. Biol., Paris*, **122**, 231.
 Roche, J., Derrien, Y. & Moutte, M. (1941). *Bull. Soc. Chim. biol.* **23**, 1114.
 Shedlovsky, T. (1943). *Ann. N.Y. Acad. Sci.* **43**, Art. 6, 257.
 Sorensen, S. P. L. (1930). *C.R. Lab. Carlsberg*, **18**, No. 5.
 Sorensen, S. P. L. & Hoyrup, M. (1917). *C.R. Lab. Carlsberg*, **12**, 213.

Observations on the Nicotinic Acid Requirements of Pigs

By R. BRAUDE AND S. K. KON, *National Institute for Research in Dairying, University of Reading*,
 AND E. G. WHITE, *Research Institute in Animal Pathology, Royal Veterinary College, London*

(Received 27 June 1946)

While a large number of methods or their modifications have been described for the chemical estimation of nicotinic acid, the specificity of the rather complicated procedure has only seldom been confirmed by biological test. The work of Harris & Raymond (1939) in this country and of Waisman, Mickelsen, McKibbin & Elvehjem (1940) in America

may be quoted as the exception rather than the rule.* Both groups of workers used dogs, and it is obvious that in this country assays of this type are difficult to carry out.

* The same remark may also be applied to the numerous methods for the microbiological assay of nicotinic acid (cf. Snell & Wright, 1941).

Chick, Macrae, Martin & Martin (1938 *a, b*) were the first to demonstrate the need for nicotinic acid in the nutrition of the pig, and since then Hughes (1938, 1943), Madison, Miller & Keith (1939) and Wintrobe (1939), have confirmed and extended these observations. Since it seemed to us that the pig might prove a suitable animal for the biological assay of nicotinic acid, the experiments described below were carried out. While this first attempt has not yet yielded a satisfactory method, information was obtained on the nicotinic acid requirements of pigs which it is thought worth while to report.

EXPERIMENTAL

Plan of experiment

Twenty-four Large White normally developed and healthy weaners, about 9 weeks old, were selected from three litters of our own breeding, reared indoors. One of the litters, no. 824, consisting of twelve pigs was divided on a weight basis into two lots of six pigs, lot 824A containing the six heavier, and lot 824B the six lighter animals. Another six pigs were chosen from a litter of nine, no. 1758, and a further six from a litter of seven, no. 1760. The initial average weights of the pigs from litters 824A, 1758, 1760 and 824B were 34.2, 33.2, 28.2 and 27.0 lb. respectively. Although there were considerable differences between the litters, the variation within litters was small. One pig from each of two litters and two from the third were allocated by lot to one of the six experimental groups; each of the groups thus contained a representative of each litter, with the large litter represented twice.

The individual feeding method was adopted. Each group occupied a pen provided with small individual feeding compartments into which the animals were let, twice daily, for about 30 min. to eat their food. Owing to shortage of accommodation, pigs in groups 4, 5 and 6 had to be kept in two pens, each of which was provided with six individual feeding compartments. This meant that two pigs from group 4 shared a pen with group 5, and two with group 6. No bedding was provided and straw was rigidly excluded. Part of the concrete floor was, however, covered with wooden boards to provide a comfortable resting place for the animals.

All six groups received the diet deficient in nicotinic acid used by Chick *et al.* (1938 *a, b*) and consisting of ground yellow maize 77.5, pea meal 10.5, purified casein* 6.5, cod-liver oil 3, mineral mixture 2.5, all parts by weight. The mineral mixture was composed of calcium carbonate 2, bone ash 2, common salt 1, ferric oxide 0.08 parts. The meal mixture was mixed by hand in the piggery. The chemical

analysis of the meal mixture is given in Table 1. The meal was weighed out and fed twice daily as a wet mash, 3 lb. of water being at first used for each

Table 1. *Analysis of the basic diet*

	(%)		(%)
Moisture	12.7	Carbohydrate	60.0
Crude protein	15.3	Fibre	1.8
Fat	6.3	Ash	3.9

lb. of meal. This amount did not mix properly with the meal, and the quantity had to be reduced to approximately 2 lb./lb. of meal. This was enough to satisfy the requirements of the pigs which on a number of occasions refused water given separately. The quantity of meal fed daily was based on a normal scale for fattening pigs (cf. Braude & Foot, 1942). The pigs were weighed on Mondays, Wednesdays and Fridays, and the quantity of food given was adjusted for live-weight after each weighing. Details of the different groups are given in Table 2.

Groups 2 and 3 were selected by lot as 'positive controls'. The pigs in group 2 received in addition to their meal 10 mg. of nicotinic acid daily. The nicotinic acid was dissolved in acidulated distilled water, the required quantity measured out each day and added directly to the individual troughs just before the afternoon feeding, together with 15 g. of casein to balance the additional protein provided by the supplement of yeast given to group 3. The pigs in group 3 received in addition to their meal 30 g./pig daily of fodder yeast,* added directly to the troughs at each afternoon feeding and mixed well with the meal. The nicotinic acid content of this yeast was by chemical test about 215 μ g./g. and about 6.5 mg. in the daily dose.

The pigs in groups 4, 5 and 6 were given at first only the basic diet. It was planned that as soon as they became depleted the pigs in group 4 would receive 2 mg., those in group 5, 4 mg. and those in group 6, 8 mg. of nicotinic acid/pig daily. It soon became obvious, however, that the speed of depletion varied markedly from pig to pig, and some pigs showed signs of deficiency at a time when their pen-mates were still growing at a normal rate. It was then decided to dose the animals individually as and when they ceased to grow. By then, however, one animal, pig no. 2072 from group 6, had already died and one or two more were too seriously affected to improve when given nicotinic acid.

The pigs in group 1 were allowed to develop the deficiency symptoms and were then given a single dose of 30 mg. of nicotinic acid/pig. This method was successfully used with dogs by Schaefer, McKibbin & Elvehjem (1942) and by Smith, Carry & Hawfield (1943) for nicotinic acid assays. It measures

* G. L. Casein Cl (Lactic casein, unextracted) purchased from Glaxo Laboratories Ltd., Greenford, Middlesex.

* Kindly provided by Dr A. C. Thaysen.

the growth response and the time required for symptoms of deficiency to reappear after the administration of a known quantity of the substance which is being assayed. As soon as the symptoms reappear the dosing can be repeated.

The experiment commenced on 6 October 1944 and continued till the end of March 1945, thus involving the full fattening period of a bacon pig.

The average individual daily live-weight gains were as follows:

Group 2, basic diet + 10 mg. nicotinic acid ...	0.88 lb.
Group 3, basic diet + 30 g. dried yeast ...	0.88 lb.
Group 1, basic diet alone ...	0.34 lb.
Group 4, basic diet alone ...	0.39 lb.
Group 5, basic diet alone ...	0.48 lb.
Group 6, basic diet alone ...	0.30 lb.

Fig. 1. Mean growth curves of groups of four pigs receiving the basic diet deficient in nicotinic acid alone or supplemented with yeast or nicotinic acid.

RESULTS

The effects of dietary deficiency and of dosing with nicotinic acid

The growth of the pigs. On the 47th day of the experiment pig no. 2072 receiving the basic diet died. The growth curves up to that date of the pigs in the six experimental groups are presented in Fig. 1.

There can be no doubt that the pigs in the two 'positive control' groups, i.e. those receiving a supplement of 10 mg. of nicotinic acid or of 30 g. of yeast, grew at a much better rate than the pigs receiving the basic diet alone. Although there was some variation between individuals in each of the groups it is clear that the basic diet was deficient in nicotinic acid. Apart from subnormal growth, some

Fig. 2. Individual growth curves of pigs receiving the basic diet deficient in nicotinic acid, and details of various treatments (see also Table 2).

Table 2. *Details of treatment, growth and autopsy findings for individual pigs*

N.A. = nicotinic acid. I.C.V. = ileo-caecal valve. N.E. = necrotic enteritis. P. = <i>Pasteurella septica</i> . H. = A member of the <i>Haemophilus</i> group. Su. = succinyl sulphathiazole. Pe. = peritonitis.									
Group	Diet	Pig no. and litter no.	Treatment of individual animals	Symptoms and growth	Autopsy findings	Bacteriology			
1	Basic diet until symptoms of deficiency, then single doses of 30 mg. N.A., repeated when signs of deficiency reappeared	2079	30 mg. N.A. on 54th, 77th, 115th and 138th day	Scoured but responded for a time after each of the first 3 doses of N.A. Grew well after 4th dose and slaughtered on 171st day	Killed at abattoir. Lungs only examined—normal	Sterile			
		824 A							
		2058	30 mg. N.A. on 56th day	Grew well and slaughtered on 171st day. May have obtained some N.A. from companions in same pen	Killed at abattoir. Lungs only examined—normal	Sterile			
		1760							
		2066	30 mg. N.A. on 54th, 82nd and 94th day	Scoured severely until given first dose of N.A. First two doses temporary effect, third dose no effect. Died on 101st day	Mouth ulcers. N.E. from I.C.V. to rectum. Half of each lung consolidated: pleurisy, Pe.	P. from lung			
		1758							
2	Basic diet, with 10 mg. N.A. daily	2078	30 mg. N.A. on 54th day	Off food and scoured. Temporary improvement after N.A. but relapsed and died on 66th day	Mouth ulcers. Nearly half of each lung consolidated: Pe.	P. from lung			
		824 B							
		2073	N.A. withdrawn on 94th day.	Grew well in spite of withdrawal of N.A. and administration of Su. Slaughtered on 157th day	Killed at abattoir. Lungs only examined. Small areas of long-standing pneumonia	1 colony of P. from lung			
		824 A	Su. 15 g. daily 139th–145th day						
		2056	As above	As above	Killed at abattoir: no autopsy	None			
		1760							
		2069	N.A. withdrawn on 94th day	Grew well and slaughtered on 157th day	As above	As above			
		1758							
		2081	N.A. withdrawn on 94th day: given again from 98th to 145th day	Grew well until N.A. withdrawn, then lost weight. Recovered when N.A. given. Second withdrawal had no effect. Slaughtered in good condition on 157th day	As above	As above			
		824 B							

N.A. = nicotinic acid.

I.C.V. = ileo-caecal valve.

N.E. = necrotic enteritis.

P. = *Pasteurella septica*.H. = A member of the *Haemophilus* group.

Su. = succinyl sulphathiazole.

Pe. = peritonitis.

Table 2 (cont.)

Group	Pig no. and litter no.	Diet	Treatment of individual animals	Symptoms and growth	Autopsy findings	Bacteriology
3	2074	Basic diet, with 30 g. fodder yeast (about 6.5 mg. N.A.) daily	Yeast withdrawn on 94th day.	Grew well in spite of withdrawal of yeast and administration of Su. Slaughtered on 157th day	Killed at a battery: no autopsy	None
	824 A		Su. 15 g. daily 139th-145th day			
	2060		Yeast withdrawn on 94th day	Grew well throughout the experimental period and slaughtered on 157th day	As above	As above
	1760					
	2065		Yeast withdrawn on 94th day.	As above	As above	As above
	1758		Su. 15 g. daily 139th-145th day			
	2084		Yeast withdrawn on 94th day:	Grew well until N.A. withdrawn, then lost weight until N.A. was replaced. Failed to react to second withdrawal of N.A. Slaughtered on 157th day	Killed at a battery. Lungs only examined: small areas of long-standing pneumonia	Few colonies of P. from lung
	824 B		given again from 98th to 145th day			
4	2085	Basic diet until symptoms of deficiency, then 2 mg. N.A. daily	2 mg. N.A. daily from 67th day	Lost weight but recovered when N.A. added. On 119th day separated from companions receiving high level of N.A. Died on 122nd day	N.E. involving most of large intestine. Half of each lung consolidated: Pe.	P. from heart blood and lung. H. from lung
	824 A					
	2062		2 mg. N.A. daily from 67th day	Lost weight, but recovered when N.A. added. On 112th day separated from companions receiving high level of N.A. Died on 118th day	More than half of each lung consolidated: pleurisy with adhesions	P. and H. from lung
	1760					
	2064		2 mg. N.A. daily from 97th to 105th day, 8 mg. 105th to 122nd day, 100 mg. on 122nd day only	Grew well until separated from companions on 89th day. Daily doses of 2 and 8 mg. N.A. failed to cure, also single dose of 100 mg. Died on 123rd day	N.E. involving most of colon. Two-thirds of each lung consolidated. Fibrinous pericarditis and Pe.	P., H. and <i>Esch. coli</i> from lung
	1758					
	2072		No N.A. given	Died on 47th day	Acute enteritis in small intestine; no N.E. Mouth ulcers. One-third of each lung consolidated	Cultures overgrown by 'spreaders'
	824 B					

5	Basic diet until symptoms of deficiency, then 4 mg. N.A. daily	2075	4 mg. N.A. daily from 98th to 111th day, then 8 mg. daily	Grew well until removed from companions on 98th day. Recovered with N.A., relapsed, recovered after larger daily dose and slaughtered on 171st day	Killed at abattoir. Lungs only examined. One-third of each lung consolidated	P. from lung
		824 A				
		2057	4 mg. N.A. daily from 54th day	Lost weight but improved with N.A. supplement. On 108th day (cold spell) became ill, and died on 114th day	N.A. confined to 10 sq.cm. of caecal mucosa. One-half to two-thirds of each lung consolidated	P. and H. from lung
		1760				
		2067	4 mg. N.A. daily from 59th to 139th day, then 8 mg. daily	Lost weight and scoured. Improved with addition of N.A. but relapsed. Regained condition with larger daily dose and slaughtered on 171st day	Killed at abattoir. Lungs only examined: normal	Sterile
6	Basic diet until symptoms of deficiency, then 8 mg. N.A. daily	2083	4 mg. N.A. daily from 52nd day	Lost weight and scoured. Died on 58th day	Mouth ulcers. One-third of each lung consolidated: Pe.	P. from lung
		824 B				
		2076	8 mg. N.A. daily from 53rd day	Lost weight and scoured. Recovered when N.A. added. Slaughtered on 171st day	Killed at abattoir. Lungs only examined. Pneumonic areas of lung standing along inferior border of each lung	Sterile
		824 A				
		2059	8 mg. N.A. daily from 63rd day	Lost weight and scoured. Recovered when N.A. added. Paraplegia developed following a fight on 150th day. Slaughtered on 171st day	Killed at abattoir. Bladder dilated, hydronephrosis right renal pelvis. One-third of each lung consolidated	P. from lung
		2063	8 mg. N.A. daily from 66th day	Lost weight and scoured. Recovered when N.A. added. Slaughtered on 171st day	Killed at abattoir. Lungs only examined. One-third of each lung consolidated and one abscess at apex of each lung	P. from lung. <i>α-strep.</i> , H. and <i>C. pyogenes</i> from abscesses
		1758				
		2080	No N.A. given	Scoured and lost weight. Died on 52nd day	Mouth ulcers. N.E. from i.c.v. to 40 cm. from anus, involving one-sixth of mucosal surface. One-quarter of each lung consolidated	P. from lung
		824 B				

of the pigs affected scoured badly, refused food and became thin and emaciated. Their breathing was at times laboured and on occasion they looked somewhat lethargic.

Fig. 2 presents the individual growth curves of the pigs, and Table 2 details of their treatment.

The 'positive control' groups. All eight pigs in the 'positive control' groups grew at a very satisfactory rate and no adverse symptoms of any kind were observed. On the 94th day of the experiment it was decided to discontinue the nicotinic acid supplements which these pigs had so far been receiving daily, in order to see whether symptoms of depletion could be produced in older pigs. The average daily gains in weight of the pigs up to that time were 1.22 lb. for group 2 (10 mg. nicotinic acid daily) and 1.21 lb. for group 3 (30 g. yeast daily). It is apparent from Fig. 2 that growth was very uniform throughout.

One pig in each group, no. 2081 in group 2 and no. 2084 in group 3, both belonging to the lighter half of litter 824, reacted immediately and drastically to the removal of the supplements. They refused a considerable portion of food, lost weight and rapidly became thin. The remaining pigs in both groups did not appear affected by the change of the diet and continued to grow at a very satisfactory rate.

The two affected pigs were taken out of their pens, put together in a separate pen and given again the nicotinic acid supplements on the 98th day of the experimental period. Both gradually took to the food and after a few days became normal, and resumed growth at a normal rate. On the 145th day their nicotinic acid supplements were again withdrawn. It can be seen from Fig. 2 (groups 2 and 3) that on this occasion the two pigs reacted only very slightly to the change and continued to grow normally until they were slaughtered on the 157th day of the experiment.

The remaining three pigs in each of groups 2 and 3 continued to grow normally though they were receiving only the basic diet. Their average daily live-weight gains up to the 138th day of experiment when the diet was again changed were 1.04 lb. in group 2 and 1.05 lb. in group 3.

It was obvious that the removal of nicotinic acid from the diet had much less effect on pigs weighing about 140–150 lb. than on young weaners. Now only two pigs out of eight showed signs of deficiency, and even these two later resisted deprivation of nicotinic acid. Three possible explanations of this behaviour were considered. (1) The pig requires nicotinic acid only for the early stage of its growth. (2) Older pigs are able to synthesize the factor either in their body or with the help of intestinal bacteria. (3) Pigs can store nicotinic acid in their organs and make use of it when given a diet deficient in this factor.

In recent years the existence of microbial synthesis of vitamins in the gut has been firmly established, and it has been shown that even non-ruminants can under certain conditions derive benefit from the vitamins so formed (cf. review by Kon, 1945). Ellinger, Coulson & Benesh (1944) indicated that the nicotinamide requirement of man can be to some extent covered by the contribution made by the

intestinal microflora. To find out whether the same was true of our pigs we gave to pigs nos. 2073 and 2056, and 2074 and 2065 respectively in groups 2 and 3, which were growing normally on the diet deficient in nicotinic acid, 15 g. of succinyl sulphathiazole/pig daily for 7 days, starting on the 139th day of the experiment. A third pig in each of the groups, selected by lot, was left untreated. Urine samples were collected before and after treatment, and analyzed for nicotinamide methochloride by the method of Coulson, Ellinger & Holden (1944). The results of the urine test were inconclusive. The pigs receiving the drug continued to grow normally, and were apparently not affected in any way. On the 157th day all six pigs were slaughtered.

The pigs receiving daily 2, 4 and 8 mg. of nicotinic acid. The growth of pigs in groups 4, 5 and 6 up to the 47th day of the experiment has already been mentioned on p. 845. On that day the first death occurred on the diet deficient in nicotinic acid. Pig no. 2072 (litter 824 B) from group 4 had stopped growing after a fortnight on the basic diet. It was not given nicotinic acid supplement as most of its pen-mates were still growing, and it was originally the intention to avoid dosing some pigs and not others within the same pen. As already stated, however, we had to modify this procedure, and subsequently started dosing each pig as the symptoms of deficiency became obvious.

Fig. 2, group 4, gives growth curves and details of dosing of pigs in group 4. From the 56th day of experiment two pigs in this group, nos. 2085 and 2062, received a supplement of 2 mg. of nicotinic acid/pig daily. When the treatment was started these pigs were showing deficiency symptoms. Both responded to the dose and resumed growth. They grew normally until they were removed from the pens in which hitherto they were together with pigs receiving higher doses of nicotinic acid. The separation was carried out as soon as alternative accommodation became available. Pig no. 2062 separated on the 112th day of the experiment died within 6 days, while pig no. 2085 separated on the 119th day died 3 days later. Extremely cold weather prevailed during the week in January when these two deaths occurred, and it is quite possible that the lowered temperature in the piggery contributed to the death of the pigs weakened by a deficient diet.

The third pig in this group, no. 2064, continued to grow, after an initial setback, without any nicotinic acid supplement. This pig was housed with the pigs of group 6, and after a time it became apparent that it was benefiting in some way from the supplements which its pen-mates were receiving. Taking this into account it was decided on the 89th day of the experiment to house this pig separately. Within a few days it showed severe symptoms of deficiency, and a daily dose of 2 mg. nicotinic acid started on the 97th day had no beneficial effect. The pig continued to decline rapidly and on the 105th day the dose was increased to 8 mg. daily, without effect, however. A 100 mg. dose given on the 122nd day was equally ineffective and the pig, which by then was in a very poor condition, scouring severely and breathing very heavily, died on the 123rd day.

In group 5, three pigs developed deficiency symptoms and were given 4 mg. of nicotinic acid/pig daily during the week which started on the 52nd day of the experiment. Pig

no. 2083 belonging to litter 824 B was given the supplement on the 52nd day. At that time it was already in a bad condition which gradually became worse, and it died 6 days later.

Pig no. 2067 was given the supplement on the 59th day. It was then scouring very badly and was generally in a poor condition. For about 10 days the treatment did not seem to have much effect, the pig getting gradually worse. After a fortnight, however, a definite improvement was observed and the pig started putting on weight at about normal rate. It suffered a setback on the 115th day of experiment, possibly associated with the very cold weather prevailing at the time. It recovered, but within a few days was scouring again and on the 140th day was in such a poor condition that it was decided to increase its nicotinic acid supplement to 8 mg./day. From that date the pig, although still scouring for some time, started to put on weight and eventually attained a normal rate of growth and was sent to slaughter on the 171st day.

Pig no. 2057 was given 4 mg. of nicotinic acid from the 54th day onward and responded with a fairly normal growth until the 108th day, which coincided with the very cold weather period. It refused food and died on the 114th day.

The fourth pig in this group, no. 2075, grew at a normal rate. Its behaviour in the pen, however, made it obvious that it was benefiting from the supplement given to its pen-mates; it would follow them about when they left their feeding pens, and lick their snouts. On the 89th day the pig was taken out of the pen and housed separately. Within a week deficiency symptoms became apparent; the pig lost appetite and weight and its breathing became laboured. On the 98th day daily dosing with 4 mg. of nicotinic acid was started. At first the pig responded well to the dose, but within a fortnight deficiency symptoms reappeared, and the dose was increased to 8 mg. The pig soon recovered and continued to grow normally until it was slaughtered as a full baconer on the 171st day of experiment.

In group 6 all four pigs gradually developed the deficiency symptoms. Pig no. 2080 from litter 824 B succumbed at a very early stage, on the 52nd day, before curative treatment was started. Within a fortnight the remaining three pigs were showing symptoms and were given the daily dose of 8 mg. of nicotinic acid. Dosing of pig no. 2076 which had previously had a setback from which it partly recovered was started on the 53rd day, that of pig no. 2059 on the 63rd day and of pig no. 2063 on the 66th day of experiment. All three pigs responded to the treatment. Pig no. 2076 had two obvious setbacks from which it recovered spontaneously, and was finally slaughtered on the 171st day of experiment, having attained bacon weight. Pig no. 2063, after a rather slow start, eventually grew at a normal rate and attained the same weight at slaughter on the 171st day as pig no. 2076. Pig no. 2059 grew at a normal rate until the 150th day of the experiment when, during a fight, it was severely bitten on the tail and bled profusely. It completely lost control of its legs, becoming paralyzed, and within a week stopped eating. It could not get up even with assistance, had to be separated from its group-mates, and be fed when lying down. It was decided not to disturb it by further weighing. The pig was kept until the 171st day of experiment when, together with its group-mates, it was sent to slaughter.

Pigs dosed intermittently with large doses of nicotinic acid. In group 1, three pigs, nos. 2079, 2066 and 2078, were showing deficiency symptoms on the 54th day of experiment. They were then given a single dose of 30 mg. nicotinic acid.

The fourth pig, no. 2058, which at that time appeared to be normal, was separated from the remainder of the pen. It would not, however, settle down in a pen on its own and had to be brought back to its group; it received 30 mg. nicotinic acid on the 56th day.

Pig no. 2078 which belonged to litter 824 B became very ill after a short initial improvement, refused food and its breathing became laboured. It died on the 66th day.

Pig no. 2066 which, when given its dose, was scouring heavily, responded to the treatment and grew fairly satisfactorily for about 18 days, the scouring having stopped. On the 74th day, however, the scouring reappeared, the pig refused food again and was losing weight when, on the 82nd day, it was given another dose of 30 mg. of nicotinic acid. It again responded to the dose, but within a week was ill again. On the 94th day another 30 mg. dose was given, but this time with no effect at all. The pig's condition became worse and it died on the 101st day.

Pig no. 2079 responded satisfactorily to the first dose of nicotinic acid, but after about a fortnight started scouring, and its condition deteriorated. On the 77th day it was given another single dose of 30 mg. nicotinic acid, responded to it again and for about 6 weeks grew at a fairly normal rate. At the end of this period it again developed deficiency symptoms and was dosed further on the 115th day. It again improved, but after about 3 weeks had a relapse and was given a fourth dose on the 138th day which enabled it to grow at a fair rate until it was slaughtered on the 171st day of experiment.

Pig no. 2058 was given a dose on the 156th day before showing any symptoms and continued to grow at a normal rate until it was slaughtered on the 171st day of experiment. Changes in the slope of its growth curve (Fig. 2, group 1) which coincide with the administration of nicotinic acid to its pen-mate suggest that it was benefiting in some way from the vitamin given to the latter.

Pathological findings

A summary of the main pathological and bacteriological findings is given in Table 2. Every pig which died was examined and cultures were made from the lungs, intestinal contents and mesenteric lymph nodes. Those pigs that survived until bacon weight and were eventually slaughtered for bacon were not studied except that the lungs of eight of them were brought back to the laboratory for bacteriological and histological examination.

Skin. Chick *et al.* (1938 *a, b*) described scurvy, scabby skin and dermatitis in pigs on a pellagra-producing diet, improving with nicotinic acid addition. Other writers have since referred to changes in the skin, but usually to a dirty and scurvy skin and not to actual dermatitis. Worden & Slavin (1944) found the skin normal in pigs receiving diets deficient in nicotinic acid.

None of the pigs in the present experiments showed dermatitis although ailing pigs had a rough staring coat. It is possible that some of the skin lesions described in pigs on deficient diets, particularly when the changes were prominent about the ears, were due to sarcoptic mange.

Mouth lesions. Ulceration of the buccal mucosa was described by Miller & Rhoads (1935) in pigs which developed achlorhydria and anaemia on a modified Goldberger black-tongue diet. Four of the pigs in the present experiments showed at autopsy small areas of ulceration of the buccal mucosa. The ulcers were 2–5 mm. in diameter and were situated in the region of the incisors and canines and where the buccal mucosa is reflected from the outer surface of the gum to pass up the inside of the cheek. The lesions would readily be missed if not particularly looked for.

Necrotic enteritis. It is agreed that necrotic enteritis is a fairly constant finding in nicotinic acid deficiency in pigs, although Miller & Rhoads (1935) and Hughes (1938) do not refer to its presence in animals on a pellagra-producing diet, though the animals of Hughes (1938) recovered when nicotinic acid was added. Birch, Chick & Martin (1937) found necrotic enteritis affecting the whole of the caecum and upper half of the large intestine of pigs receiving a diet deficient in nicotinic acid.

We found the condition in five of the nine pigs which died in our experiment. The extent of the lesions varied from an area of about 10 sq.cm. in the caecum in one animal (pig no. 2057) to the whole of the caecum and large intestine down to within a few cm. of the anus, with about one-quarter of the mucosa necrotic (pig no. 2066).

Anaemia. Birch *et al.* (1937) found a progressive simple anaemia in their deficient pigs. Addition of iron salts was without effect. Worden & Slavin (1944) made a similar observation, and most workers seem agreed on this point.

Unfortunately, no haemoglobin determinations were made in our experiments and, although slight anaemia was a constant finding at autopsy, it was never as severe as is found in piglet anaemia due to deficiency of iron.

Association with Salmonella suispestifer. Birch *et al.* (1937) failed to isolate *S. suispestifer* or any other pathogenic organism from affected pigs, and this finding was confirmed later (Chick *et al.* 1938*b*). Worden & Slavin (1944) isolated the organism from the mesenteric lymph nodes only of one of two pigs which died on the deficient diet. It seems unlikely therefore that this organism is concerned in producing the lesions. This is confirmed by our own inability to isolate it from any of the nine pigs which died, including the five pigs with lesions of necrotic enteritis. It seems likely that the infectious enteritis of swine associated with *S. suispestifer* is a different condition.

Breed (1942), for instance, attempted to differentiate necrotic enteritis from infectious enteritis (salmonellosis). The former disease is characterized by slow onset, little or no pyrexia, necrotic enteritis of the caecum and large intestine, absence of acute inflammatory changes, no splenic enlarge-

ment and no changes in any lymph nodes other than those draining the affected part of the intestine. Infectious enteritis, on the other hand, is sudden in onset, associated with pyrexia, gastric haemorrhage and often ulceration, acute enteritis with necrotic changes, enlarged and congested spleen, renal haemorrhages, and lymphadenitis of all the lymph nodes.

In this connexion it is of interest to note that recent attempts to produce enteritis by administration of cultures of *S. suispestifer* from naturally affected pigs, using a variety of strains, have been unsuccessful (Gwatkin & Moynihan, 1944, 1945). Even by lessening resistance by chilling the animals it was not possible to reproduce the condition. It is difficult to explain the apparently consistent success of Murray, Biester, Purvin & McNutt (1927) and of Biester, Murray, McNutt & Purvin (1928) in reproducing necrotic enteritis by feeding *S. suispestifer* to pigs. These authors mentioned ulceration of the buccal mucosa in three of the pigs and gave an excellent account of the pathology of the condition and of the part played by *S. suispestifer* and by *Fusiformis necrophorus* in causing the necrotic lesions. They give, unfortunately, no adequate details of the actual organism used and no information as to the diet employed.

Lung lesions in the present experiments. All nine pigs which died when receiving diets deficient in nicotinic acid showed pneumonic changes in the lungs. Both lungs were always involved, and the extent of the lesions varied from one-quarter to two-thirds of the total volume of lung tissue. Affected areas were consolidated, sank in water, and were usually covered by an inflamed pleura with a fibrinous deposit and, in cases of some standing, by fibrous adhesions between visceral and parietal layers. All lungs affected with pneumonia yielded *Pasteurella septica*, except in one animal in which all cultures were contaminated by 'spreaders'. In five animals a member of the *Haemophilus* group was also obtained, and in one of them *Esch. coli* was isolated in addition.

A sero-fibrinous pleurisy was present in one of the two pigs which died on a nicotinic acid-deficient diet in the experiment described by Worden & Slavin (1944), but it is not stated whether a *Pasteurella* was isolated.

The lungs of eight animals which reached bacon weight and were quite healthy were examined bacteriologically. Five showed small pneumonic areas and from three of them a *Haemophilus* was isolated; cultures from the fourth animal remained sterile. The lungs of the fifth pig showed areas of consolidation and an apical abscess in each lung; *Pasteurella septica* was isolated from the lung tissue and a *Haemophilus*, α -haemolytic streptococci and *Corynebacterium pyogenes* from the abscesses. Three of the eight animals had macroscopically normal lungs, and cultures from these were sterile.

The pneumonia did not appear to be contagious, since sick animals remained in the same pen as healthy ones without the latter becoming affected. The ultimate cause of death in most of our animals

was pneumonia. A fibrinous pleurisy and peritonitis with adhesions was a common finding. The exact part played by *Pasteurella septica* and the haemophilus is not clear. The presence of a haemophilus suggested the possibility of the swine influenza virus being present, but several attempts to demonstrate it by intranasal instillation of lung material into ferrets were unsuccessful.

The deaths usually occurred during cold spells, and it seemed that chilling, together with a deficient diet, allowed the *Pasteurella*, *Haemophilus* organisms, and possibly a virus, to multiply.

That a pasteurella is widespread in the piggery is shown by the fact that it was isolated from the lungs of several apparently normal pigs and also from pneumonic lungs of pigs in several other experiments during recent months.

DISCUSSION

Our findings confirm the view of Chick *et al.* (1938 *a, b*) that the diet used by them is deficient in nicotinic acid and does not by itself support the growth of pigs. All sixteen of our pigs which received from the beginning of the experiment the basic diet alone sooner or later ceased to grow and showed symptoms of deficiency. We have also confirmed that the shortcomings of this diet can largely be made good by the addition of yeast or of nicotinic acid. In fact, pigs which received the basic diet supplemented with either 30 g. of fodder yeast or 10 mg. of nicotinic acid daily grew at rates well within the normal range of those obtained under our conditions with good practical rations. Chemical tests showed that the daily supplement of yeast provided about 6.5 mg. of nicotinic acid, and it may be taken therefore that from 6.5 to 10 mg. daily is ample to satisfy the nicotinic acid requirements of pigs receiving the diet of Chick *et al.* (1938 *a, b*).

Further light on the need of the pig for nicotinic acid was thrown by the curative tests. These showed that 2 or 4 mg./pig daily did not allow satisfactory recovery once deficiency symptoms became manifest. On the other hand, provided the animal was not too far gone, 8 mg. generally restored normal growth and performance.

The results of the curative tests agree well with the findings in the prophylactic experiments and confirm the view that the needs for nicotinic acid of young pigs receiving the diet of Chick *et al.* (1938 *a, b*) are met by a daily supply of 6.5–10 mg. It is possible that a dose of 4–5 mg. would be sufficient.

The puzzling observations with pigs in the later stages of fattening make it much more difficult to assess their requirements. With two exceptions pigs which had reached the weight of about 140 lb. continued to grow at a normal rate when the nicotinic acid supplement was withdrawn, and the rate of growth was not affected by the administration of

succinyl sulphathiazole. With dogs receiving a diet of the same type as ours, West (1941) noticed that sulphapyridine inhibited the curative action of nicotinic acid. Very recently, however, Najjar, Holt, Johns, Medairy & Fleischmann (1946) reported that they had been unable to suppress the biosynthesis of nicotinamide in man by administration of succinyl sulphathiazole.

Even the two pigs which were at first affected by the withdrawal of nicotinic acid but recovered when the supplement was resumed, were later able to grow satisfactorily without further additions. It is of interest that Wintrobe, Stein, Folis & Humphreys (1945), who used synthetic diets, found that pigs which received only vitamin B₁ and riboflavin benefited from an addition of nicotinic acid, but that they could dispense with this factor when pyridoxin, choline and calcium pantothenate were included. On the other hand, Schaefer *et al.* (1942) produced typical black-tongue symptoms in dogs with a synthetic diet similarly supplemented but lacking in nicotinic acid. Wintrobe *et al.* (1945) criticized the use of the diet of Chick *et al.* (1938 *a, b*) on the score that it is deficient in factors other than nicotinic acid. While not denying this possibility we would point out that our pigs which received this diet supplemented only with nicotinic acid grew at a normal rate. According to Wintrobe *et al.* (1945) protein is the main shortcoming of this diet. As compounded by us it contained 15.3 % crude protein of good quality. Moreover, the supplement of yeast supplied approximately 15 g. of protein/pig daily. In the initial stages of the experiment when the pigs were consuming daily approximately 1 kg. of the diet this supplement would increase the dietary protein to about 17 %. The protein in the diet of the pigs which received crystalline nicotinic acid was similarly raised by the addition of extra casein.

Recent work has shown that tryptophan can replace nicotinic acid in the diet of the rat (Krehl, Teply, Sarma & Elvehjem, 1945; Krehl, Sarma, Teply & Elvehjem, 1946; Krehl, Sarma & Elvehjem, 1946; Singal, Briggs, Sydenstricker & Littlejohn, 1946) and of the chick (Briggs, 1945), and it is highly probable that the same will prove true for the pig. The nicotinic acid requirement of this animal may therefore vary with the tryptophan content of its diet, and the values now quoted may be only of limited application.

The pigs in groups 2 and 3 received some additional tryptophan in the casein and yeast supplements, which both contain about 1.2 % of the amino-acid (Block & Bolling, 1945). This small amount may have contributed to the effect of nicotinic acid which the pigs received simultaneously, but the performance of the pigs in group 6 which received only nicotinic acid shows that the effect could not have been marked.

It will be recalled that in pigs receiving the diet deficient in nicotinic acid the symptoms were failure to maintain normal weight gain, refusal of food, scouring, and laboured respiration. No skin disorders of any kind were observed. The only constant lesion in pigs which died on the deficient diet was pneumonia; necrotic enteritis was present in only five of the nine animals. The effect of the diet deficient in nicotinic acid seems to have been to lower general resistance to infection so that the animals readily succumbed to pneumonia and septicaemia associated with a pasteurella, particularly during cold weather, whereas litter-mates in the same pen but on a diet supplemented with nicotinic acid or yeast remained healthy. A haemophilus was also isolated from the lungs, and the presence of a virus such as that of swine influenza cannot be entirely ruled out. Pneumonia, with associated pleurisy and pericarditis, has been found in a number of animals in the piggery during various nutritional experiments and seems in no way specific to nicotinic acid deficiency. On the other hand, necrotic enteritis not associated with salmonella infection, and ulcers of the buccal mucosa have not been seen among the large number of animals which died or were slaughtered during the past five years, and it seems that they may be specific lesions.

We have pointed out in the body of this paper that the condition of some of the deficient pigs improved when their pen-mates were given nicotinic acid. The benefit may have been derived from the consumption of excreta, though this was never observed to any extent. Whatever the cause, this factor made quantitative measurements more difficult, and there is no doubt that for such purpose individual pig pens are necessary. Apart from that, our results indicate that

the response of the pig to different levels of nicotinic acid would be sufficiently graded to form a basis of a quantitative growth test. Whether the relatively large number of pigs necessary to provide groups of sufficient size would make it practicable is another matter.

The method of intermittent dosing successfully used by Schaefer *et al.* (1942) and Smith *et al.* (1943) with dogs might prove more satisfactory, and our preliminary results show that in certain cases the method might be made to work with pigs. It would certainly be worth while to investigate this possibility with individual pigs kept separately.

SUMMARY

1. The findings of Chick *et al.* (1938 *a, b*) that the diet described by them causes symptoms of nicotinic acid deficiency in pigs have been confirmed.
2. The amount of nicotinic acid needed to prevent or cure the deficiency caused by this diet is probably between 5 and 10 mg. daily, supplied either as food yeast or as the crystalline substance.
3. Six out of eight pigs which received supplements of nicotinic acid until they reached a weight of 140–150 lb. continued to grow at a normal rate when the supplement was discontinued, and their subsequent growth was not affected by the administration of succinyl sulphathiazole.
4. The lesions found at autopsy in nine pigs which died during the experiments are described.

We are indebted to Prof. Bosworth, Dr Lovell and Miss C. Clark, of the Research Institute in Animal Pathology, for the bacteriological examinations, to Dr S. J. Rowland, of the National Institute for Research in Dairying, for the analysis of the diet, and to Dr A. C. Bottomley, of the same Institute, for the chemical estimation of the nicotinic acid in the yeast.

REFERENCES

- Biester, H. E., Murray, C., McNutt, S. H. & Purvin, P. (1928). *J. Amer. vet. med. Ass.* **72**, 1003.
- Birch, T. W., Chick, H. & Martin, C. J. (1937). *Biochem. J.* **31**, 2065.
- Block, J. & Bolling, D. (1945). *Arch. Biochem.* **6**, 277.
- Braude, R. & Foot, A. S. (1942). *J. agric. Sci.* **32**, 70.
- Breed, F. (1942). *J. Amer. vet. med. Ass.* **100**, 27.
- Briggs, G. M. (1945). *J. biol. Chem.* **161**, 751.
- Chick, H., Macrae, T. F., Martin, A. J. P. & Martin, C. J. (1938*a*). *Biochem. J.* **32**, 10.
- Chick, H., Macrae, T. F., Martin, A. J. P. & Martin, C. J. (1938*b*). *Biochem. J.* **32**, 844.
- Coulson, R. A., Ellinger, P. & Holden, M. (1944). *Biochem. J.* **38**, 150.
- Ellinger, P., Coulson, R. A. & Benesh, R. (1944). *Nature, Lond.*, **154**, 270.
- Gwatkin, R. & Moynihan, I. W. (1944). *Canad. J. comp. Med.* **8**, 66.
- Gwatkin, R. & Moynihan, I. W. (1945). *Canad. J. comp. Med.* **9**, 71.
- Harris, L. J. & Raymond, W. D. (1939). *Biochem. J.* **33**, 2037.
- Hughes, E. H. (1938). *Hilgardia*, **11**, 595.
- Hughes, E. H. (1943). *J. Anim. Sci.* **2**, 23.
- Kon, S. K. (1945). *Proc. Nutrit. Soc.* **3**, 217.
- Krehl, W. A., Sarma, P. S. & Elvehjem, C. A. (1946). *J. biol. Chem.* **162**, 403.
- Krehl, W. A., Sarma, P. S., Teply, L. J. & Elvehjem, C. A. (1946). *J. Nutrit.* **31**, 85.
- Krehl, W. A., Teply, L. J., Sarma, P. S. & Elvehjem, C. A. (1945). *Science*, **101**, 489.
- Madison, L. C., Miller, R. C. & Keith, T. B. (1939). *Science*, **89**, 490.
- Miller, D. K. & Rhoads, C. P. (1935). *J. clin. Invest.* **14**, 153.
- Murray, C., Biester, H. E., Purvin, P. & McNutt, S. H. (1927). *J. Amer. vet. med. Ass.* **72**, 34.
- Najjar, V. A., Holt, L. E., Jr., Johns, G. A., Medairy, G. C. & Fleischmann, G. (1946). *Proc. Soc. exp. Biol., N.Y.*, **61**, 371.

- Schaefer, A. E., McKibbin, J. M. & Elvehjem, C. A. (1942). *J. biol. Chem.* **144**, 679.
- Singal, S. A., Briggs, A. P., Sydenstricker, V. P. & Littlejohn, J. (1946). *Fed. Proc.* **5**, 154.
- Smith, S. G., Carry, R. & Hawfield, H. (1943). *J. Nutrit.* **25**, 341.
- Snell, E. E. & Wright, L. D. (1941). *J. biol. Chem.* **139**, 675.
- Waisman, H. A., Mickelsen, O., McKibbin, J. M. & Elvehjem, C. A. (1940). *J. Nutrit.* **19**, 483.
- West, R. (1941). *Proc. Soc. exp. Biol., N.Y.*, **46**, 369.
- Wintrobe, M. M. (1939). *Amer. J. Physiol.* **126**, 375.
- Wintrobe, M. M., Stein, H. J., Follis, R. H., Jr. & Humphreys, S. (1945). *J. Nutrit.* **30**, 395.
- Worden, A. N. & Slavin, G. (1944). *J. comp. Path.* **54**, 57.

Haemoglobin of *Gastrophilus* Larvae. Purification and Properties

By D. KEILIN AND Y. L. WANG, *Molteno Institute, University of Cambridge*

(Received 25 May 1946)

The object of this paper is the study of the haemoglobin of the larvae of *Gastrophilus* (horse bot fly) which live as parasites in the stomach of the horse. As the haemoglobin occurs only in certain stages of the development of this insect and is localized in highly specialized cells, a brief account will be given of the life history of *Gastrophilus intestinalis* and the structure of the cells harbouring the pigment.

Our knowledge of the life history of these flies is based partly upon an extensive study of the parasite in its normal host and partly upon its behaviour in small experimental animals (see Dinulescu, 1932). The latter applies especially to the stages immediately following the emergence of the larvae from their eggs. The life history of *Gastrophilus* thus reconstructed can be summarized as follows (Keilin, 1944):

'The female fixes its eggs to the hairs of the horse, usually those of the legs, breast and abdomen. In about 5-6 days the larvae are completely formed but they emerge only on contact with the tongue of the host and penetrate under its epithelial layer, forming a superficial gallery. They migrate towards the pharynx, undergo the first moult, leave this gallery and become attached to the mucosa of the pharynx, especially under the epiglottis. They stay in this position only for a short time and then migrate into the stomach where they are found attached by means of their mouth hooks to the mucosa of the cardiac region. The larvae undergo a second moult and pass into the third stage which is usually reached within 45-60 days after the first moult. The larva grows rapidly and full-grown larvae are found already between October and November. They remain, however, in a quiescent stage at least 7-8 months, until May or June, and during this long period of quiescent life they feed very little and their respiratory activity is at its lowest. The larvae finally leave the alimentary tract, burrow themselves into the soil or crevices in the ground, and pupate. The adult fly emerges from the pupa within 20-30 days.'

During the first period of life until the larva reaches the stomach its respiratory requirements are well satisfied by the rich oxygen supply in the surrounding medium. During the next 7-8 months of life within the stomach of the host the larva is

immersed in a semi-fluid medium deficient in oxygen and depends on an intermittent contact with air bubbles mixed with ingested food. The respiratory system of the larva is, nevertheless, highly developed and specialized for the aerobic mode of life. It exhibits several important features which enable the larva to overcome the difficulties inherent in its peculiar surroundings. One of those features is the presence of an intracellular haemoglobin.

The young larvae of *Gastrophilus* are of a uniformly red colour which, owing to the great transparency of the cuticle, shows that the haemoglobin is not restricted to one kind of tissue. In fact, on dissection the larva shows the presence of haemoglobin in the whole of the fat body, parietal musculature and hypoderm. As the larva grows the haemoglobin becomes localized in special cells forming a large red mass almost entirely occupying the posterior third of the body. These cells, which are known as tracheal cells, form an integral part of the respiratory or tracheo-spiracular system of the larva. Their distribution in the body, their connexion with the tracheal system, their structure and development were dealt with in papers by Enderlein (1899), Prenant (1900), Portier (1911), Kemnitz (1916), Dinulescu (1932) and Keilin (1944). Of the five pairs of tracheae which originate from postabdominal spiracles, four pairs are exceptionally wide at their base but rapidly decrease in diameter, thus acquiring a distinctly conical shape. All along their surface they give off numerous branches of smaller tracheae which ramify and finally break up into numerous tufts of intraprotoplasmic tracheoles filling the large tracheal cells (Fig. 1 A, B). The latter vary somewhat in size but the majority of them reach about 350-400 μ . in diameter, thus being visible to the naked eye. These cells are of a distinctly red colour owing to the presence of oxyhaemoglobin and are known as haemoglobin or tracheal cells. They probably originate from fat-body cells by gradual differentiation and invasion by tracheae (Dinulescu, 1932). In addition to haemoglobin, these cells are