of rat-serum γ -globulin. Fluorescent antibody studies revealed that the antiserum against lysine-rich rat-liver histone reacted with the contents of the nuclei of cryostat sections or of cell-suspension smears of rat liver (Plate 1).

Different results were obtained with the antiserum against lysine-poor rat-liver histones (fraction 2a + 3). In the first place this antiserum did not react with lysine-rich rat-liver histone but reacted weakly with one or two rat serum components. After sufficient absorption with rat-serum powder, the antiserum revealed two lines with rat-liver histone fraction 2a + 3, and the same two lines with liver homogenate. Both lines were also observed with separate fraction 2a but only one with fraction 3. This indicated that histone fractions 2a and 3 were immunologically different from each other, and that fraction 2a was contaminated with fraction 3. Immunoelectrophoretically the two lines were localized towards the cathode, where they crossed each other completely.

Our finding that rat-liver homogenates, prepared in 0.15M-NaCl, reacted with antisera against rat-liver histones suggests that part of the histone of rat liver dissociates from DNA after homogenization of the tissue in 0.15M-NaCl. The possibility that

saline-soluble protein contaminating our histone preparations had acted as antigen seems unlikely, since the histones were prepared from rat liver after removal of saline-soluble material (Sluyser *et al.* 1965). Immunoelectrophoresis also showed that the antigen had a basic character and fluorescent antibody studies showed that it was located in rat-liver nuclei.

It can thus be concluded that: (1) rat-liver histones, prepared according to our procedure, are antigenic in the rabbit; (2) rat-liver histone fractions 1, 2a and 3 are immunochemically distinct.

The authors thank Miss T. Bos and Miss M. L. Janse for expert technical assistance.

- Black, M. M., Ansley, H. R. & Mandl, R. (1964). *Arch. Path.* **78**, 350.
 Freund, J. (1947). *Annu. Rev. Microbiol.* **1**, 291.
 Scheidegger, J. J. (1955). *Int. Arch. Allergy*, **7**, 103.
 Sluyser, M., Thung, P. J. & Emmelot, P. (1965). *Biochim. biophys. Acta*, **108**, 249.
 Stedman, E. & Stedman, E. (1950). *Nature, Lond.*, **166**, 780.
 Weller, T. H. & Coons, A. H. (1954). *Proc. Soc. exp. Biol., N. Y.* **86**, 789.
 Wilson, R. H., Jurevics, V., Delaney, R., Ritter, T., Rencz, K. & Schram, A. (1966). *Nature, Lond.*, **209**, 1102.

Biochem. J. (1966) **101**, 20

Isolation of 5-Amino-4-ribitylaminoouracil from a Riboflavineless Mutant of *Aspergillus nidulans*

By J. SADIQUE, RADHA SHANMUGASUNDARAM and E. R. B. SHANMUGASUNDARAM
Biochemistry Research Laboratory, University of Madras, India

(Received 1 July 1966)

Studies to elucidate the mode of the formation of riboflavine in the riboflavine-synthesizing organism *Ermothecium ashbyii* (Goodwin & Treble, 1957) have indicated that a pyrimidine compound, 4,5-diaminoouracil, is a probable precursor in flavinogenesis. This observation has been extended by Katagiri, Takeda & Imai (1959) and Kishi, Asai, Masuda & Kuwada (1959), who showed that the ribityl derivative of 4,5-diaminoouracil is converted by extracts of *E. ashbyii* and *Aerobacter aerogenes* into 6,7-dimethyl-8-ribityl-lumazine, which has been reported to be the immediate precursor of riboflavine (Plaut, 1963).

Our observations on the utilization of the ribityl derivative of 4,5-diaminoouracil by the riboflavineless mutants of *Aspergillus nidulans* (Sadique,

Shanmugasundaram & Shanmugasundaram, 1966) also indicate that it can be a precursor of riboflavine. The identification of 4,5-diaminoouracil in a riboflavineless mutant of *Aspergillus nidulans* was reported by Sadique *et al.* (1966). The present communication deals with the isolation and identification of the ribityl derivative of 4,5-diaminoouracil in growing mycelia of a riboflavineless mutant of *Aspergillus nidulans*.

A riboflavineless mutant designated as wribo₆ was grown in 2l. of the minimal medium, as used by Pontecorvo (1953), in Hoffkin flasks supplemented with riboflavine. After 7 days' growth at 37° the mycelia were harvested. About 100g. wet wt. of mycelia were homogenized with 50ml. of water in a cold room (3–4°) with a Servall Omnimixer. The

Plate 1. Staining of rat-liver nuclei with the indirect fluorescent antibody test. Incubation of a smear of a rat-liver cell suspension with antiserum to rat-liver histone fraction 1 (*a*) and with normal rabbit serum (*b*). Experimental details are given in the text.

proteins were denatured by heating the extract in a boiling-water bath for 3–5 min. and then precipitated completely at pH 2.5 by adding 25 g. of $(\text{NH}_4)_2\text{SO}_4$. The precipitated proteins were centrifuged off and the flavines were extracted by the method of Crammer (1948), with phenol–ether–water (8:12:1, by vol.). The aqueous phase containing flavines were extracted with 10 ml. of benzyl alcohol to remove riboflavine and other lumazines and then with 5 ml. of chloroform to remove excess of benzyl alcohol used. The aqueous phase was finally passed through a Dowex 50 (X8) column equilibrated with 0.05N-HCl (Wacker, Harvey, Winestock & Plaut, 1964). The column was eluted with about 50 ml. of each of 0.05N- and 0.1N-HCl successively. The eluents were concentrated by pervaporation in the cold room at 3–4° and then chromatographed on Whatman no. 1 paper along with an authentic sample of 5-amino-4-ribitylaminouracil with butanol–acetic acid–water (4:1:5, by vol.) as solvent. A dark-brown spot detected under ultraviolet light with R_f 0.15 corresponded to the authentic sample of ribitylated uracil compound. The material eluted from the spot with 0.1N-HCl at 60° gave an ultraviolet spectrum similar to that of the authentic sample when examined in a Unicam SP.700 recording spectrophotometer. The eluent containing the ribitylated uracil was also subjected to electrophoresis on Whatman no. 1 paper (46 cm. $\times$ 4 cm.) with a solvent system containing 11.6 ml. of pyridine and 9.8 ml. of acetic acid at pH 4.8 (Plaut, 1963), a potential difference of 400v being applied for 3 hr. The isolated ribitylated compound moved 10.6 cm. towards the anode from the point of application, as detected under ultraviolet light, and, under these conditions, the authentic sample moved the same distance.

Further evidence for the formation of this ribitylated uracil compound was obtained by converting the isolated material into the DNP derivative by the method of Levy (1955). On paper chromatography the yellow DNP derivative of the isolated material had R_f 0.23, which was the same as that of the DNP derivative prepared from authentic 5-amino-4-ribitylaminouracil. About 1 mg. of the ribityl derivative of 4,5-diaminouracil was obtained from the mycelia harvested from 5 l. of medium.

The isolated compound was tested for growth-promoting activity with five riboflavineless mutants of *Aspergillus nidulans*. In auxanographic tests it promoted good growth of all the riboflavineless mutants, as did a sample of synthetic 5-amino-4-ribitylaminouracil. Thus the formation of 5-amino-4-ribitylaminouracil in the riboflavineless mutant and its growth-promoting activity establish its role in the biosynthesis of riboflavine.

We thank Dr Masuda for the generous gift of an authentic sample of 5-amino-4-ribitylaminouracil.

- Crammer, J. L. (1948). *Nature, Lond.*, **161**, 349.
 Goodwin, T. W. & Treble, D. H. (1957). *Biochem. J.* **67**, 10P.
 Katagiri, H., Takeda, I. & Imai, K. J. (1959). *J. Vitaminol. Japan*, **5**, 287.
 Kishi, J., Asai, M., Masuda, T. & Kuwada, S. (1959). *Chem. pharm. Bull., Tokyo*, **7**, 515.
 Levy, A. C. (1955). *Meth. biochem. Anal.* **11**, 360.
 Plaut, G. W. R. (1963). *J. biol. Chem.* **238**, 2225.
 Pontecorvo, G. (1953). *Advanc. Genet.* **5**, 14.
 Sadique, J., Shanmugasundaram, R. & Shanmugasundaram, E. R. B. (1966). *Naturwissenschaften*, **53**, 282.
 Wacker, H., Harvey, R. A., Winestock, C. H. & Plaut, G. W. E. (1964). *J. biol. Chem.* **239**, 3493.

Biochem. J. (1966) **101**, 3c

Multiple Branching in Glycogens

By G. N. BATHGATE and D. J. MANNERS

*Department of Brewing and Biochemistry, Heriot-Watt University,
 Edinburgh*

(Received 29 June 1966)

Although it is generally accepted that glycogen has a multiply branched 'tree-type' structure as originally suggested by Meyer (1943), formal proof of this is limited. The most convincing evidence arises from the enzymic studies of Larnier, Illing-

worth, Cori & Cori (1952), who examined the step-wise degradation of rabbit-liver and -muscle glycogen by phosphorylase and amylo-1,6-glucosidase. The results indicated that the branch points were arranged in tiers, and that approximately half