Rat-Urine Glycosidases and Kidney Damage

By D. ROBINSON, R. G. PRICE AND N. DANCE

Department of Nutrition, Queen Elizabeth College (University of London), London, W.8

(Received 13 June 1966)

1. The activities of β -galactosidase, β -glucosidase, β -glucuronidase and *N*-acetyl- β -glucosaminidase were estimated in normal and pathological rat urine, with 4-methylumbelliferyl glycosides as substrates. 2. Kidney damage induced by injections of uranium nitrate, mercuric chloride, potassium dichromate or 4-nitrophenylarsonic acid causes a marked increase in the urinary excretion of all four enzymes. 3. The rise in β -glucosidase activity was associated with the appearance of a new urinary enzyme species, which was examined by starch-gel electrophoresis, DEAE-cellulose chromatography and filtration on Sephadex G-75 and G-200. 4. This enzyme appears to be identical with its counterpart in the kidney, and it is suggested that it arises in the urine as a result of renal tubular breakdown. 5. The other glycosidases examined also show some physical similarities to the corresponding enzymes of the rat kidney.

In the preceding paper (Robinson, Price & Dance, 1967) some characteristics of rat-kidney β -galactosidase (β -D-galactoside galactohydrolase, EC 3.2.1.23), β -glucosidase (β -D-glucoside glucosylhydrolase, EC 3.2.1.21), β -glucuronidase (β -D-glucuronide glucuronohydrolase, EC 3.2.1.31) and *N*-acetyl- β -glucosaminidase (β -2-acetamido-2-deoxy-D-glucoside acetamidodeoxyglucosylhydrolase, EC 3.2.1.30) were described. The occurrence of β -glucuronidase in rat urine has been demonstrated by Mead, Smith & Williams (1955), and Furth & Robinson (1965) found similarities between rat-urine and rat-kidney β -galactosidases. Some urinary enzymes have found diagnostic applications because of the variations in amount excreted under pathological conditions (Long, 1961; Amador, Zimmerman & Wacker, 1963). These variations may arise from serum enzymes by glomerular leakage (Wilding, 1965) or by cellular breakdown in the kidney (Rosalki & Wilkinson, 1959), or other parts of the uro-genital tract. Urinary alkaline phosphatase is increased after experimental kidney damage by X-rays (Asscher & Anson, 1960) or uranium nitrate (Breedis, Flory & Furth, 1943). In the latter case Alibert, Rigal, Bastide, Turchini & Dastugue (1963) showed that there was a concurrent decrease in the alkaline phosphatase (orthophosphoric monoester phosphohydrolase, EC 3.1.3.1) of the kidney.

In the present work, kidney damage in rats was induced by mercuric chloride, uranium nitrate and potassium dichromate, the histological effects of which have been described (Tapp, Carrol & Kovacs, 1965), and by 4-nitrophenylarsonic acid, a substance

used in poultry feeds as a growth promoter and therapeutic agent (Moody & Williams, 1964). The nephrotoxicity of this substance in mammals has not been previously reported. We present evidence for the excretion of elevated amounts of β -galactosidase, β -glucosidase, β -glucuronidase and *N*-acetyl- β -glucosaminidase in the urine of rats as a consequence of kidney damage by the above agents. Enzyme activities are low in normal urine, but can be readily estimated fluorimetrically, and for β -glucuronidase the urine can be diluted before assay sufficiently to eliminate the effect of naturally occurring inhibitors (Marsh, 1963) without need for dialysis.

EXPERIMENTAL

Materials. Glycosidase substrates and inhibitors were obtained as described in the preceding paper (Robinson *et al.* 1967). All salts used for buffer solutions and kidney poisons were A.R. grade (British Drug Houses Ltd., Poole, Dorset) and 4-nitrophenylarsonic acid was a gift from the Salsbury Laboratories, Charles City, Iowa, U.S.A.

Urine collection. Adult male Sprague-Dawley rats were maintained in metabolism cages over silicone-treated funnels. The 24hr. samples were collected in tubes maintained at 4° to minimize enzyme inactivation. Because bran is known to have high β -galactosidase activity (Wohnlich, 1963) feeding was restricted to 1 hr. each day and care was taken to avoid contamination of the urine with food particles. There was no restriction on drinking water. Urine samples were filtered through Whatman no. 54 paper and made up to 50 ml. with distilled water before assay.

Urine powder. Rat urine (1250 ml.) was concentrated by dialysis under vacuum against 0.04M-sodium phosphate buffer, pH 7.0 (Huehns & Shooter, 1961). The urine concentrate was then freeze-dried to a light-brown powder

(2.7 g.), which retained 78% of the β -galactosidase activity of the original urine and was a convenient enzyme source for further characterization. This powder is stable for many weeks at -10° but rapidly denatures at more elevated temperatures or in dilute solutions.

DEAE-cellulose and Sephadex G-75 and G-200 columns. These were prepared and used as described in the preceding paper (Robinson *et al.* 1967).

Gel electrophoresis. After quantitative assays had been performed the urine from experimental animals was concentrated 40–50-fold under vacuum and subjected to starch-gel electrophoresis. The conditions for electrophoresis and the technique of identification of glycosidase bands are described in the preceding paper (Robinson *et al.* 1967).

Enzyme assays. All glycosidases were estimated fluorimetrically as described in the preceding paper (Robinson *et al.* 1967), with suitably diluted urine samples.

K_m values. These were calculated by the double-reciprocal plot method of Lineweaver & Burk (1934).

pH optima. These were determined in sodium phosphate-citrate (McIlvaine, 1921) and in 0.1 M-acetate buffers.

Protein. This was measured by the extinction at 280 m μ .

Chloride. This was assayed by the method of Sanderson (1952).

Nephrotoxic agents. All materials were injected as solutions in 1 ml. of sterile water adjusted to pH 7.5 by the addition of NaHCO₃. 4-Nitrophenylarsonic acid (5 mg.) and uranium nitrate (5 mg.) were administered intraperitoneally, and mercuric chloride (1 mg.) and potassium dichromate (5 mg.) were given subcutaneously to rats of 150–200 g. body weight.

RESULTS

Properties of glycosidases

pH optima and effects of buffers. Fig. 1 shows that *N*-acetyl- β -glucosaminidase and β -glucosidase are not greatly affected by the buffer system used and have pH optima about 5.0–5.5, but β -glucuronidase is less active in phosphate-citrate buffer than in acetate, and the pH optimum shifts from 4.5 in acetate to 5.5 in phosphate-citrate. This inhibition and apparent change in optimum pH has been described by Mills, Paul & Smith (1953) for ox spleen and by Woollen & Walker (1965) for human-plasma β -glucuronidase.

In acetate buffer, β -galactosidase shows two optima at pH 5.0 and at 3.5, whereas in phosphate-

Fig. 1. Effect of pH on urinary β -glycosidase activities in citrate-phosphate (solid symbols) or acetate (open symbols) buffers. Enzymes were assayed by the fluorimetric methods described in the text. (a) β -glucosidase; (b) β -galactosidase; (c) *N*-acetyl- β -glucosaminidase; (d) β -glucuronidase.

Table 1. *Estimation of β -glucuronidase, N-acetyl- β -glucosaminidase, β -galactosidase and β -glucosidase in freshly voided normal rat urine and in freeze-dried non-diffusible material*

Activities were measured under optimum conditions as described in the text and expressed as μ moles of the appropriate 4-methylumbelliferyl glycoside hydrolysed/hr.

Enzyme	Activity/mg. of urine powder	Activity/24 hr. urine sample	K_m (mM)
β -Glucosidase	0.008	0.017 ± 0.006 (11)	0.27
β -Galactosidase	0.16	2.10 ± 0.49 (20)	0.16
N-Acetyl- β -glucosaminidase	0.10	1.03 ± 0.66 (20)	0.47
β -Glucuronidase	0.40	4.09 ± 2.90 (18)	0.02

citrate buffer a single optimum pH of 4.0 is obtained.

Normal urine activities. Average values for the daily excretion of the four β -glycosidases are shown in Table 1 along with the activities found in urine powder. β -Glucosidase only occurs in trace amounts in urine in comparison with the other three enzymes, having less than one-hundredth of the activity of β -galactosidase. In the urine powder the relative activity of β -glucosidase to β -galactosidase is increased fivefold, and since the recovery of the latter was 78% during the process of concentration and freeze-drying it may be that the elevated activities of β -glucosidase represent an activation or the removal of an inhibitor.

DEAE-cellulose chromatography. Urine powder was fractionated on a DEAE-cellulose column with a chloride gradient elution (Fig. 2). Each enzyme appears to contain at least two components. In each case a major component is not retained by the column under these conditions. Similar results have been described for β -galactosidase by Furth & Robinson (1965). The individual peaks were concentrated and rechromatographed on fresh columns to ensure that they did not arise from overloading or as artifacts produced during the chromatographic run. Recoveries of 50–70% of the applied enzyme activity were obtained.

Gel filtration. A separation of the β -glucuronidase, N-acetyl- β -glucosaminidase and β -galactosidase activities of rat-urine powder was obtained on a Sephadex G-200 column (Fig. 3), but the β -glucosidase activity could not be separated from the N-acetyl- β -glucosaminidase and β -galactosidase fractions. Estimates of molecular weights obtained by separation on a column that had been calibrated with proteins of known molecular weights as described by Andrews (1965) were: β -glucuronidase 200 000–210 000; N-acetyl- β -glucosaminidase, 130 000–140 000; β -galactosidase, 80 000.

Gel electrophoresis. The β -glucosidase activity was too weak to be detectable on the gels by the methods employed. Of the other three enzymes β -glucuronidase and N-acetyl- β -glucosaminidase were negatively charged at pH 7.0 and gave diffuse bands near the origin whereas the β -galactosidase

Fig. 2. Chloride-gradient elution from a DEAE-cellulose column of β -glycosidases of rat-urine powder. The column (20 cm. $\times$ 1 cm.) was equilibrated with 0.01 M-phosphate buffer, pH 6.0, and eluted with an NaCl gradient (in the same buffer) up to 0.3 M. Each 3 ml. fraction was assayed for the four β -glycosidases and for protein and chloride as described in the text. (a) β -Glucosidase; (b) β -galactosidase; (c) N-acetyl- β -glucosaminidase; (d) β -glucuronidase; (e) $E_{280\text{m}\mu}$ (—) and chloride gradient (·····).

activity moved more rapidly towards the anode and appeared somewhat faster than the slow-moving bands of kidney β -galactosidase.

Effects of kidney damage

There was a marked rise in the urinary excretion of all four enzymes after injections of uranium nitrate, mercuric chloride, potassium dichromate or 4-nitrophenylarsonic acid. The changes in enzyme activities during one such experiment are shown in Fig. 4, and typical experiments with each of the four compounds are compared in Table 2. All the substances produced similar transient increases, reaching a peak on the first or second day after injection. In each case the urine volumes increased and proteinuria was detected, but in non-fatal experiments the enzyme activities subsequently returned to near-normal values. A notable feature is the relative increase in β -glucosidase, which rises from barely detectable levels in the normal animal to substantial values after kidney damage. The rise in β -glucuronidase was occasionally delayed by up to 24 hr. in comparison with the peak excretion of the other enzymes. The rats were killed on the

eighth day and the kidneys were found to be greatly enlarged and discoloured.

Gel electrophoresis. A comparison of the gel-electrophoresis patterns of the enzymes from normal and pathological urines showed that the increase in activity of β -glucosidase is due to the appearance of a new species, which is characterized by its rapid anodic mobility and by the fact that it appears to carry both β -glucosidase and β -galactosidase activity (Fig. 5). Development of the band on the gel was completely inhibited by incorporating glucono-(1 $\rightarrow$ 5)-lactone or galactono-(1 $\rightarrow$ 4)-lactone into the substrate solutions used irrespective of whether β -glucosidase or β -galactosidase was being examined. Its electrophoretic mobility was identical with that of the β -glucosidase enzyme of the kidney, reported in the preceding paper (Robinson *et al.* 1967). Apart from an increase in intensity of the bands due to the higher enzyme activities no marked differences were observed in the electrophoretic behaviour of the other enzymes examined in the pathological specimens when compared with normal samples.

Gel filtration. When a sample of abnormal urine was filtered through a Sephadex G-75 column (Fig. 6) the β -glucosidase activity could be clearly

Fig. 3. Gel filtration of rat-urine-powder β -glycosidases on Sephadex G-200 (200–300 mesh). The column (30 cm. $\times$ 3 cm.) was prepared in 0.01M-sodium phosphate buffer, pH 7.6, containing 0.4M-NaCl and 0.01M-EDTA. The sample (15 mg.) was dissolved in 2 ml. and run in the same buffer at 4° at a flow rate of 12 ml./hr. Samples of the 3 ml. fractions collected were suitably diluted and assayed as described in the text. The void volume (57 ml.) was determined by using Blue Dextran 2000 and the internal volume (111 ml.) by using sucrose. Assays were carried out in citrate-phosphate buffer (solid symbols) or in acetate buffer (open symbols). $\bullet$, β -Glucosidase; $\blacksquare$, β -galactosidase; $\blacktriangle$, *N*-acetyl- β -glucosaminidase; ∇ , β -glucuronidase.

Fig. 4. Daily excretion of β -glycosidases by a rat after a single dose of 5 mg. of 4-nitrophenylarsonic acid on day 2. Units of activity are as given in Table 2, and symbols for β -glycosidases are as given in Fig. 3. The broken line indicates the volume of the 24 hr. urine samples.

Table 2. *Effect of drugs on excretion of β -glycosidases in rat urine*

Results are expressed as units of activity excreted in the 24hr. before injection and on that day after dosing when the maximum rise occurs. One unit of activity is defined as that amount of enzyme that will hydrolyse 1 μ mole of the appropriate 4-methylumbelliferyl β -glycoside/hr. under the standard conditions of assay described in the text.

Nephrotoxic agent		Total activity excreted in a 24hr. urine sample			
		β -Galactosidase	β -Glucosidase	<i>N</i> -Acetyl- β -glucosaminidase	β -Glucuronidase
4-Nitrophenylarsonic acid (5mg. intraperitoneally)	Before injection	1.56	0.0075	1.2	7.5
	After injection	22.25	4.17	23.4	62.5
Mercuric chloride (1mg. subcutaneously)	Before injection	1.55	0.0014	1.45	0.75
	After injection	3.10	0.810	5.13	3.1
Potassium dichromate (5mg. subcutaneously)	Before injection	2.35	0.015	1.95	8.75
	After injection	6.09	0.840	13.8	21.1
Uranium nitrate (5mg. intraperitoneally)	Before injection	1.46	0.020	1.0	2.61
	After injection	3.75	0.570	9.75	17.2

Fig. 5. Starch-gel electrophoresis of β -galactosidase in rat-urine concentrate before and after dosing on day 2 with 4-nitrophenylarsonic acid. Separation was carried out in 0.04M-sodium phosphate buffer, pH7.0, for 3hr. at 4°. Bands were developed by flooding with 1mM-4-methylumbelliferyl β -galactoside in phosphate-citrate buffer, pH3.7, and photographed under u.v. light. The rapidly migrating band also showed all the β -glucosidase activity.

separated from the bulk of the β -galactosidase activity. The eluate samples containing the peaks of activity were concentrated and subjected to starch-gel electrophoresis, when it was found that the β -glucosidase peak contained all the fast-moving β -galactosidase activity (cf. Fig. 5) and none of the slow-moving β -galactosidase components. The latter appeared in the concentrates of the main β -galactosidase peak. Molecular-weight estimations on Sephadex G-200 gave results similar to those obtained with normal urine powder for β -glucuronidase, *N*-acetyl- β -glucosaminidase and β -galactosidase. The β -glucosidase had a molecular weight of 40000, which is the same as that of the corresponding kidney enzyme.

DEAE-cellulose. A further identification of the β -glucosidase component of the pathological urines

Fig. 6. Separation on Sephadex G-75 of β -galactosidase and β -glucosidase in rat urine collected 48hr. after injection of 5mg. of 4-nitrophenylarsonic acid. Details of column procedure (V_0 51 ml., V_i 75 ml.) and assays are as described for Fig. 3. ●, β -Glucosidase; ■, β -galactosidase.

with that normally found in the kidney was achieved by a stepwise-elution experiment on DEAE-cellulose. Whereas in normal urine the only detectable β -glucosidase is that which is not retained by the column, the bulk of the activity appearing after injection of 4-nitrophenylarsonic acid consisted of a component that was firmly bound to the ion-exchanger and required 0.15M-chloride for its displacement.

DISCUSSION

Except for amylase in pancreatitis (Budd, Walter, Harris & Knight, 1959) and β -glucuronidase

in bladder-cancer (Boylard, Wallace & Williams, 1955) measurement of the urinary excretion of glycosidases has so far not found clinical use. One reason for this is, no doubt, the very low enzyme activities that occur and therefore the highly sensitive methods required for their detection. For diagnosis it is important that the origin of the urinary enzyme is clearly recognized. To do this, the physical and biochemical characteristics of specific tissue and urine enzymes must be compared. A combination of the fluorescence methods of assay with characterization of the enzyme forms by gel electrophoresis suggests here that the new enzyme species appearing in the urine after dosage with nephrotoxic agents originated in the kidney cells, and not by serum filtration through damaged glomeruli. The β -glucosidase component of the pathological urines has the characteristics of the kidney enzyme and has no counterpart in the spleen, liver, epididymis or intestine, which are the other major sites of activity in rat tissues. Histochemical examination of the kidney shows most intense β -galactosidase and β -glucosidase activity in the epithelial cells of the proximal and distal tubules (Rutenburg, Rutenburg, Monis, Teague & Seligman, 1958). Severe necrosis of the proximal tubular epithelium is caused by mercuric acetate and potassium dichromate, whereas the effect of uranium nitrate is to produce generalized tubular necrosis (Tapp, *et al.* 1965). There is thus strong circumstantial evidence for the elevated activities of β -glucosidase and β -galactosidase being an indicator of tubular breakdown.

It is noteworthy that in their examination of the alkaline phosphatase of human urine Butterworth, Moss, Pitkanen & Pringle (1965) found a low-molecular-weight component, which bore a close similarity to that present in the cells excreted after exfoliation by aspirin. They were, however, unable to show such a component in kidney extracts and considered it might be a minor contribution to the total kidney alkaline-phosphatase activity. In the present case, however, a component closely resembling the abnormal urinary β -glucosidase is found in crude kidney extracts. It has a similar

optimum pH, the same low molecular weight, identical behaviour on DEAE-cellulose chromatography and starch-gel electrophoresis and is active towards both β -glucosides and β -galactosides. This kidney enzyme accounts for all the β -glucosidase and 15–20% of the β -galactosidase of that organ.

REFERENCES

- Alibert, A., Rigal, M., Bastide, P., Turchini, J. & Dastugue, G. (1963). *C.R. Acad. Sci., Paris*, **157**, 1439.
Amador, E., Zimmerman, T. S. & Wacker, W. E. C. (1963). *J. Amer. med. Ass.* **185**, 769.
Andrews, P. (1965). *Biochem. J.* **96**, 595.
Asscher, A. W. & Anson, S. G. (1960). *Lancet*, **i**, 1109.
Boylard, E., Wallace, D. M. & Williams, D. C. (1955). *Brit. J. Cancer*, **9**, 62.
Breedis, C., Flory, C. M. & Furth, J. (1943). *Arch. Path.* **36**, 402.
Budd, J. J., Walter, K. E., Harris, M. L. & Knight, W. A. (1959). *Gastroenterology*, **36**, 333.
Butterworth, P. J., Moss, D. W., Pitkanen, E. & Pringle, A. (1965). *Clin. chim. Acta*, **11**, 220.
Furth, A. J. & Robinson, D. (1965). *Biochem. J.* **97**, 59.
Huehns, E. R. & Shooter, E. M. (1961). *J. molec. Biol.* **3**, 257.
Lineweaver, H. & Burk, D. (1934). *J. Amer. chem. Soc.* **56**, 658.
Long, C. (1961). *Biochemists' Handbook*, p. 931. London: E. and F. N. Spon Ltd.
McIlvaine, T. C. (1921). *J. biol. Chem.* **40**, 183.
Marsh, C. A. (1963). *Biochem. J.* **86**, 77.
Mead, J. A. R., Smith, J. N. & Williams, R. T. (1955). *Biochem. J.* **61**, 569.
Mills, G. T., Paul, J. & Smith, E. E. B. (1953). *Biochem. J.* **53**, 232.
Moody, J. P. & Williams, R. T. (1964). *Fd Cosmet. Toxicol.* **2**, 695.
Robinson, D., Price, R. G. & Dance, N. (1967). *Biochem. J.* **102**, 525.
Rosalki, S. B. & Wilkinson, J. H. (1959). *Lancet*, **ii**, 327.
Rutenburg, M., Rutenburg, S. H., Monis, B., Teague, R. & Seligman, A. M. (1958). *J. Histochem. Cytochem.* **6**, 122.
Sanderson, P. H. (1952). *Biochem. J.* **52**, 502.
Tapp, E., Carrol, R. & Kovacs, K. (1965). *Arch. Path.* **79**, 629.
Wilding, P. (1965). *Clin. chim. Acta*, **8**, 918.
Wohnlich, J. (1963). *Bull. Soc. Chim. biol., Paris*, **45**, 1171.
Woollen, J. W. & Walker, P. G. (1965). *Clin. chim. Acta*, **12**, 659.