EXPLANATION OF PLATE I

Electron micrographs of cell-wall fraction of *B. polymyxa* shadowed with gold-palladium: (a) wall from 31 hr. culture; (b) walls from 18 hr. culture; (c) and (d) material extracted with urea, dialysed and freeze-dried. The scale marks represent 0.25μ .

Short Communications

The Structure of the Cell Wall of *Bacillus polymyxa* (N.C.I.B. 4747)

By J. GOUNDRY, A. L. DAVISON, A. R. ARCHIBALD and J. BADDILEY
Microbiological Research Laboratory, Department of Organic Chemistry,
University of Newcastle upon Tyne

(Received 14 April, 1967)

The presence of a regular fine-structure has been reported (Baddiley, 1964) in the cell wall of the Gram-variable bacterium *Bacillus polymyxa*. Electron micrographs (Plate 1a) show a regular rectangular packing of particles of about 75 Å diameter. A similar type of fine-structure has previously been observed in the walls of an unidentified Gram-positive bacterium (Labaw & Mosley, 1954). Related fine-structure with hexagonal packing has been found in the walls of several Gram-negative bacteria (Murray, 1963; van Iterson, 1954; Salton & Williams, 1954) and also in *Micrococcus radiodurans* (Glauert, 1962). *B. polymyxa* was grown in yeast-peptone-water [yeast extract (Difco), 0.5%; peptone (Oxoid), 1.0%; NaCl (A.R.), 0.5%] at 30° with stirring and aeration (5l. of air/min. for 10l. of medium) and cells were harvested after 5, 18 and 31 hr., giving cells showing respectively positive, mixed and negative reaction to the Gram stain. The cells were disintegrated in a Shockman shaker head and walls were collected by centrifugation, washed with water and freeze-dried.

Walls from all three batches of organism show the same type of structure, although this was partially obscured in many of the walls prepared from the 18 hr. culture (Plate 1b). Treatment of these walls (100mg.) with chloroform-methanol (2:1, v/v) (50ml.) for 1 hr. at 20° removed lipid (10mg.), which consisted mainly of neutral lipid, phosphatidylethanolamine and phosphatidylglycerol. The amino acid composition of the wall was unaffected by this extraction, but electron microscopy now showed regular structure on all walls, indicating that in the original sample the component obscuring the regular structure was lipid. On the other hand, detergents removed the component responsible for the appearance of regular structure. Walls (100mg.) from the 31 hr. culture were suspended in 1% (w/v) sodium dodecyl sulphate (50ml.) for 1 hr. at room temperature. The residual walls were then recovered by centrifugation, washed with water and freeze-dried (yield, 32.7mg.). Wall residue (3mg.) and extract (2ml.) were heated at 100° for 16 hr. *in vacuo* in 6N-HCl (3ml.). The HCl was removed *in vacuo* over KOH and samples of the hydrolysate

were analysed for amino acids with a Technicon amino acid analyser.

Wall residue (5mg.) and extract (4ml.) were also hydrolysed for 3 hr. at 100° in 2N-HCl and the dried hydrolysates were examined for sugars by paper chromatography in butan-1-ol-pyridine-water (6:4:3, by vol.). The detergent removed 67% of the original weight of the wall, and electron microscopy showed that the residual wall material was completely devoid of regular structure and, although it had the same shape as the original walls, it was much thinner.

The residual material contained typical components of mucopeptide and was susceptible to the action of lysozyme; it is concluded that this material is the rigid mucopeptide component of the wall. The extract (Table 1) contained almost all of the aspartic

Table 1. Amino acid composition of wall samples

	Amino acid composition (μmoles/g. of original wall)		
	31 hr. cell walls	Residue after detergent treatment	Detergent extract
Aspartic acid	553	33	519
Threonine and serine	754	21	683
Muramic acid	191	171	0
Glutamic acid	412	171	264
Proline	*	0	*
Glycine	352	15	341
Alanine	623	259	396
Glucosamine	211	213	8
Methionine	20	2	23
Diaminopimelic acid	161	140	Trace
Leucine	272	11	255
Tyrosine	100	11	96
Phenylalanine	151	5	136
Lysine	322	30	246
Histidine	22	3	21

* Present but not estimated. In addition, substances tentatively identified as valine and galactosamine were present in each sample.

acid and lysine of the walls but little or no diaminopimelic acid. Hydrolysis of the extract gave glucose, identified with Glucostat reagent, and two or three other sugars, which have not yet been identified. No structure was detected after treatment of walls (5mg.) with papain (2mg. in 5ml. of 0.05M-phosphate buffer, pH 7.0) or with trypsin (2mg. in 5ml. of 0.05M-phosphate buffer, pH 7.0).

Walls (100mg.) from the 31hr. culture were suspended in 6M-urea (50ml.) at 20° for 1 hr. and then recovered by centrifuging as before. The urea extract was dialysed against distilled water and freeze-dried. Electron microscopy of the residual walls showed that some but not all of the structure had been removed. The extract contained thin layers or strings of structural material (see Plates 1c and 1d). Prolonged extraction (20° for 30hr.) caused complete disintegration of the walls.

Walls (77mg.) were suspended in water (100ml.) at 100° for 25min. They were then centrifuged and freeze-dried; these 'boiled' walls (71mg.) did not show fine structure. A sample (23.7mg.) was stirred with 6M-urea solution (12ml.) at 20° for 1 hr. The remaining walls were centrifuged and washed with water (yield, 12mg.); areas possessing regular structure were observed on some of these walls.

These observations show that, contrary to the earlier suggestion (Baddiley, 1964; cf. Martin, 1966), the structured component of the cell wall is not a

typical mucopeptide. The removable material is rich in aspartic acid and lysine and contains little or no muramic acid, glucosamine or diaminopimelic acid. It is not yet known whether the neutral sugars present in extracts are associated with the protein or polypeptide, or whether these are present in a separate polysaccharide component. The disappearance of regular structure on boiling, and its subsequent partial reappearance on brief treatment with urea, suggest that heating may cause reversible changes in the arrangement of the structured component.

Murray & Nermut (personal communication) have recently examined the structured layer in strains of *B. polymyxa* and will describe their results elsewhere.

We thank Mr E. H. Boulton for electron microscopy. The work was supported by grants from the Science Research Council and Glaxo Laboratories Ltd.

Baddiley, J. (1964). *Endeavour*, **23**, 33.

Glauert, A. M. (1962). *Brit. med. Bull.* **18**, 245.

Labaw, W. & Mosley, V. M. (1954). *Biochim. biophys. Acta*, **15**, 325.

Martin, H. H. (1966). *Annu. Rev. Biochem.* **35**, 457.

Murray, R. G. E. (1963). *Canad. J. Microbiol.* **9**, 381.

Salton, M. R. J. & Williams, R. C. (1954). *Biochim. biophys. Acta*, **14**, 455.

van Iterson, W. (1954). *Proc. int. Congr. Electron Microscopy, London*, p. 602.

Biochem. J. (1967) **104**, 20

Glycerol Kinase Activities in Rat Heart and Adipose Tissue

By JANICE ROBINSON and E. A. NEWSHOLME*

*Medical Research Council Unit for Research in Cell Metabolism,
Department of Biochemistry, University of Oxford*

(Received 19 April 1967)

Wieland & Suyter (1957) reported that glycerol kinase (ATP-glycerol phosphotransferase, EC 2.7.1.30) was absent from adipose tissue and heart muscle of the rat. These authors assayed glycerol kinase activity by coupling to α -glycerophosphate dehydrogenase and measuring NAD reduction at pH 9.5. During studies on glycerol kinase from rat liver the *in vitro* half-life of this enzyme at 25° was found to be only 6.5 min. at pH 9.8, compared with 3.5 hr. at pH 7.5. It therefore seemed possible that

* Present address: Agricultural Research Council Unit of Insect Physiology, Department of Zoology, University of Oxford.

failure to demonstrate even a low activity of glycerol kinase in adipose tissue and muscle might have been due to rapid inactivation by the high pH of the assay conditions. Recently Newsholme, Robinson & Taylor (1967) described a sensitive radiochemical technique for measuring the activity of glycerol kinase, and this method has now been used to demonstrate the presence of this enzyme in rat adipose tissue and heart.

Rat hearts and epididymal fat pads were cut into small pieces and homogenized in 2 vol. of 1% (w/v) KCl and 1 mM-EDTA at 2° in a Silverson homogenizer (Silverson Machines Ltd., London, S.E. 1)