between corneal and skeletal keratan sulphates (Mathews & Cifonelli, 1965).

The immunological characteristics of the keratan sulphate-protein complex and of the pooled supernatants from the hyaluronidase digestion were entirely different. With antisera to the starting proteoglycan preparation (Loewi & Muir, 1965), only the pooled supernatants contained the several antigens of the original material, none of which was revealed in the keratan sulphate-protein by double-diffusion at various dilutions. Nevertheless, antisera to the keratan sulphate-protein complex were successfully raised in rabbits, and showed no cross-reaction with any of the antigens in the pooled supernatants. The keratan sulphate-protein complex was thus immunologically distinct from these antigens, and furthermore appeared to be a single antigen on double-diffusion and immuno-electrophoresis. In the original complex, this protein to which both keratan sulphate and some chondroitin sulphate is attached is therefore different from the proteins to which most of the chondroitin sulphate is bound. Since the latter comprise the greater part of the protein in the whole complex, they are responsible for its overt antigenic properties. The keratan sulphate-protein complex, although a minor component, appears to be more prevalent in larger complexes since these contain more glucosamine than smaller ones. The

keratan sulphate-protein complex may therefore either vary greatly in size or be in some way involved building in up larger molecules.

We are grateful to Dr B. Högberg, A. B. Leo, Halsingborg, Sweden, and Professor A. G. Lloyd for purified hyaluronidase, and to Dr R. Pain for sedimentation experiments. The competent technical assistance of Mr R. Ewins and the support of C. P. T. by the Nuffield Foundation is gratefully acknowledged.

- Anderson, B., Hoffman, P. & Meyer, K. (1965). *J. biol. Chem.* **240**, 156.
 Gregory, J. D. & Rodén, L. (1961). *Biochem. biophys. Res. Commun.* **5**, 430.
 Hoffman, P., Meyer, K. & Linker, A. (1956). *J. biol. Chem.* **219**, 653.
 Lindahl, U. & Rodén, L. (1964). *Biochem. biophys. Res. Commun.* **17**, 254.
 Loewi, G. & Muir, H. (1965). *Immunology*, **9**, 119.
 Mathews, M. B. & Cifonelli, J. A. (1965). *J. biol. Chem.* **240**, 4140.
 Muir, H. (1958). *Biochem. J.* **69**, 195.
 Muir, H. & Jacobs, S. (1967). *Biochem. J.* **103**, 367.
 Partridge, S. M. & Elsdon, D. F. (1961). *Biochem. J.* **79**, 26.
 Rosenblum, E. L. & Cifonelli, J. A. (1967). *Fed. Proc.* **26**, 282.
 Seno, N., Meyer, K., Anderson, B. & Hoffman, P. (1965). *J. biol. Chem.* **240**, 1005.
 Tsiganos, C. P. & Muir, H. (1965). *Proc. int. Symp. Biochimie et Physiologie du Tissu Conjonctif, Lyon*, p. 27. Ed. by Comte, P.

Biochem. J. (1967) **104**, 28c

Variations in the C-Terminal Half of Immunoglobulin λ -Chains

By C. MILSTEIN

Medical Research Council Laboratory of Molecular Biology, Cambridge

(Received 17 May 1967)

The immunoglobulin light chains contain multiple variations of amino acid sequence localized in the N-terminal half of the molecule, whereas the C-terminal half is essentially invariant. This is true for κ -chains (Hiltschmann & Craig, 1965; Titani, Whitley, Avogardo & Putnam, 1965; Milstein, 1966a) and also for λ -chains (Milstein, Clegg & Jarvis, 1967). In κ -chains, however, one variant occurs at position 191 that can be correlated with the allotypic marker Inv. Thus Inv *a* chains contain leucine whereas Inv *b* chains contain valine in that position (Milstein, 1966a,b; Baglioni, Zonta, Cioli & Carbonara, 1966). Position 122 is still controversial

(discussed by Milstein, 1966c). The invariancy of the C-terminal half of light chains has not been surveyed very extensively. In κ -chains the C-terminal half of three proteins (Hiltschmann & Craig, 1965; Titani *et al.* 1965) and stretches around the cysteine residues of five more (Milstein, 1966c) have been studied in detail. 'Fingerprints' of a larger number of proteins have not established any other variation (Baglioni & Cioli, 1966). The sequences of the C-terminal half of two λ -proteins have been established independently in two different Laboratories and there is complete agreement in the two reported sequences with the exception of the

position 190 is indicated by the fact that it is present only when the arginine-containing peptide T3 is absent.

One of the proteins (Mz) contained two variants, one in peptide T8 (valine instead of alanine) and the other in peptide T4/T5 (see Fig. 1). The variant in peptide T8 was located at residue 143. This was shown by isolating a subtilisin peptide derived from peptide T8 with the sequence Val-Val-Thr. This was confirmed by determining the *N*-terminus of peptide T8 (alanine) and isolating a subtilisin peptide Val-Ala, which accounts for the two alanine residues of peptide T8 of protein Mz. The variant peptide of T4/T5 appears to involve a substitution of lysine for asparagine at position 171, giving only one peptide instead of the two observed in the other proteins studied. Asparagine rather than aspartic acid is indicated by the fact that the peptide was neutral at pH 6.5. Apart from these two variants the composition of all the tryptic peptides of this protein (Mz) corresponding to the *C*-terminal half was in full agreement with the sequence depicted in Table 1.

The results suggest that position 190 of human chains may be present in the population in two forms, one containing lysine and the other arginine. It is noteworthy that the Inv character of κ -chains is correlated with a leucine/valine substitution at position 191.

I acknowledge the helpful technical assistance of Mr J. M. Jarvis and thank Professor S. Cohen, Dr J. R. Hobbs and Dr A. Feinstein for providing the Bence-Jones proteins.

- Baglioni, C. & Cioli, D. (1966). *J. exp. Med.* **124**, 307.
 Baglioni, C., Zonta, L. A., Cioli, D. & Carbonara, A. (1966). *Science*, **152**, 1519.
 Hilschmann, N. & Craig, L. C. (1965). *Proc. nat. Acad. Sci., Wash.*, **53**, 1403.
 Milstein, C. (1966a). *Nature, Lond.*, **209**, 370.
 Milstein, C. (1966b). *Biochem. J.* **101**, 338.
 Milstein, C. (1966c). *Biochem. J.* **101**, 352.
 Milstein, C., Clegg, J. B. & Jarvis, J. M. (1967). *Nature, Lond.*, **214**, 270.
 Titani, K., Whitley, E., Avogardo, L. & Putnam, F. W. (1965). *Science*, **149**, 1090.
 Wikler, M., Titani, K., Shinoda, T. & Putnam, F. W. (1967). *J. biol. Chem.* **242**, 1668.

Biochem. J. (1967) **104**, 30c

The Amino Acid Sequence of the *N*-Terminal 84 Residues of a Human Heavy Chain of Immunoglobulin G (Daw)

By E. M. PRESS

Department of Immunology, St Mary's Hospital Medical School, London, W. 2

(Received 19 May 1967)

The chemical structure of the heavy chain of a human pathological immunoglobulin G has been studied by splitting with cyanogen bromide at the four methionine residues and separation of the resulting five fragments. The method of isolation, the composition and the order of the five fragments have been reported, together with the amino acid sequence of the *N*-terminal fragment (2b) consisting of 34 residues (Piggot & Press, 1967).

The next fragment from the *N*-terminal end is fragment 4, having 50 residues, and the amino acid sequence of this fragment has now been determined by using the methods described by Piggot & Press (1967). The α -amino group of this fragment is blocked.

Tryptic peptides. Fragment 4 was digested with trypsin (200 μ g./ μ mole) for 5 hr. at 37° at pH 8.3, and the digest fractionated on a column of Sephadex G-25 in aq. 20mN-NH₃. Five fractions were

obtained and from these five peptides were isolated; one fraction was pure (peptide 4T1) and the remaining fractions were purified by paper electrophoresis at pH 3.5. The compositions and mobilities of the peptides are given in Table 1.

Peptide 4T2 α contained homoserine and is therefore the *C*-terminal peptide. The partial sequence was established by the 'dansyl'-Edman technique to be: Asx*-Glx-Val-(Val,Leu,Ser)-homoserine.

Peptide 4T2 β was neutral at pH 6.5 and the sequence was established by the 'dansyl'-Edman technique to be: Asp-Thr-Ser-Lys.

Peptide 4T3 was shown by the 'dansyl'-Edman technique to have the sequence: Leu-Ala-Val-Ser-Lys.

Peptide 4T5 contained tryptophan and was

*Abbreviations: Asx, aspartic acid or asparagine; Glx, glutamic acid or glutamine; CM-Cys, *S*-carboxymethylcysteine; PCA, pyrrolid-2-one-5-carboxylic acid.