accuracy of the method decreases with decreasing concentration of riboflavin in the material examined. This low variation was confirmed by Bolton (private communication), who found with our procedure a coefficient of variation of 5.3% when analyzing ground-nut cake in 12 repeats (2.38 μ g. riboflavin/g., standard deviation ± 0.127 , or 5.3%).

RESULTS

Various foodstuffs were examined for their content of riboflavin. As may be seen in Table 2 the method is applicable to a wide variety of material. A number of foodstuffs have been assayed both chemically and microbiologically. The latter method is described in a following paper (Kodicek, 1949). Good agreement was found.

The method described above obviates the lengthy process of adsorption. The extraction of riboflavin when combined with enzymic digestion seems to be complete. The resulting extracts are clear and almost colourless.

It is important to point out that when dithionite is used as the reducing agent, blue light is

essential for obtaining the correct blank reading. We have observed that under ultraviolet light $\text{Na}_2\text{S}_2\text{O}_4$ reduces very strongly, or rather extinguishes almost completely, the total fluorescence of the extract. When a cell containing this reducing agent is placed in a beam of ultraviolet light in front of a cell containing riboflavin, the solution does not fluoresce. When blue light is used as the exciting source the fluorescence is almost unaltered.

SUMMARY

1. A modification of a fluorimetric method for the estimation of riboflavin has been described.

2. It estimates directly the fluorescence of riboflavin by using a sensitive photocell-galvanometer circuit without recourse to adsorption.

3. Washing of the extracts with chloroform and oxidation by potassium permanganate and hydrogen peroxide reduces the interference by unspecific fluorescent substances. The reduction of riboflavin by sodium dithionite has been used to obtain the value of the blank.

REFERENCES

- Andrews, J. S. (1943). *Cereal Chem.* **20**, 613.
Barton-Wright, E. C. & Booth, R. G. (1943). *Biochem. J.* **37**, 25.
Bolton, W. (1944). *J. agric. Sci.* **34**, 198.
Conner, R. T. & Straub, G. J. (1941). *Industr. Engng Chem.* (Anal. ed.), **13**, 380.
Hodson, A. Z. & Norris, L. G. (1939). *J. biol. Chem.* **131**, 621.
Kodicek, E. (1949). *J. gen. Microbiol.* (In the Press.)
Morell, D. B. (1947). *Aust. J. exp. Biol. med. Sci.* **25**, 251.
Morell, D. B. & Slater, E. C. (1946). *Biochem. J.* **40**, 644.
Rubin, S. H. & DeRitter, E. (1945). *J. biol. Chem.* **158**, 639.
Rubin, S. H., DeRitter, E., Schuman, R. L. & Bauernfeind, J. C. (1945). *Industr. Engng Chem.* (Anal. ed.), **17**, 136.
Scott, M. L., Hill, F. W., Norris, L. C. & Heuser, G. F. (1946). *J. biol. Chem.* **165**, 65.
Slater, E. C. & Morell, D. B. (1946). *Biochem. J.* **40**, 644.

The Fluorimetric Estimation of Nicotinamide in Biological Materials

BY D. K. CHAUDHURI AND E. KODICEK

Dunn Nutritional Laboratory, University of Cambridge and Medical Research Council

(Received 31 August 1948)

Scudi (1946) has reported that when a solution of nicotinamide is treated with cyanogen bromide (CNBr) and then extracted with *isobutanol* at an alkaline pH it fluoresces like the F_2 compound formed from N^1 -methylnicotinamide. Pyridoxal, vitamin B_1 , N^1 -methylnicotinamide and certain alkaloids also produce a similar fluorescence under these experimental conditions. Kodicek (1947) has modified the conditions by using methyl ethyl ketone in a single phase at an alkaline pH instead of

extracting with *isobutanol*. Even then vitamin B_1 , pteroylglutamic acid, N^1 -methylnicotinamide, pyridoxal and quinolinic acid interfere.

On re-examination of the reaction we have found that a fluorescent compound can be obtained from nicotinamide on treatment with cyanogen bromide, and on standing at an alkaline pH, without recourse to an organic solvent. By further modifications it was possible to eliminate the interference of other fluorescent substances. A method has thus been

worked out for the specific estimation of nicotinamide in various materials in the presence of other nicotinic acid derivatives.

EXPERIMENTAL

Fluorimeter. The photoelectric fluorimeter used in these experiments was of the 'one photocell type', with Wood's and calorex filters fitted between the incident light and the solution, and Wratten no. 47 and no. 2A filters placed between the solution and the photocell. A sensitive mirror galvanometer (3800 mm./1 μ amp.) was placed at a distance of 2 m. from the scale to obtain a high degree of sensitivity.

Reagents. NaOH (5N); CNBr, prepared fresh just before the experiment by adding ice-cold 10% aqueous NaCN drop by drop from a burette to ice-cold saturated bromine water until it is just decolorized; 0.2M-KH₂PO₄-NaOH buffer solution, pH 7.2; metaphosphoric acid (25%), freshly prepared; nicotinamide standard containing 2.5 mg./100 ml. water, prepared weekly from a stronger nicotinamide solution.

are kept in the dark at room temperature for 45 min., after which the readings of fluorescence can be taken.

The blank correction. For the blank, the solution undergoes the same treatment except that the CNBr reagent is not added ('No CNBr' blank). The galvanometer deflexion due to the blank was usually 1 mm. when working with mixtures of pure substances.

Variation of factors

The conditions of the reaction described above have been studied by varying the individual factors which may affect the formation of the fluorescent compound.

(1) Effect of varying concentrations of CNBr and NaOH and of time in alkali on the development of fluorescence

Fig. 1 shows the effect of 2 and 4 ml. CNBr and also of 4, 6 and 8 ml. 5N-NaOH. It will be seen that 4 ml. CNBr (added in the first stage) and 8 ml. 5N-NaOH (added in the second stage) give the best readings. The maximum fluorescence was attained in 30 min. and was constant for at

Fig. 1. Effects of varying concentrations of CNBr and of NaOH, and of time in alkali on the development of fluorescence of nicotinamide solutions. A = 4 ml. 5N-NaOH; B = 6 ml. 5N-NaOH; C = 8 ml. 5N-NaOH.

Principle of method for solutions of pure substances

First stage (CNBr treatment). The nicotinamide standard (2 ml. containing 50 μ g. nicotinamide) and 2 ml. phosphate buffer, are mixed with the CNBr solution and distilled water to about 14 ml. in 15 ml. glass-stoppered measuring flasks. The flasks are immersed for 4 min. in a water bath at 56–58°, then cooled in ice-cold water for 5 min. and the volume made up to 15 ml. with water.

Second stage (development of fluorescence). From 5 to 15 ml. of the CNBr-treated solution are taken, 8 ml. 5N-NaOH added, and the volume made up to 30 ml. with water in a measuring cylinder. The contents are mixed thoroughly and transferred to a conical flask. The solutions

least another 90 min. A period in the dark of 45 min. has therefore been adopted.

Higher concentrations of CNBr (6 ml.) caused a decrease of fluorescence. Slightly larger amounts of NaOH (10 and 12 ml.) did not change the maximal fluorescence, but 16 ml. caused a diminution. Therefore 4 ml. CNBr and 8 ml. 5N-NaOH have been adopted.

The concentration of the phosphate buffer was found to be of no consequence so long as its final concentration was kept above 0.01M. We used 1 ml. 0.2M-phosphate buffer in 15 ml. of the reaction mixture.

(2) Fluorescence with increasing amounts of nicotinamide

Fig. 2 shows the fluorescence obtained with increasing concentrations of nicotinamide. The fluorescence is proportional

to the concentration of nicotinamide up to a final value of about 0.6 $\mu\text{g./ml.}$

Fig. 2. Fluorescence obtained with increasing amounts of nicotinamide

(3) Elimination of interference by other fluorescent compounds

The following compounds were tested and found not to fluoresce: *N*¹-methylnicotinamide, nicotinic acid, coramine, quinolinic acid, α -picolinic acid, trigonelline, nipecotic acid (piperidine-3-carboxylic acid), urea, uric acid, yeast nucleic acid, choline chloride, *p*-aminobenzoic acid, adenosinetriphosphate, inositol, biotin, calcium pantothenate, riboflavin and indole. On the other hand, vitamin B₁ and diphosphopyridine nucleotide gave a fluorescence under the conditions of our test. Pyridoxal, pteroylglutamic acid and nicotinic thioamide (Karrer & Schukri, 1945) gave a slight fluorescence.

The fluorescence due to vitamin B₁ and coenzyme I was particularly strong. To eliminate this interference, two possible ways were explored: (1) the use of another blank, i.e. the 'No NaOH' blank, and (2) pre-treatment of the vitamin solution, prior to the CNBr treatment, by acidification to remove acid-insoluble compounds, like folic acid, followed by weakly alkaline digestion which destroys the vitamin B₁ fluorescence but does not affect that due to nicotinamide.

The 'No NaOH' blank, which was obtained by omitting the addition of 8 ml. 5*N*-NaOH in the second stage, proved to be unsuitable. On the other hand, the pre-treatment eliminated the interference by other substances, and did not diminish the fluorescence of nicotinamide, present as such or formed from other pyridine compounds. The results are seen in Table 1 which gives galvanometer readings due to the fluorescence of various compounds with and without pre-treatment.

Pre-treatment of the solution. The solution to be tested is adjusted to pH 2 with conc. HCl and allowed to stand for 5 min. at room temperature. If any precipitate is formed, it is centrifuged off. The solution, in a conical flask, is then brought to pH 9.4-9.6 with 20% NaOH (thymol blue as external indicator) and immersed in a boiling water bath for 30 min., after which it is cooled, adjusted to pH 7.2, and made up to a known volume. The steps described under 'Principle of method' are then performed.

From Table 1 it can be seen that this 'pre-treatment' does not affect the intensity of the fluorescence of nicotinamide. Diphosphopyridine nucleotide, when tested in equimolar concentration, gave a fluorescence of the same magnitude as nicotinamide, indicating that the molecule had been split between the pyridine ring and ribose, and free nicotinamide formed. Nicotinic thioamide was converted to a compound giving a fluorescence, but of a lower intensity than that of nicotinamide. As there is no evidence that this compound occurs in nature, this finding is only of theoretical interest. All the other compounds tested by the final procedure showed either no, or little, fluorescence, which in every case was accounted for by the 'No CNBr' blank. Similar

Table 1. Specificity of fluorescence

Substance	Final conc. ($\mu\text{g./ml.}$)	Galvanometer reading (mm.)					
		Without pre-treatment			With pre-treatment: final procedure		
		'No CNBr' blank	'No NaOH' blank	Full fluorescence	'No CNBr' blank	'No NaOH' blank	Full fluorescence
Nicotinamide	0.55	1	1	71	1	1	71
Coenzyme I (48% purity)	6.2*	46	1	70	2.5	1	72
Nicotinic thioamide	1.1	1	1.5	2	1	1.5	45
Aneurin	0.55	5	103	101	1	2	1
Nicotinic acid	1.1	1	1	1	1	1	1
Coramine	1.1	0.5	0.5	0.5	0.5	0.5	0.5
<i>N</i> ¹ -Methylnicotinamide chloride	1.1	1	1	1	1	1	1
Quinolinic acid	1.1	1	1	1	1	1	1
α -Picolinic acid	1.1	—	—	—	1.5	3	1.5
Trigonelline	1.1	—	—	—	1	1	1
Urea	1.1	—	—	—	1	1	1
Pteroylglutamic acid	1.1	1	21	1	1	16	1
Pyridoxal	1.1	1	46	1.5	2	42	2
Riboflavin	0.78	1	11	1	4	16	4

* Equimolar with nicotinamide.

results were also obtained when mixtures of the above substances were tested. The only exception was nicotinic acid, the CNBr-addition compound of which decreased the fluorescence of CNBr-treated nicotinamide. The nature of this finding was examined in the following experiments.

(4) *Effect of different concentrations of nicotinic acid on the fluorescence of nicotinamide*

From Fig. 3 it can be seen that there is a proportional decrease of fluorescence with increasing concentration of nicotinic acid. The depression is non-competitive; it does not depend on the concentration of nicotinamide, since

Fig. 3. Effect of different concentrations of nicotinic acid on the fluorescence of nicotinamide. A = nicotinamide; B = nicotinamide + 25 μ g. nicotinic acid; C = nicotinamide + 50 μ g. nicotinic acid; D = nicotinamide + 100 μ g. nicotinic acid.

when this is varied the same percentage inhibition of fluorescence is obtained, depending on the concentration of nicotinic acid (Table 2). Fortunately, even with large amounts of nicotinic acid, the fluorescence of nicotinamide is proportional to concentration within limits. We shall describe below how the fluorescence of nicotinamide can be quantitatively measured even in the presence of nicotinic acid by using an internal standard of nicotinamide.

Nature of 'quenching' effect of nicotinic acid. The depressing effect of the nicotinic acid CNBr-addition compound on the fluorescence of nicotinamide seems to be an 'inner filter' effect. It occurs only with nicotinic acid; no other pyridine derivative gives a similar result. The 'inner filter' effect was obtained only when nicotinic acid was treated with CNBr, irrespective of whether the nicotinic acid was added, as such, at the start of the procedure or added as the CNBr complex immediately before the reading of the fluorescence. Similar results were obtained when a nicotinic acid-CNBr solution was placed between the fluorescent nicotinamide solution and the incident ultraviolet light.

Table 2. *Percentage inhibition of fluorescence of nicotinamide by nicotinic acid*

Nicotinic acid (μ g.)	Nicotinamide (μ g.)	Galvanometer reading (mm.)	Inhibition (%)
0	12.5	16	0
0	25	32	0
0	50	64	0
Mean			0
25	12.5	14	12.5
25	25	28.5	11.0
25	50	56.5	11.7
Mean			11.7
50	12.5	12	25
50	25	24	25
50	50	48.5	24.2
Mean			24.7
100	12.5	8	50
100	25	16	50
100	50	32.5	49.3
Mean			49.8

Final procedure for the estimation of nicotinamide

Extraction and pre-treatment

The material to be tested (5 g.) is weighed out, cut up finely with scissors if necessary, and ground with sand together with a small amount of 0.1 N-HCl (1-2 ml.). The ground tissue is transferred with 40 ml. water into a 100 ml. beaker, and heated in a boiling water bath for 30 min. The solution is cooled, brought to pH 2 with concentrated HCl, centrifuged, and the residue washed with 10 ml. 0.1 N-HCl and centrifuged. The combined centrifugates are brought to a known volume with 0.1 N-HCl (usually 40 ml.). Freshly prepared 25% metaphosphoric acid (6 ml.) is added. The precipitate is centrifuged after standing for 5-10 min. at room temperature. The clear solution is brought to pH 9.4-9.6 (thymol blue as external indicator), and is then heated in a beaker in a boiling water bath for 30 min. The solution is cooled, carefully adjusted to pH 7.2, and the volume made up to 50 ml. It is then filtered through Whatman no. 5 filter paper, and is ready for the next stage of the estimation. An amount of the extract (referred to as U) is taken to contain about 5-25 μ g. of nicotinamide.

Development of fluorescence

First stage (CNBr treatment). Three 15 ml. glass-stoppered measuring flasks are taken (preferably in duplicate), and the reagent mixed according to the following scheme:

Flask A (blank). x ml. U + 2 ml. 0.2 M-phosphate buffer + (12 - x) ml. water.

Flask B (unknown). x ml. U + 2 ml. 0.2 M-phosphate buffer + 4 ml. CNBr solution + (8 - x) ml. water.

Flask C (unknown + internal standard). 1 ml. (25 μ g.) nicotinamide standard + x ml. $U + 2$ ml. 0.2M-phosphate buffer + 4ml.CNBr solution + $(7-x)$ ml. water.

The flasks, after their contents have been thoroughly mixed, are immediately placed for 4 min. in a water bath at 56–58°. It is convenient to suspend them on the rim of the water bath by a string counterbalanced by the stopper. The solutions are then cooled in ice-cold water for 5 min. and the volume made up to 15 ml. with water.

Second stage. The CNBr-treated solution (5 ml.) is placed in a measuring cylinder (up to 15 ml. can be taken, depending on the concentration of nicotinamide and the clearness of the solution). NaOH (8 ml., 5N-) is added, and the volume made up to 30 ml. with water. The contents are mixed and transferred to conical flasks which are kept for 45 min. at room temperature in the dark. The readings of fluorescence are then taken.

Calculation

Let a = galvanometer reading corresponding to flask A; b = galvanometer reading corresponding to flask B; c = galvanometer reading corresponding to flask C; x = ml. of U taken for analysis; and f = dilution factor. Then

$$\text{nicotinamide } (\mu\text{g./g.}) = \frac{b-a}{c-b} \cdot \frac{25}{x} \cdot f.$$

Reproducibility of results

Replicate estimations of the nicotinamide content of a specimen of rat muscle using the method described here have been statistically analyzed and compared with nicotinic acid results obtained with a modified *p*-aminoacetophenone procedure (Wang & Kodicek, 1943; Harris & Raymond, 1939). It will be seen that the standard error is small. The results are shown in Table 3.

Table 3. *Reproducibility of results*

Rat muscle (sample no.)	Nicotinamide (μ g./g. tissue)	Total nicotinic acid (μ g./g. tissue)
1	62.5	71.2
2	61.0	69.6
3	65.8	68.3
4	64.3	68.9
5	69.5	70.8
6	62.5	72.5
Mean	64.3 (± 3.05 , standard error of an indi- vidual estima- tion)	70.2 (± 1.66 , standard error of an indi- vidual estima- tion)
Standard error of the mean	1.24	0.68

RESULTS AND DISCUSSION

The method as described has proved a convenient and rapid procedure for the estimation of nicotinamide. It seems to be reasonably specific, as no naturally occurring substances capable of fluorescence which we have tried so far have been found to interfere. The quenching which is experienced in extracts is corrected for by the internal standard.

The method estimates total nicotinamide either present as such or derived from nicotinamide-containing compounds such as coenzyme I. These break down during the pre-treatment and nicotinamide is liberated. Precautions must be taken which have been already described by Wang & Kodicek (1943) in the CNBr reaction: the pH must be adjusted carefully to 7.2, and only fresh cyanogen bromide, prepared from ice-cold solution, should be used.

Table 4. *Nicotinamide and nicotinic acid content of biological materials*

Material analyzed	Present authors		Krehl <i>et al.</i> (1946)	
	Nicotin- amide (μ g./g.)	Total nicotinic acid (μ g./g.)	Nicotin- amide (μ g./g.)	Total nicotinic acid (μ g./g.)
Rat:				
Liver	125	119	148	154
Muscle	55	52	73	78
Kidney	83	78	79	87
Heart	91	94	—	—
Spleen	40	42.7	—	—
Lung	19	17	—	—
Brain	35	41.8	—	—
Testis	18	21.6	—	—
Wheat bran	0	165	48	245
Wheat flour (100% ex- traction)	<5	42	15	49.7
Wheat germ	20.8	39.2	20.7	43.5
Dried yeast	185.5	380	—	—
Baker's yeast	100.4	199	—	—
Brewer's yeast	62.8	124.8	—	—
Marmite	0	640	—	—

Table 4 shows the results obtained by our method on various materials. The values were compared with the total nicotinic acid content estimated chemically by a modified *p*-aminoacetophenone procedure. The results of Krehl, Huerga, Elvehjem & Hart (1946) on similar materials have been listed in this table for comparison so far as they have been reported. The method used by Krehl *et al.* is based on a microbiological procedure using *Leuconostoc mesenteroides*, which estimates only free nicotinic acid. The nicotinamide and compounds other than nicotinic acid are therefore calculated by difference between values for the hydrolyzed

and unhydrolyzed sample. Our figures agree fairly well with those of the American workers.

The rat tissues which we have studied seem to contain almost all their nicotinic acid in the form of amide, bound or free. Yeast contains about 50 % of the vitamin as amide compounds. From our results we do not know whether the free nicotinic acid found in yeast is an artifact, caused by rapid breakdown of the amide to nicotinic acid, though the same proportion of the acid and amide has been observed both in dried and fresh yeasts. Autolyzed yeast and marmite evidently contain only the nicotinic acid and no amide.

Cereal products give interesting results. Krehl *et al.* (1946) reported that a certain proportion of the high amount of nicotinic acid compounds in bran is possibly present as the amide, though they qualify their statement that possibly the precursor of nicotinic acid (Kodicek, 1940; Krehl & Strong, 1944; Krehl, Elvehjem & Strong, 1944) may contribute to the apparent nicotinamide value. We did not find any nicotinamide in bran and conclude that the 'precursor' of nicotinic acid present in this cereal must break down on our pre-treatment to free nicotinic acid and not to the amide. On the other hand, wheat germ contained about 50 % of its vitamin content as nicotinamide. The negative results for 100 % extraction whole-wheat flour seem to indicate that only small amounts of nicotinamide are present which would not be detected because of the large experimental error occurring in low-vitamin extracts.

SUMMARY

1. A quantitative method for the estimation of nicotinamide has been developed, based on the formation of a fluorescent compound from nicotinamide on treatment with cyanogen bromide under specified conditions.

2. The method eliminates possible interference by other compounds giving a similar fluorescence under the conditions of the test by the use of acid and alkaline pre-treatment and of a special blank. It estimates the total nicotinamide content, including the free and bound forms.

3. The procedure gives specific and reproducible results for biological materials and for cereals, and the results agree well with the reported microbiological values.

4. Practically all the vitamin in rat organs and muscles seems to be present in the form of the amide, bound or free. In bran, no nicotinamide could be detected before or after digestion. It is assumed that the breakdown product of the 'precursor' of nicotinic acid present in bran is the free acid and not the amide. Yeast and wheat germ seem to contain about 50 % of the vitamin present in the amide form, bound or free.

Our thanks are due to Dr L. J. Harris for his criticism and helpful suggestions, to Prof. P. Karrer for kindly supplying us with nicotinic thioamide, to Dr T. H. Jukes of Lederle Laboratories for supplies of pteroylglutamic acid, and to Dr K. Folkers of Merck and Co., Rahway, U.S.A., for the pyridoxal hydrochloride. One of us (D. K. Chaudhuri) is an Indian Government Scholar.

REFERENCES

- Harris, L. J. & Raymond, W. D. (1939). *Biochem. J.* **33**, 2037.
Karrer, P. & Schukri, J. (1945). *Helv. chim. Acta*, **28**, 820.
Kodicek, E. (1940). *Biochem. J.* **34**, 724.
Kodicek, E. (1947). *Analyst*, **72**, 385.
Krehl, W. A., Elvehjem, C. A. & Strong, F. M. (1944). *J. biol. Chem.* **156**, 13.
Krehl, W. A., Huerga, J., Elvehjem, C. A. & Hart, E. B. (1946). *J. biol. Chem.* **166**, 53.
Krehl, W. A. & Strong, F. M. (1944). *J. biol. Chem.* **156**, 11.
Scudi, J. V. (1946). *Science*, **103**, 567.
Wang, Y. L. & Kodicek, E. (1943). *Biochem. J.* **37**, 530.