Solutions of glutamine were incubated in buffer solutions at 37.4°. Samples were withdrawn at intervals, and the extent of decomposition measured by estimation of ammonia; there was no increase in formol acidity. The results are given in Table 2.

It is clear that on long incubation there is appreciable decomposition of glutamine. This is greater in phosphate buffer, in which approximately 28 % of glutamine is decomposed in 20 hr., than in a phosphate-free buffer (veronal), in which only 6 % is decomposed. The absence of increase in formol acidity suggests that pyrrolidonecarboxylic acid is the product. This we have confirmed by isolation of the compound from a solution of glutamine (400mg.) incubated in phosphate buffer (100 ml.) for 42 hr. The acid was extracted with ethyl acetate (cf. Woodward & Reinhart, 1942) and had m.p. 157–160°: melting point mixed with authentic pyrrolidonecarboxylic acid, 157–160°.

Attempted isolation of end product of the action of liver extracts on glutamine. For these experiments short-term digests were used so as to avoid the complication of the formation of considerable amounts of pyrrolidonecarboxylic acid by spontaneous decomposition of glutamine.

We were unable to detect any pyrrolidonecarboxylic acid in our digests by the method used successfully

by Woodward & Reinhart (1942) for its isolation from digests of glutathione. Several estimations of the loss of amino nitrogen supported Krebs's (1935) findings that glutamic acid and not pyrrolidonecarboxylic acid is formed. Attempts to isolate glutamic acid from glutamine digests have so far been unsuccessful, but paper chromatography (cf. Consden, Gordon & Martin, 1944) of the material extracted by *n*-butanol from such digests suggests that the product is indeed glutamic acid. We thus have no reason to dispute the conclusion of the earlier workers that glutamic acid is the principal product of the enzymic hydrolysis of glutamine.

SUMMARY

The nature of the products of the action of tissue extracts on glutamine has been examined. Whilst glutamic acid appears to be the main product, significant amounts of pyrrolidonecarboxylic acid may be present in long-term digests, owing to spontaneous decomposition of glutamine.

The authors' thanks are due to the Royal Society for a Government Grant which defrayed part of the cost of this investigation. We are indebted to Mr P. B. Wood for carrying out some of the amino-N estimations. We wish to thank Prof. H. A. Krebs, F.R.S., for providing a specimen of pure glutamine for comparison with our preparation.

REFERENCES

- Archibald, R. M. (1944). *J. biol. Chem.* **154**, 643.
Archibald, R. M. & Hamilton, P. B. (1945). Cited in R. M. Archibald, *Chem. Rev.* **37**, 161.
Bray, H. G., James, S. P., Raffan, I. M., Ryman, B. E. & Thorpe, W. V. (1949). *Biochem. J.* **44**, 618.
Chibnall, A. C. & Westall, R. G. (1932). *Biochem. J.* **26**, 122.
Consden, R., Gordon, A. H. & Martin, A. J. P. (1944). *Biochem. J.* **38**, 224.
Hamilton, P. B. (1945). *J. biol. Chem.* **158**, 375.
Harris, L. J. (1923). *Proc. Roy. Soc. B*, **95**, 440, 500.
Krebs, H. A. (1935). *Biochem. J.* **29**, 1951.
Léuthardt, F. M. (1940). *Hoppe-Seyl. Z.* **265**, 1.
Melville, J. (1935). *Biochem. J.* **29**, 184.
Vickery, H. B., Pucher, G. W., Clarke, H. E., Chibnall, A. C. & Westall, R. G. (1935). *Biochem. J.* **29**, 2710.
Woodward, G. E. & Reinhart, F. E. (1942). *J. biol. Chem.* **145**, 471.

The Effect of Various Diets on the Metabolism of Nicotinic Acid and Nicotinamide in the Rabbit

By P. ELLINGER AND M. M. ABDEL KADER, *Lister Institute of Preventive Medicine, London*

(Received 7 January 1949)

It has recently been shown (Ellinger & Abdel Kader, 1948, 1949) that nicotinic acid and nicotinamide are not metabolized in an identical manner by various mammalian species. Man, dog, cat and rat aminate nicotinic acid to nicotinamide and methylate the latter to nicotinamide methochloride, whilst the rabbit and guinea pig deaminate nicotinamide to nicotinic acid and eliminate small amounts of an

unidentified compound which was provisionally called 'pseudo nicotinamide methochloride'. Whilst the guinea pig eliminates nicotinic acid unchanged, the rabbit methylates part of it to trigonelline. It appears that the species which aminate nicotinic acid are omnivorous or carnivorous, while those which deaminate nicotinamide are strictly herbivorous. It appeared possible that these differences

are related to the type of diet and not to the constitution of the species. In order to decide which was the real cause, the work described in this paper was carried out. Rabbits were chosen as test objects for the following reasons. It can be expected that a change in diet will affect the omnivorous less than the strictly herbivorous animals, and, among the latter, the rabbit has shown a more complex metabolism than the guinea pig. In addition, it is known that rabbits can be trained to eat meat (Aberhalden & Brahm, 1909).

regained 330 and 390 g., respectively, during the meat period and at the end their weights were 2730 and 2753 g. respectively. At the end of the second cabbage period their weights were 2350 and 2420 g. respectively. Without extradietary intake of either acid or amide the only metabolites detected were nicotinic acid and 'pseudo nicotinamide methochloride' (Table 2), as was found in the earlier experiments (Ellinger & Abdel Kader, 1949). The amounts of nicotinic acid eliminated were lowest during the cabbage, somewhat higher during the oat

Table 1. *Protein, fat, carbohydrate, nicotinic acid and nicotinamide contents and caloric values of the diets used*

Diet		Protein		Fat		Carbohydrate	
Type	(g./day)	(g./100 g.)	(g./day)	(g./100 g.)	(g./day)	(g./100 g.)	(g./day)
Cabbage	200	1.6*	3.2	0.3*	0.6	5.6*	11.2
Rolled oats	100	16.7*	16.7	7.3*	7.3	66.2*	66.2
Meat meal + national bread	50	64.2§	32.1	26.4§	13.2	—	—
	50	8.5	4.25	1.2	0.6	57.4	28.7
			36.35		13.8		

Diet		Calories		Nicotinic acid (mg./day)	Nicotinamide (mg./day)
Type	(g./day)	(per 100 g.)	(per day)		
Cabbage	200	31.6†	63.2	0.204‡	0.392‡
Rolled oats	100	399.5†	399.5	0.530‡	0.720‡
Meat meal + national bread	50	503.2†	251.6	1.250‡	1.492‡
	50	250.0	125.0		
			376.6		

* From Tibbles (1912).

† Calculated (protein, 4.0 cal.; fat, 9.3 cal.; carbohydrate, 4.0 cal.).

‡ Determined.

§ Analysis supplied by Mr F. C. Hymas, Spratt Patent Ltd.

|| From Medical Research Council (1945).

EXPERIMENTAL

Two male rabbits were kept in metabolism cages at 22–27°. The urine was collected every 24 hr. and analyzed for nicotinic acid, nicotinamide, trigonelline and 'pseudo nicotinamide methochloride' as described (Ellinger & Abdel Kader, 1949). The rabbits were fed on cabbage alone for 3 weeks, on rolled oats for 4 weeks, on national bread and meat meal for 29 days and again on cabbage for another 22 days. Oats, bread and meat meal were moistened with water.

When the daily urinary output of the various metabolites was approximately constant for at least 5 days, nicotinic acid or nicotinamide (100 or 200 mg.) was injected intraperitoneally at intervals of 3–5 days. The increase in metabolite elimination following the extradietary administration of the compounds was calculated by deducting the average daily predosing output from that found after administration of the compound. In all cases the predosing level was reached again after 24 hr. In two experiments choline chloride (0.15 g.) was injected intraperitoneally together with 0.2 g. of nicotinic acid.

RESULTS

Both rabbits reacted similarly to changes in the diet. On the cabbage diet the rabbits kept their weight of 2810 and 2850 g. respectively. They lost 540 and 560 g. respectively, during the oat period; they

and highest during the meat period, in accordance with the daily dietary intake of nicotinic acid and

Table 2. *Daily urinary elimination of nicotinamide metabolites by rabbits kept on different diets*

(Values for 'pseudo nicotinamide methochloride' are expressed in terms of nicotinamide methochloride.) (Neither nicotinamide, nicotinuric acid nor trigonelline were eliminated.)

Diet	Rabbit no.	Days	Metabolites eliminated (mg./day)	
			Nicotinic acid	'Pseudo nicotinamide methochloride'
Cabbage	1	5	0.793	0.501
	2	5	0.577	0.537
Oats	1	5	0.992	0.565
	2	5	0.983	0.500
Meat-bread	1	8	1.755	0.619
	2	8	1.770	0.645

nicotinamide during each of these periods. The amount of 'pseudo nicotinamide methochloride' eliminated did not change markedly with the diet. Since the nature of this compound is unknown it could not be decided whether the output of meta-

bolites exceeded the intake of the compounds, indicating biosynthesis.

As in the earlier experiments (Ellinger & Abdel Kader, 1949) on cabbage or oat diet extradietary nicotinic acid was nearly completely eliminated, but nicotinamide was eliminated to a much lesser extent, particularly after doses of 100 mg. The meat diet did not markedly affect the total metabolite output with the exception that after 100 mg. amide, output was considerably higher than on either of the other diets (Table 3).

Both the ability to methylate and deaminate were, therefore, affected during the meat period. To obtain information as to whether this was due directly to the kind of food taken, or to changes caused by the diet on the tissues responsible for the metabolism, the metabolism of both nicotinic acid and nicotinamide was studied at different times after changing from the meat to the cabbage diet (Table 5). On the eighth day after the change, total output was still low, methylation was still absent and deamination as low as during the meat period. Methylation to

Table 3. *Effect of diet on increase in the total urinary elimination of nicotinamide metabolites (except 'pseudo nicotinamide methochloride') following the extradietary administration of nicotinic acid or nicotinamide*

(Figures are given as percentage of administered dose.)

		Total elimination							
		Rabbit 1				Rabbit 2			
Compound administered ...		Nicotinic acid		Nicotinamide		Nicotinic acid		Nicotinamide	
Dose (mg.) ...		100	200	100	200	100	200	100	200
Diet:									
Cabbage		99.1	93.0	23.5	78.4	81.7	95.1	28.1	70.5
Oat		95.0	96.8	18.0	70.0	84.9	97.1	22.8	63.6
Meat-bread		92.1	96.3	36.5	55.3	76.0	99.3	43.2	61.9

Table 4. *Effect of diet on increase in the urinary elimination of individual nicotinamide metabolites (except 'pseudo nicotinamide methochloride') following the extradietary administration of nicotinic acid or nicotinamide*

(Figures are given as percentage of administered dose.)

		Metabolite eliminated (%)							
		Rabbit 1				Rabbit 2			
Compound administered ...		Nicotinic acid		Nicotinamide		Nicotinic acid		Nicotinamide	
Dose (mg.) ...		100	200	100	200	100	200	100	200
Metabolite eliminated	Diet								
Nicotinic acid	Cabbage	50.4	64.0	11.0	18.0	59.9	69.3	20.3	33.0
	Oats	45.3	62.4	9.5	17.2	63.7	74.3	17.9	30.3
	Meat-bread	92.1	96.3	14.1	7.8	76.0	99.3	16.0	12.7
Nicotinamide	Cabbage	0	0	2.6	17.7	0	0	0	0
	Oats	0	0	0.6	12.7	0	0	0	0
	Meat-bread	0	0	22.4	47.6	0	0	27.2	49.2
Trigonelline	Cabbage	48.5	29.0	9.9	42.7	25.8	25.7	7.8	36.6
	Oats	49.6	34.4	7.9	40.2	21.1	22.7	5.8	33.3
	Meat-bread	0	0	0	0	0	0	0	0

On a cabbage or oat diet, nicotinic acid injection was followed by the output of nicotinic acid and trigonelline only; this was also the case in one rabbit after nicotinamide injection, whilst the other eliminated in addition small amounts of unchanged nicotinamide (Table 4). During the meat period the results were very different, nicotinic acid was entirely eliminated as such and nicotinamide mostly unchanged, but to a small degree deaminated to the acid. Methylation was completely absent. There was no methylation either when choline was injected together with the acid.

a reduced extent began again on the eleventh day, but methylation and deamination had not yet reached the pre-meat level on the sixteenth day of the cabbage diet. On the twenty-first day the pre-meat level was almost attained in both respects.

The output of 'pseudo nicotinamide methochloride' after extradietary nicotinic acid administration was hardly affected by the meat diet; that after nicotinamide application was, however, 2-3 times higher during the meat than during the other periods.

Table 5. *Changes in the urinary elimination of products of nicotinic acid and nicotinamide metabolism of rabbits on replacing the meat-bread by a cabbage diet following extradietary administration of the compounds*

(Nicotinic acid or nicotinamide (200 mg.) administered. Figures for elimination are given as percentage of administered dose. Column 2 denotes days after end of meat-bread period.)

Compound administered ...		Metabolites eliminated			
		Rabbit 1		Rabbit 2	
		Nicotinic acid	Nicotinamide	Nicotinic acid	Nicotinamide
Total output	8th	—	58.6	—	54.6
	11th	87.7	—	87.3	—
	16th	—	56.2	—	63.3
	21st	—	73.8	—	68.1
Nicotinic acid eliminated	8th	—	10.1	—	5.2
	11th	68.5	—	73.4	—
	16th	—	18.2	—	29.0
	21st	—	20.8	—	35.9
Nicotinamide eliminated	8th	—	48.4	—	49.3
	11th	0	—	0	—
	16th	—	13.6	—	7.6
	21st	—	17.1	—	0
Trigonelline eliminated	8th	—	0	—	0
	11th	19.3	—	13.9	—
	16th	—	24.4	—	26.7
	21st	—	35.9	—	32.2

Table 6. *Effect of diet on urinary elimination of 'pseudo nicotinamide methochloride' following extradietary administration of nicotinic acid or nicotinamide to the rabbit*

(The values of 'pseudo nicotinamide methochloride' are expressed in terms of nicotinamide methochloride.)

Compound administered ...		'Pseudo nicotinamide methochloride' elimination (mg.)							
		Rabbit 1				Rabbit 2			
		Nicotinic acid		Nicotinamide		Nicotinic acid		Nicotinamide	
Dose (mg.)	...	100	200	100	200	100	200	100	200
Diet:									
Cabbage		0.500	0.384	0.600	0.800	0.460	0.484	0.750	0.825
Oats		0.480	0.456	0.651	0.680	0.432	0.405	0.600	0.660
Meat-bread		0.575	0.488	1.920	2.011	0.600	0.501	1.230	1.698

DISCUSSION

Two different effects of the three diets, cabbage, oats and meat-bread, were studied, i.e. that on body weight and that on nicotinic acid and nicotinamide metabolism. The caloric intake of the rabbits during the two cabbage periods was very low; but in spite of this the body weight remained constant on 63 cal./day during the first of these periods. The basal metabolic rate of the rabbits, calculated according to Evans (1945) using Meeh's (1879) formula for the surface area, was 87 cal. The apparent adequacy of this low food intake might have been due to the high temperature of the room and the static conditions of the rabbits in the limited space of the metabolism cages or to a reduced metabolic rate caused by a possible goitrogenic effect of cabbage in rabbits (Chesney, Clawson & Webster, 1928;

Webster, Clawson & Chesney, 1928; Webster & Chesney, 1928; Marine, Baumann & Cipra, 1929; Webster & Chesney, 1930; Marine, Baumann & Webster, 1930; Webster, Marine & Cipra, 1931). The far higher caloric intake of oats did not prevent a loss of body weight. Oats proved, therefore, to be inadequate for the rabbit, whilst the meat-bread mixture was much better utilized. The inadequacy of the 63 cal. during the second cabbage period might be due either to an increased metabolic rate caused by the preceding oat or meat-bread diet or to a depletion of the rabbit during the oat and meat periods of an essential factor which could not be supplied in sufficient amounts by the small cabbage ration.

The results of the three diets on the nicotinic acid and nicotinamide metabolism were different. Here feeding of oats after cabbage did not cause a marked

change, but feeding of the meat-bread mixture altered the metabolism essentially. No new function like amidation of nicotinic acid, characteristic of omnivorous species, was developed, but the ability to methylate was lost completely and that to deaminate was lost to a large extent when meat was fed. Both injected nicotinic acid and amide were found to be eliminated mainly unchanged in the urine. The persistence of the change in metabolism long after the return to the cabbage diet shows that this alteration is not caused directly by the actual food administered, but is due to changes, caused by the diet, in the tissues responsible for methylation and deamination. In the rat, the methylation takes place in the liver (Perlzweig, Bernheim & Bernheim, 1943; Ellinger, 1946, 1948), while the site of deamination is unknown, but might be the kidney and brain which are able to carry out the amination of nicotinic acid (Ellinger, 1946, 1948). From the fact that methylation is more quickly restored than deamination, it might be concluded that the two functions are independent of each other. The most likely explanation of the effect of the meat diet on nicotinic acid and nicotinamide metabolism is a damage of enzymic systems involved in methylation and deamination caused either by the inadequacy of the animal protein or by the lack of an essential factor in the diet.

The considerable increase in the output of 'pseudo

nicotinamide methochloride' after nicotinamide intake cannot be explained as long as the nature of this compound remains unknown. But this increased elimination may make it possible to collect sufficient amounts of this compound to study its nature.

SUMMARY

1. A change from a cabbage to an oat diet had relatively little effect on the metabolism of nicotinic acid and nicotinamide in rabbits. However, a change to a meat-bread diet suppressed the methylating mechanism completely, and the deaminating mechanism to a large extent; but no amidation of nicotinic acid as in omnivorous animals was developed. Replacing the meat-bread by a cabbage diet restored the original metabolic condition within 3 weeks. It is concluded that this altered metabolism is not due directly to the food fed, but to changes caused by the altered food intake on the tissues responsible for the metabolism.

2. The body weight was affected by the three diets in different ways. 63 cal./day were sufficient to maintain weight on the cabbage diet, while 400 cal. were insufficient on the oat diet, whilst with the meat-bread diet weight was gained with 380 cal.

We wish to thank Messrs Spratts Patent, Ltd., for a generous gift of meat meal.

REFERENCES

- Abderhalden, E. & Brahm, C. (1909). *Hoppe-Seyl. Z.* **62**, 133.
- Chesney, A. M., Clawson, T. A. & Webster, B. (1928). *Johns Hopk. Hosp. Bull.* **43**, 261.
- Ellinger, P. (1946). *Biochem. J.* **40**, xxxi.
- Ellinger, P. (1948). *Biochem. J.* **42**, 175.
- Ellinger, P. & Abdel Kader, M. M. (1948). *Biochem. J.* **42**, xxxiii.
- Ellinger, P. & Abdel Kader, M. M. (1949). *Biochem. J.* **44**, 78.
- Evans, L. C. (1945). *Principles of Human Physiology*, 9th ed. London: Churchill, p. 762.
- Marine, D., Baumann, E. J. & Cipra, A. (1929). *Proc. Soc. exp. Biol., N.Y.*, **26**, 822.
- Marine, D., Baumann, E. J. & Webster, B. (1930). *Proc. Soc. exp. Biol., N.Y.*, **27**, 1029.
- Medical Research Council (1945). *War Memorandum*, no. 14. London: H.M.S.O.
- Meeh, K. (1879). *Z. Biol.* **15**, 428.
- Perlzweig, W. A., Bernheim, M. L. C. & Bernheim, F. (1943). *J. biol. Chem.* **150**, 401.
- Tibbles, W. (1912). *Foods*. London: Baillière, Tindall and Cox.
- Webster, B. & Chesney, A. M. (1928). *Johns Hopk. Hosp. Bull.* **43**, 299.
- Webster, B. & Chesney, A. M. (1930). *Amer. J. Path.* **6**, 275.
- Webster, B. & Clawson, T. A. & Chesney, A. M. (1928). *Johns Hopk. Hosp. Bull.* **43**, 278.
- Webster, B., Marine, D. & Cipra, A. (1931). *J. exp. Med.* **53**, 81.