REFERENCES

- Beiler, J. M. & Martin, G. J. (1948). *J. biol. Chem.* **174**, 31.
 Beilinson, A. (1929). *J. biol. Méd. exp. Moscou*, **11**, 52; cited from *Chem. Abstr.* (1930), **24**, 1427.
 Bournsnel, J. C., Dangerfield, W. G. & Wormall, A. (1939). *Biochem. J.* **33**, 81.
 Bournsnel, J. C. & Wormall, A. (1939). *Biochem. J.* **33**, 1191.
 Curd, F. H. S. & Davey, D. G. (1949). *Nature, Lond.*, **163**, 89.
 Dangerfield, W. G., Gaunt, W. E. & Wormall, A. (1938). *Biochem. J.* **32**, 59.
 Dewey, H. M. & Wormall, A. (1946). *Biochem. J.* **43**, 24.
 Hawking, F. (1939). *Ann. trop. Med. Parasit.* **33**, 13.
 Hawking, F. (1940). *Trans. Roy. Soc. trop. Med. Hyg.* **34**, 37.
 Issekutz, B. (1933). *Arch. exp. Path. Pharmac.* **173**, 499.
 Jancsó, N. & Jancsó, H. (1934). *Z. Bakt. (1 Abt. Orig.)*, **132**, 257; cited from *Trop. Dis. Bull.* (1935), **32**, 22.
 Kabat, E. A. & Mayer, M. M. (1948). *Experimental Immunology*. Springfield, Illinois: Charles C. Thomas.
 Mackworth, J. K. (1948). *Biochem. J.* **42**, 82.
 Markham, R. (1942). *Biochem. J.* **36**, 790.
 Quastel, J. H. (1931). *Biochem. J.* **25**, 1121.
 Stuber, B. & Lang, K. (1926). *Biochem. Z.* **179**, 70.
 Sumner, J. B. & Somers, G. F. (1947). *Chemistry and Methods of Enzymes*, 2nd ed. New York: Academic Press Inc.
 Tcherniakofsky, P. & Nattan-Larrier, R. (1937). *C.R. Soc. Biol., Paris*, **126**, 1070.
 Town, B. W., Wills, E. D. & Wormall, A. (1949a). *Nature, Lond.*, **163**, 735.
 Town, B. W., Wills, E. D. & Wormall, A. (1949b). *Nature, Lond.*, **164**, 233.
 Town, B. W. & Wormall, A. (1949). *Biochem. J.* **44**, xxxviii.
 Wills, E. D. & Wormall, A. (1949). *Biochem. J.* **44**, xxxix.
 Wills, E. D. & Wormall, A. (1950). *Biochem. J.* **47**, 158.
 Wilson, E. J. & Wormall, A. (1949). *Biochem. J.* **45**, 224.

Studies on Suramin

9. THE ACTION OF THE DRUG ON SOME ENZYMES

By E. D. WILLS AND A. WORMALL

Department of Biochemistry and Chemistry, Medical College of St Bartholomew's Hospital, London

(Received 21 December 1949)

Evidence is available that suramin inhibits a few enzymes (for a review of the literature see Town, Wills, Wilson & Wormall, 1950). For reasons given in the preceding paper (Town *et al.* 1950) we have studied the action of very small amounts of the drug on a considerable number of enzymes under conditions as closely related as possible to the physiological. This paper is concerned mainly with experiments on non-proteolytic enzymes, with a few observations on the milk-clotting enzymes.

The primary object of the experiments described here was to find out whether suramin, in amounts comparable with those which are sufficient to destroy trypanosomes in the blood and tissues of man and other animals, has any inhibitory effect on enzymes likely to be present in the trypanosome. Where marked enzyme inhibition occurs, a study has been made of the mechanism of the inhibition and the conditions under which the drug reacts with the enzyme. In view of the importance of carbohydrate for the nutrition of trypanosomes, special attention has been paid to those enzymes taking part in carbohydrate metabolism, i.e. the enzyme systems concerned with yeast fermentation and muscle metabolism. In this connexion it is of interest to note that Marshall (1948) has recently found that trivalent arsenicals and straight-chain diamidines inhibit

certain enzyme systems concerned with carbohydrate metabolism in *Trypanosoma evansi*, and he has shown that the breakdown of glucose by trypanosomes follows almost exactly the pattern of glucose metabolism by yeast.

EXPERIMENTAL

Materials

Urease. (a) *Soya bean extract.* Soya bean flour (British Drug Houses) (1 g.) was extracted with 95 ml. of water and the mixture centrifuged; the clear solution was mixed with 5 ml. of 0.2 M-acetate buffer (pH 5.0) and centrifuged; the clear supernatant solution was used for the tests. (b) *Crystalline jack bean urease.* This was prepared as described by Sumner & Somers (1947); a solution containing 9 mg. of protein/l. was used.

Succinic oxidase, succinic dehydrogenase and cytochrome oxidase. The preparation of these enzyme systems from pig heart was carried out as described by Straub (1939) as far as the precipitation at pH 4.6. The precipitate was suspended in 5 vol. of 0.1 M-phosphate (pH 7.4) and 1 ml. of the mixture used for each test.

Yeast juice. For different experiments baker's yeast, brewer's top yeast and brewer's bottom yeast were used. After it had been well washed with water, the yeast (100 g.) was mixed with kieselguhr (50 g.) and washed sand (100 g.), and the mixture ground to a dough in a heavy mechanical mortar. The dough was kept at room temperature for

30 min. and then squeezed in a hand press. The juice contained all the factors necessary for the fermentation of glucose.

Hexokinase. This enzyme was prepared from 500 g. of yeast by using the first stage of the method of Bailey & Webb (1948), with precipitation by ethanol at -5° instead of by $(\text{NH}_4)_2\text{SO}_4$. The precipitate was centrifuged, dried and dissolved in water (5 mg./ml.) just before use.

Choline dehydrogenase. Following the method of Mann & Quastel (1937), fresh rat liver (10 g.) was ground with sand (5 g.) and 0.2M-phosphate, pH 7.4 (30 ml.), and the extract strained through muslin and then centrifuged. The undiluted supernatant solution was used in the tests.

Arginase and D-amino-acid oxidase. The extract of rat liver prepared as described under choline dehydrogenase was used as a source of these two enzymes.

Catalase. (a) *From blood.* Washed red blood cells were haemolysed with water, and the cell debris removed by centrifuging; the supernatant solution, suitably diluted (usually corresponding to about a 250-fold dilution of the red cells), was used in the tests. (b) *From liver.* Beef liver was sliced and washed thoroughly to remove as much blood as possible. It was then ground with sand and 1 vol. of water. The mixture was squeezed through muslin and the centrifuged extract was diluted about 1000-fold.

Amylases. Malt α - and β -amylases were prepared by a method based on that of Ohlsson (1926, 1930). Fresh malt was ground with water and the extraction allowed to continue for 30 min. For the preparation of α -amylase the reaction of the crude extract was adjusted to pH 6.5 and the mixture was kept at 70° for 15 min. and then cooled quickly. (The β -amylase is destroyed by this treatment.)

The β -amylase was prepared by adjusting the reaction of the crude malt extract to pH 3.3 and maintaining it at that point for 15 min., followed by adjustment to pH 6.0. (This treatment destroys nearly all the α -amylase.)

Xanthine oxidase. The cream of unpasteurized milk or the whey preparation described by Dixon & Kodama (1926) was used.

Peroxi-dase. Minced horse-radish (40 g.) was extracted with water (250 ml.) at 0° overnight and the extract was strained through muslin. It was diluted, usually with about 20 vol. of water, before use.

Carbonic anhydrase. Washed sheep or ox red blood cells (5 ml.), water (3 ml.) and ethanol (2 ml.) were well mixed, and CHCl_3 (0.5 ml.) was then added. The mixture was shaken for 0.5 min., and centrifuged. The supernatant solution, filtered if necessary, was diluted with 4 vol. of water before use.

Ascorbic acid oxidase. This was prepared from cucumber and cabbage as described by Johnson & Zilva (1937).

Glyoxalase. Fresh beef liver (70 g.) was minced and extracted with water (50 ml.) overnight at about 4° . The extract was squeezed through muslin and centrifuged; the supernatant solution was diluted 50-fold before use.

Carboxylase (yeast). Two enzyme solutions were used in these tests: (a) yeast juice prepared as described above and then suitably diluted, and (b) a preparation made from dried yeast using the first stage of the method of Green, Herbert & Subrahmanyam (1941).

Invertase. Dilute solutions of several commercial samples of invertase were used.

Emulsin (β -glucosidase). This was prepared from dried defatted almond meal (Sumner & Somers, 1947).

Papain. A 0.5% (w/v) aqueous solution of commercial papain (Harrington Bros.) was used.

Rennin. 'Essence of Rennet' (Benger's) was diluted with 9 vol. of water.

Pepsin. 'Liquor Pepticus' (Allen and Hanburys) was diluted with 4 vol. of water.

Trypsin. The tests employed involved the measurement of the milk-clotting power. Although it has a high proteolytic activity, the enzyme preparation used (Liquor Trypsin Co.; Allen and Hanburys) has a low clotting power, and it was therefore used undiluted.

Buffer solutions. The following buffers were used: 0.2M-Na acetate-acetic acid (Clark and Lubs), 0.2M-phosphate (KH_2PO_4 plus NaOH: Clark and Lubs), 0.2M-boric acid-KCl-NaOH and 1% (w/v) NaHCO_3 gassed with CO_2 to pH 7.4.

Methods

The methods used for the measurement of enzyme activities in the experiments described in this paper were mainly manometric. The details given below refer to the control experiments. Where suramin was added, the water content of the rest of the mixture was reduced by an amount equal to the volume of the suramin solution. The concentration of suramin recorded in these inhibition tests is that in the reaction mixture just before the addition of the substrate. The enzyme-suramin mixtures (and the controls) were kept at the temperature at which the enzyme tests were made, before the addition of the substrate; the time of contact of suramin and enzyme before the addition of substrate ranged from 10 to 60 min. but was usually 20–30 min. All enzyme tests were carried out at 37° except where otherwise mentioned, and the mixtures were buffered; where no pH is specified below, the tests were carried out over a range of pH.

The extent of inhibition in all the experiments described in this paper was calculated from the first steady rates of the reactions in the suramin-containing mixture and the control experiment.

Urease. (a) *Manometric*, by the estimation of CO_2 liberated. The main compartment of the Warburg apparatus contained enzyme solution (1 ml.), 0.2M-acetate of pH 5.0 unless otherwise stated (1 ml.) and water (1 ml.). The substrate was 0.25% (w/v) urea solution (0.2 ml.). (b) *Conway method.* A mixture of urea solution (0.2 ml. of a 0.25% solution), the enzyme (0.5 ml.) and buffer (0.5 ml. of 0.1M-phosphate of pH 7.4) was placed in the outer compartment of the Conway unit and 1 ml. of 1% boric acid plus mixed indicator in the centre compartment. The dish was kept at 37° for a standard period (usually 15 min.) and saturated K_2CO_3 solution (1 ml.) added to the outer compartment. After a further 1 hr. at 37° the boric acid was titrated with N/150 HCl.

Succinic oxidase system. The enzyme suspension (1 ml.), 0.1M-phosphate, pH 7.4 (1 ml.) and water (1 ml.) were placed in the main compartment of the Warburg flask and 0.2 ml. of 20% (w/v) KOH in the centre cup. The reaction was started by the addition of 2% (w/v) Na succinate (0.3 ml.) from the side bulb and the enzyme action measured by the O_2 uptake.

Succinic dehydrogenase. The method was the same as that for succinic oxidase, with the addition of neutralized 5% (w/v) KCN solution (0.1 ml.), to poison the cytochrome oxidase, and 0.5% methylene-blue solution (0.3 ml.), to link the system directly to oxygen.

Cytochrome oxidase; as for succinic oxidase with 1% *p*-phenylenediamine (0.3 ml.) instead of the succinate, to reduce the cytochrome. The rate of re-oxidation of cytochrome by cytochrome oxidase was measured by following the O_2 uptake.

Cholinesterase. This enzyme action was studied manometrically by following the acid production as recorded by the liberation of CO_2 . The enzyme (0.5 ml. of horse serum), 1% $NaHCO_3$ (gassed with CO_2 to give pH 7.4) (0.5 ml.) and water (1 ml.) were placed in the main compartment, and 0.5% acetylcholine solution (0.4 ml.) in the side bulb of the Warburg flask. The gas phase was 95% N_2 plus 5% CO_2 .

Tyrosinase. Activity was measured by following O_2 uptake at both 20° and 37°. The enzyme (1 ml. of freshly prepared potato juice), 0.2M-buffer (1 ml.) and water (1 ml.) were placed in the main part of the Warburg flask and the following substrates were used: (i) saturated tyrosine solution (0.5 ml.), (ii) 2.5% *p*-cresol (0.3 ml.), and (iii) 2.5% catechol solution (0.3 ml.).

Whole yeast. The fermentation of glucose (0.2 ml. of a 6% solution) by a mixture of a 2% suspension of baker's yeast (1 ml.), 0.1M-phosphate, pH 6.0 (1 ml.) and water (1 ml.) was measured by the CO_2 production.

Yeast juice. The method was the same as that for whole yeast. The substrates used were 2–6% solutions (0.2 ml.) of glucose, fructose, sucrose and maltose.

Hexokinase. (a) *Using yeast juice as the enzyme source and measuring the utilization of glucose*. A mixture of yeast juice (1 ml.), 0.1M-phosphate, pH 6.0 (1 ml.) and water (1.3 ml.) was kept in a test tube at 37° for 15 min. Glucose (1 ml. of a 0.2% solution) was then added, and samples (1 ml.) were withdrawn from the mixture after 0, 3, 8 and 15 min. Each sample was quickly blown into a mixture of water (3 ml.) and 0.3N- $Ba(OH)_2$ (0.75 ml.) to precipitate phosphate esters, and 2 min. later 5% (w/v) $ZnSO_4$ solution (0.75 ml.) was added to precipitate proteins (cf. Nelson, 1944). The mixture was centrifuged and the glucose content of 1 ml. of the supernatant solution was determined by the method of Haslewood and Strookman as described by King (1946), using a photoelectric colorimeter.

(b) *Using a partially purified hexokinase preparation* (cf. Bailey & Webb, 1948). A mixture of the enzyme solution (0.3 ml.), prepared as described above (p. 159), bicarbonate, pH 7.4 (2.6 ml.), 0.1M- $MgSO_4$ (0.5 ml.) and 2% glucose solution (0.6 ml.) was kept at 37° for 15 min. ATP (Na salt: 0.4 mg. pyrophosphate-P/ml.) (0.5 ml.) was then added to start the reaction, and samples (1 ml.) were withdrawn after 0, 3, 8 and 15 min. Each sample was quickly blown into 2N-HCl (1 ml.) and the mixture heated at 100° for 7 min. The liberated inorganic P was then determined by the method of Fiske & Subbarow (1925), and the hexokinase activity was calculated from the decrease in the '7 min. hydrolysable' P compounds.

Choline dehydrogenase. Activity was measured by the O_2 uptake of a mixture of enzyme solution (1 ml.), 0.2M-phosphate, pH 7.4 (1 ml.) and water (1 ml.), after the introduction of 0.033M-choline chloride (0.4 ml.); the centre cup contained 20% (w/v) KOH (0.2 ml.).

D-Amino-acid oxidase. The method used was similar to that described for choline dehydrogenase, but with 5% DL-alanine (0.3 ml.) as substrate.

Arginase. The method of Weil & Russell (1934), involving the manometric estimation of urea production, was used. The main compartment of the Warburg flask contained enzyme solution (1 ml.), borate, pH 9.0 (0.3 ml.), 0.1M-

arginine (0.2 ml.) and water (0.7 ml.); urease solution (0.5 ml.) was placed in the side bulb. The manometer was kept at 37° for 15 min. and M-acetate buffer of pH 5.6 (1 ml.) was then added. After the manometer had been re-equilibrated, the urease solution was 'tipped in' and the total CO_2 liberated was measured. (Buffer of pH 5.6 was used because urease is not inhibited by suramin at this pH.)

Catalase. The enzyme solution (1 ml.), 0.2M-phosphate or acetate (1 ml.) and water (1 ml.) were kept at 20° in the Warburg apparatus for 20 min.; 0.27% (w/v) H_2O_2 (0.3 ml.) was then added from the side bulb, and the O_2 liberated measured.

Amylase. A mixture of enzyme solution (1 ml.), M-acetate (2 ml.) and water (12 ml.) was kept in a small flask at 37° for 20 min., and 1% starch solution (5 ml.) was then added. Samples (2 ml.) were withdrawn after 0, 2, 4, 6 and 8 min. and the liberated maltose was determined by the Somogyi-Shaffer-Hartmann method (as described by Hawk, Oser & Summerson, 1947). The samples were mixed with water (3 ml.) and Cu reagent (5 ml.), and the mixture heated in a boiling water bath for 15 min. and then cooled; 5N- H_2SO_4 (1 ml.) was added and free I_2 titrated with 0.01N-thiosulphate.

Xanthine oxidase. Activity was measured manometrically by the O_2 uptake with acetaldehyde or xanthine as substrate (cf. Ball, 1939). A mixture of enzyme solution (1 ml.) or milk (2 ml.), 0.2M-phosphate or acetate (1 ml.) and water (1 ml.) was kept at 37° for 20 min. and 0.1M-acetaldehyde solution (0.4 ml.) or 0.05M-xanthine in 0.05N-NaOH (0.4 ml.) was then added.

Peroxidase. The manometric method involving the measurement of the CO_2 liberated from pyrogallol (Ettori, 1949) was used. A mixture of enzyme solution (1 ml.), 1% H_2O_2 (0.2 ml.), 0.2M-acetate (1 ml.) and water (0.8 ml.) was kept at 37° for 20 min. and 5.5% pyrogallol solution (0.3 ml.) added from the side bulb.

Carbonic anhydrase. The enzyme solution (1 ml.), 0.2M-phosphate, pH 6.8 (1 ml.) and water (1 ml.) were equilibrated with rapid shaking in the Warburg apparatus. The reaction was started by the addition of 0.5M- $NaHCO_3$ in 0.02N-NaOH (0.3 ml.), and the CO_2 liberation at 20° was recorded every 15 sec. for 1.5 min. A blank experiment (without enzyme) was carried out in every case.

Ascorbic acid oxidase. A mixture of the enzyme solution (1 ml.), 0.2M-phosphate or acetate (1 ml.) and water (1 ml.) was kept at 37° for 20 min. and a neutralized 1.25% ascorbic acid solution (0.3 ml.) was then added from the side bottle. The rate of O_2 uptake was recorded at intervals; a blank experiment was always made to give the rate of autoxidation of the ascorbic acid.

Glyoxalase. The enzyme solution (1 ml.), 2% $NaHCO_3$ (gassed with CO_2 to pH 7.4) (1 ml.), 0.1M-glutathione (0.1 ml.) and water (0.9 ml.) were kept with a gas phase of 95% N_2 /5% CO_2 at 37° for 20 min., and the reaction started by the addition of neutralized 3% methylglyoxal solution (0.2 ml.). The rate of CO_2 evolution was recorded.

Yeast carboxylase. The enzyme solution (1 ml.), 0.1M-phosphate (1 ml.) and water (1 ml.) were kept at 37° for 20 min., and 0.8% Na pyruvate (0.3 ml.) was then added. The rate of evolution of CO_2 was measured.

Invertase. The enzyme solution (1 ml.), M-acetate (2 ml.) and water (15 ml.) were kept in a small flask at 37° for 15 min., and 2% sucrose (2 ml.) was then added. Samples (2 ml.) were withdrawn after 0, 2, 4, 6 and 8 min. and

the reducing sugar determined by the Somogyi-Shaffer-Hartmann method as described for amylase.

β -Glucosidase. The enzyme solution (1 ml.), 0.2 M-acetate (3 ml.) and water (4 ml.) were kept in a small flask at 37° for 15 min., and 2% salicin (2 ml.) was then added. Samples (2 ml.) were taken after 0, 2, 4, 6 and 8 min. and the liberated glucose was determined by the Somogyi-Shaffer-Hartmann method as described under amylase.

Papain, rennin, pepsin and trypsin. In the experiments described in this paper these enzymes were studied by the method of Balls & Hoover (1937), which involves a measurement of the milk-clotting power. A mixture of the enzyme solution (1 ml.), 0.2 M-acetate, pH 4.8 (1 ml.) and water (0.5 ml.), plus 0.5 ml. of 3% cysteine hydrochloride solution in the case of papain, was kept at room temperature for 10 min. Milk (3 ml.) was then added, and the time required to give the first recognizable clotting was noted.

RESULTS

Inhibition of urease by suramin

Typical results of our experiments with soya bean extracts are shown in Fig. 1. With M/3000 suramin, contact of the drug and the enzyme at pH 5.0 for 30 min. causes almost complete loss of urease activity and M/15,000 suramin gives easily detectable inhibition.

Fig. 1. Inhibition of soya bean urease by suramin at pH 5.0.

The duration of contact of drug and enzyme before addition of the substrate is an important factor, and although the drug reacts to some extent immediately with enzymes, as with other proteins (Dewey & Wormall, 1946), maximum inhibition is usually observed when the drug-enzyme mixture is kept for 20–60 min. before addition of the substrate. In a typical experiment with urease, the inhibitory effect of the drug increased strongly during the first 30 min. of contact and reached a maximum in about 60 min.

(Table 1). No significant increase in inhibitory activity occurred after the enzyme had been exposed to the suramin for 2 hr.

Table 1. *Influence of time of contact on the inhibitory action of suramin on soya bean urease*

(The enzyme was exposed to M/5000 suramin at pH 5.0 and 37° for varying lengths of time before the addition of urea. For further details, see the text, p. 159.)

Duration of contact of urease and suramin before addition of urea (min.)	Inhibition (%)
0	41
15	50
30	80
60	93

Inhibition of urease by suramin with varying substrate concentrations. Experiments were carried out to determine the effect of varying the substrate concentration, and the results were examined by the method of Lineweaver & Burk (1934). Using a soya bean extract and concentrations of suramin ranging from M/2500 to M/5000, the results shown in Fig. 2

Fig. 2. Effect of substrate concentration in inhibition of urease at pH 5.0.

were obtained. These showed that when the reciprocal of the reaction velocity (v) is plotted against the reciprocal of the substrate concentration (S), all the curves intersect at a point on the vertical axis ($1/V$, where V is the maximum velocity). This is a criterion of competitive inhibition (Lineweaver & Burk, 1934).

Effect of pH on the inhibitory action of suramin on soya bean urease. Suramin is reported to have no effect on urease (Quastel, 1931), but no details were given about the pH at which the tests were made. We presumed that these were made at pH 7.4, the pH for urease tests, and decided, therefore, to make

a more extensive study of the effect of pH on the inhibition of urease by suramin. Tests at pH 7.4, involving the use of the Conway unit, gave results which completely confirmed those of Quastel; the manometric method cannot be used at that pH. Suramin had no inhibitory effect even in concentrations as high as $M/500$.

Using the manometric method for the measurement of urease activity we studied the inhibition over the range pH 4.8–5.3, and found that, whereas inhibition by $M/3000$ suramin was about 100% at pH 5.0, it was about 50% at pH 5.1, about 10% at pH 5.2 and zero at pH 5.3 (cf. Fig. 3). The change in

Fig. 3. Effect of varying pH on the inhibition of urease by suramin ($M/3000$).

the inhibitory effect occurs very near the isoelectric point of urease (pH 5.0–5.1 according to Sumner & Hand, 1929), and this observation led us to investigate more fully the possibility that other enzymes would show the same phenomenon. These experiments were briefly reported earlier (Wills & Wormall, 1949) and are described more fully by Wills (1950).

Inhibition of crystalline urease. By the courtesy of Prof. J. B. Sumner we obtained a supply of jack bean meal of high urease activity, and from this material we prepared the crystalline enzyme (cf. Sumner & Somers, 1947). As we had expected, this pure urease was much more sensitive to the action of suramin than crude soya bean extract. For example, $M/5,000,000$ and $M/30,000,000$ suramin gave, respectively, 90 and 53% inhibition of crystalline jack bean urease (Fig. 4). In a comparable experiment a concentration of $M/7500$ suramin was required to give 60% inhibition of a soya bean extract. Thus the crystalline enzyme was about 4×10^3 times as sensitive as was the crude preparation.

Experiments with crystalline urease showed that the inhibition with suramin is dependent on pH, and also that the inhibitory change effected at pH 5 is not

reversed when the solution is neutralized. It was reported that with crude urease regeneration of activity occurred when the enzyme inactivated by suramin at pH 5.0 was neutralized (Wills & Wormall, 1949), but subsequently we have been unable to

Fig. 4. Inhibition of crystalline urease by various suramin concentrations at pH 5.0.

Fig. 5. The effect on the reaction velocity of changing the inhibitor (suramin) concentration. Crystalline urease at pH 5.0. A, assuming $n=1$; B, assuming $n=2$.

confirm this with either pure or crude urease. All enzymes inhibited by suramin in acid solution do not, however, behave like urease in this respect, for the activity of invertase which has been inactivated by suramin at pH 3.8 is restored when the pH is adjusted to 5.2.

By adopting the method used by Eadie (1942) for cholinesterase and by Cori, Cori & Green (1943) for phosphorylase, the data obtained in our experiments with crystalline urease exposed to varying amounts of inhibitor (suramin) were used to calculate the number of molecules of inhibitor which combine with each molecule, or rather active centre, of urease. By rearrangement and adaptation of the usual Michaelis-Menten equation, the equation $1/v = a + bi^n$ (where v = the initial reaction velocity, a and b are constants, i is the molar concentration of inhibitor and n is the molecular ratio of inhibitor to enzyme) can be derived (Lineweaver & Burk, 1934; Eadie, 1942). When the reciprocal of the velocity is plotted against the square of the suramin concentration, a straight line is obtained, suggesting that $n = 2$, i.e. that two molecules of suramin are needed to block each active centre of urease (Fig. 5).

Effect of suramin on fermentation by yeast

As trypanosomes metabolize glucose with great avidity, we decided to study the action of suramin on yeast fermentation and on the various enzymes concerned with this process.

Whole yeast. Our experiments have shown that suramin in concentrations as high as $M/500$ has no effect on the anaerobic fermentation of glucose by whole yeast (baker's or brewer's) or the respiration, at pH 6.0. Furthermore, exposure of the yeast to $M/500$ suramin for 24 hr. affects neither the fermenting nor the respiratory capacities at pH 6.0. The proteins of yeast juice, like those of haemolysed red blood cells, readily combine with the drug.

Yeast juice. Suramin strongly inhibits the fermentation of glucose by yeast juice at pH 6.0 and 7.0 (Fig. 6), and similar inhibition occurs when fructose or maltose is used as substrate. Nearly all the yeast juice preparations used showed a steady autofermentation, presumably due to autogenous glycogen, and the autofermentation was likewise inhibited by the drug, but not as strongly as the fermentation of added glucose; thus $M/15,000$ suramin caused 42% inhibition of the autofermentation and 90% inhibition of the glucose fermentation. This difference suggested that one of the early stages in the utilization of glucose (e.g. the hexokinase action) is strongly inhibited by suramin.

The extent of the inhibition obtained in these experiments with yeast juice varied with different preparations of the juice, but every preparation tested was sensitive to the drug. Occasionally, 100% inhibition occurred with $M/60,000$ suramin, but a slightly higher concentration of the drug was usually necessary to give complete inhibition; 50% inhibition was usually obtained with $M/30,000$ suramin, and occasionally with $M/75,000$.

The above-described tests with yeast juice were carried out in the presence of phosphate buffer, and

replacement of this by an arsenate buffer gave interesting results (Fig. 7). The autofermentation process is apparently little affected by arsenate, for the inhibition effected by suramin is essentially the same as that observed when a phosphate buffer is used, but

Fig. 6. The effect of various suramin concentrations on yeast juice fermentation of glucose in phosphate buffer at pH 6.0. Glucose (0.2 ml. of a 6% solution) added at X.

Fig. 7. The action of suramin on yeast juice fermentation of glucose in arsenate buffer at pH 6.0. Glucose (0.2 ml. of a 6% solution) added at X.

on addition of glucose the response is quite different from that occurring in the comparable experiment with phosphate buffer. Fermentation in the control mixture (no suramin) slows down after addition of glucose and soon ceases, whereas the mixtures containing suramin are little affected, particularly those containing most suramin. In the mixture containing $M/10,000$ suramin, where the inhibition of

autofermentation was 90 %, the rate of fermentation after the addition of glucose actually became greater than that in the control.

It is considered likely that the explanation of these results with arsenate is connected with adenosinetriphosphate (ATP) formation (cf. Meyerhof & Junowicz-Kocholaty, 1942; Meyerhof, 1945). In the presence of arsenate, ATP production is retarded, and when excess of glucose is added to this system all the available ATP is rapidly used for the formation of glucose-6-phosphate by the action of hexokinase; lack of ATP will soon stop the fermentation. If suramin inhibits hexokinase, and our experiments

Fig. 8. The effect of the addition of ATP to yeast juice fermentation of glucose in arsenate buffer at pH 6.0. Glucose (0.2 ml. of a 6 % solution) either alone or with 0.2 ml. ATP (Na salt: 0.4 mg. pyro-P/ml.) added after 9 min. (X). Suramin concn. $m/10,000$.

described below show that this is true, the addition of glucose to the arsenate-containing juice will not be followed by increased utilization of ATP and the autofermentation can continue unimpaired. That this is the probable explanation has been confirmed by the results of an experiment in which extra ATP was added at the same time as the glucose. It was found (Fig. 8) that whereas the added ATP had no effect on the suramin-inhibited autofermentation, it allowed the autofermentation in the control mixture to continue at its normal rate for an appreciable time after the addition of glucose. These experiments with arsenate-buffered yeast fermentation strongly suggested, therefore, that inhibition of hexokinase is a very important factor in the inhibitory action of suramin on fermentation by yeast.

Effect of suramin on hexokinase

In view of the above-mentioned results we studied the action of suramin on hexokinase, using a partially purified preparation; we have also tested a hexokinase preparation, kindly provided by Dr M. Dixon, with results similar to those given by our own preparation. These experiments, the results of some of which are given in Table 2, showed that suramin strongly inhibits hexokinase. Of the two methods used for measuring the activity of the enzyme: (a) the rate of utilization of ATP, and (b) the rate of conversion of glucose to glucose-6-phosphate, the latter showed a slightly greater inhibition by suramin than the former.

Table 2. *Inhibition of hexokinase by suramin*

Suramin concn.	Inhibition (%)	
	Measured by ATP utilization	Measured by glucose utilization
$m/10,000$	80	100
$m/20,000$	70	80
$m/30,000$	60	75

Table 3. *Comparison of the inhibitory effects of suramin on hexokinase and on autofermentation and glucose fermentation by yeast juice*

(From the results of experiments showing the inhibitory effect of suramin on hexokinase (Table 2) and on yeast juice autofermentation (Fig. 6) the expected inhibition of glucose fermentation can be calculated. These calculated values and those obtained experimentally are given below.)

Suramin concn.	Inhibition (%) of			
	Hexokinase (a)	Autofermentation (b)	Glucose fermentation Calculated from (a) and (b)	Found
$m/10,000$	80	70	94	100
$m/20,000$	70	50	85	95
$m/30,000$	60	30	60	50

A comparison of the inhibition, by suramin, of autofermentation (presumably of glycogen), glucose fermentation and hexokinase action suggests that only about half the inhibitory effect of the drug on the total fermentation by yeast juice is due to inhibition of hexokinase (cf. Table 3). So far we have not fully studied all the yeast enzymes to find which are sensitive to suramin, but we have observed that carboxylase is inhibited by the drug. For example, $m/6000$ suramin gives 70 and $m/10,000$ gives 25 % inhibition of this enzyme at pH 6.0. The fairly high concentration of suramin required to inhibit yeast carboxylase, and the relatively low degree of inhibi-

tion suggest, however, that inhibition of carboxylase plays only a minor role in the inhibition of yeast fermentation by suramin.

Effect of suramin on some oxidizing enzymes

In view of the rapid oxygen uptake of trypanosomes, we have studied the effect of suramin on a variety of oxidizing enzymes; where the manometric method can be used and the enzyme is active at pH 7.4, the tests were made at this pH. Of the oxidizing enzymes tested only succinic dehydrogenase and choline dehydrogenase are inhibited to an extent likely to be of pharmacological interest, i.e. by concentrations of the drug likely to be attained in the blood or inside the trypanosome (Table 4).

Table 4. *Effect of suramin on some oxidizing enzymes*

Enzyme	pH	Suramin concn.	Inhibition (%)
D-Amino-acid oxidase	7.4	M/500	0
Xanthine oxidase	7.4	M/500	0
Peroxidase	6.0	M/500	0
Tyrosinase	6.0	M/500	0
Catalase	7.0	M/500	0
Succinic oxidase system	7.4	M/4000	75
Succinic dehydrogenase	7.4	M/4000	70
Choline dehydrogenase	7.4	M/5000	40
Ascorbic acid oxidase	7.4	M/1000	40
Cytochrome oxidase	7.4	M/3000	25

Effect of suramin on some non-oxidizing enzymes, at or about their optimum pH

Tests on a variety of non-oxidizing enzymes, made at or about the optimum pH of each enzyme, have shown that the majority of enzymes are unaffected by suramin under these conditions (Table 5). Glyoxalase is inhibited to the extent of 50 % by 0.001 M-suramin, but the inhibition of this enzyme is very slight with 'physiological' concentrations of the drug. However, the milk-clotting activities of commercial papain, rennin, trypsin and pepsin are fairly strongly inhibited by M/6000-M/10,000 suramin at pH 4.8 (Table 6). This inhibition may be related to the inhibition of the proteolytic action of trypsin at pH 8.9 (Town & Wormall, 1949), or it may be connected with a reaction between the drug and the substrate. It is also important to remember that suramin is a powerful inhibitor of blood clotting.

The specificity of enzyme inhibition with suramin

In order to obtain more information about the specificity of this inhibition, a study was made of the effect of suramin on each of three enzymes or enzyme systems present in an aqueous extract of rat liver. It was found (Table 7) that the drug strongly inhibited the choline dehydrogenase (at pH 7.4), but had no detectable effect on the D-amino-acid oxidase (at pH 7.4) or arginase (at pH 8.9) in the extract.

Table 5. *Some enzymes not inhibited by M/500 suramin at or near their optimum pH's*

Enzyme	pH
β -Glucosidase	5.5
Invertase	4.8
Carbonic anhydrase	6.8
Liver esterase	6.8
Cholinesterase	7.4
Arginase	8.9
α -Amylase (malt)	5.5
Urease	7.4
D-Amino-acid oxidase	7.4
Xanthine oxidase	7.4
Peroxidase	6.0
Catalase	7.0
Tyrosinase	6.0

Table 6. *Inhibition of milk-clotting enzymes by suramin*

(All the tests were made at pH 4.8.)

Enzyme	Suramin concn.	Inhibition (%)
Papain	M/5000	67
	M/10,000	43
Rennin	M/3000	87
	M/10,000	50
Trypsin	M/3000	65
	M/6000	45
Pepsin	M/3000	75
	M/6000	60

Table 7. *Specificity of enzyme inhibition by suramin, as shown by the effect of the drug on different enzymes in a rat-liver preparation*

Enzyme	pH	Suramin concn.	Inhibition (%)
Choline dehydrogenase	7.4	M/5000	40
D-Amino-acid oxidase	7.4	M/500	0
Arginase	8.9	M/500	0

Effect of some suramin analogues on urease and the succinic oxidase system

The inhibitory effect of some suramin analogues has been tested with urease and the succinic oxidase system, enzymes which are very sensitive to the action of suramin. Some typical results of these tests with compounds containing sulphonic acid groups are given in Table 8 and with those not containing sulphonic acid groups in Table 9.

The two enzyme systems studied show considerable differences in their sensitivity to these suramin analogues. Slight changes in the suramin structure do not appreciably reduce the ability of the compound to inhibit the two enzymes, but a marked reduction in the size of the molecule reduces very considerably the inhibitory effect towards urease but not towards the succinic oxidase system.

From a study of the results with urease, the following conclusions have been reached: (i) The sulphonic acid groups are essential for urease inhibition by

compounds which are structurally related to suramin. (ii) The methyl groups of suramin are not essential for this inhibition. (iii) Suramin analogues containing the intact end groups of suramin, but

capacity. (v) The capacity to inhibit urease is not a common property of all substituted ureas of the suramin type. The action of suramin in inhibiting the succinic oxidase system is apparently quite different

Table 8. *Effect, on urease and the succinic oxidase system, of some suramin analogues containing sulphonic acid groups*

(The results given are those with the following concentrations of the compound tested. Urease, $\text{m}/5000$; succinic oxidase, (a) $\text{m}/4000$; (b) $\text{m}/1000$. The urease tests were made at pH 5.0, and the succinic oxidase tests at pH 7.4.)

Formula of compound ($\text{X}=\text{SO}_3\text{Na}$)	Name of compound	Urease	Inhibition (%) of	
			Succinic oxidase	
			(a)	(b)
Suramin	80	75	95
Demethylated-suramin	73	—	—
82	60	90
28	68	90
1-Naphthylamine-4:6:8-trisulphonic acid	10	0	60

having fewer benzene rings in the molecule, are appreciably less inhibitory than is suramin. (iv) A slight alteration in the molecule involving the replacement of $-\text{NH}-\text{CO}-$ by $-\text{NH}-\text{SO}_2-$ does not necessarily reduce the enzyme-inhibiting

from that on urease, though here also the sulphonic acid groups appear to be essential. The spatial arrangement and the size of the molecule do not, however, seem to be so important in the inhibition of the succinic oxidase system.

Table 9. *Effect, on urease and the succinic oxidase system, of some suramin analogues not containing sulphonic acid groups*

(Experimental details as in Table 8, except that the tests with urease were made with $m/3000$ solutions of the compounds studied.)

Compound	Urease	Inhibition (%) of Succinic oxidase	
		(a)	(b)
$\left(\begin{array}{c} \text{HOOC}-\text{C}_6\text{H}_3(\text{CH}_3)-\text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}- \\ \text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}- \end{array} \right)_2 \text{CO}$	20	0	0
$\text{HOOC}-\text{C}_6\text{H}_3(\text{CH}_3)-\text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}_2$	0	—	—
$\left(\begin{array}{c} \text{Cl}^-(\text{CH}_3)_3 \text{N}^+ -\text{C}_6\text{H}_3(\text{CH}_3)-\text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}- \\ \text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}- \end{array} \right)_2 \text{CO}$	7	0	65
$\left(\begin{array}{c} \text{Cl}^-(\text{CH}_3)_3 \text{N}^+ -\text{C}_6\text{H}_3(\text{CH}_3)-\text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}-\text{CO}-\text{C}_6\text{H}_3(\text{CH}_3)-\text{NH}- \\ \text{NH}-\text{CO}-\text{C}_6\text{H}_4-\text{NH}- \end{array} \right)_2 \text{CO}$	0	0	30

Protection of enzymes against inhibition by suramin

Under certain conditions proteins, by competing with the enzyme for the added drug, protect enzymes such as urease and succinic dehydrogenase against the inhibiting effect of suramin. Urease, for example, is protected at pH 5.0 by the serum proteins and by egg albumin (Table 10). At pH 7.4, however, the succinic oxidase system is not protected by egg albumin though it is protected by the serum proteins. The protective effect of proteins may depend on the pH of the medium in relation to the isoelectric point of the protein (pH 4.8 for egg albumin and about or slightly above pH 7.0 for serum γ -globulin). The non-enzymic protein in a crude enzyme preparation likewise reduces the sensitivity of an enzyme, as is shown by the fact that crystalline urease is about 4000 times as sensitive as is a crude preparation of the enzyme. Protection is also given by a large excess of urea, for the inhibition of urease by suramin is apparently competitive.

We have also studied the effect of adding certain thiol compounds, for both urease and succinic dehydrogenase are SH-enzymes. Tests were also made

with arginine, histidine and lysine, to see whether these basic amino-acids would reduce the inhibition by combining with the sulphonic acid groups of suramin. The results of these experiments showed that no protection of urease at pH 5.0 is given by cysteine (0.01M) or arginine (0.01M), and that 0.01M-cysteine, glutathione, arginine, histidine and lysine do not protect the succinic oxidase system against the inhibitory effect of $m/3000$ suramin at pH 7.4. It thus appears unlikely that the inhibition of these two enzymes by suramin is due to the combination of the drug with essential SH groups of the enzyme molecule or simply to the inactivation of an essential basic group in the active centre of the enzyme.

Influence of pH on the inhibition of enzymes by suramin and a method for the determination of isoelectric points of some enzymes

Although most enzymes are not significantly inhibited by suramin at or about their optimum pH, many of them behave like urease in being strongly inhibited at a lower pH. All the enzymes included in the list in Table 5 could not be tested over a suitable

pH range, for some are inactive in acid solution, but of those tested tyrosinase, the catalases, the amylases, peroxidase, carbonic anhydrase, invertase and β -glucosidase behave like urease; they are strongly inhibited by suramin in acid solution and they all give inhibition-pH curves of the same type as that given by urease (Fig. 3). Curves of similar nature were obtained by Quastel & Yates (1936) in experiments with invertase and several acid dyes.

Table 10. *Effect of proteins in protecting urease and succinic dehydrogenase against the inhibitory action of suramin*

(Varying amounts of rabbit serum (RS) or a solution of crystalline egg albumin (EA) were added to suramin solutions and the mixtures allowed to stand for 5–10 min. before addition of the enzyme. The enzyme tests were made at pH 5.0 for urease and 7.4 for the succinic oxidase system. The egg albumin solution had an inhibitory effect on both enzymes, but not sufficient to interfere with the tests.)

Suramin concn.	Added protein (% concn. in final mixture)	Inhibition of	
		Urease (%)	Succinic oxidase system (%)
m/3000	0	80	87
	0.42 EA	40	76
	1.05 EA	—	95
	2.1 EA	—	87
m/6000	0	50	72
	0.42 EA	31	72
	2.1 EA	—	55
m/3000	0	80	87
	0.36 RS	—	68
	0.55 RS	33	—
	0.91 RS	—	50
m/6000	0	50	72
	0.36 RS	—	54
	0.55 RS	6	—
	0.91 RS	—	20

Our experience of these inhibition experiments with suramin suggests that, where an enzyme is inhibited by suramin in acid solution and not at a pH which is on the alkaline side of the isoelectric point of the enzyme, a sharp drop in the inhibition-pH curve enables one to establish the isoelectric point. The method is quite simple and it does not require a pure preparation of the enzyme. Our crude and crystalline ureases, for example, gave the same isoelectric point by this method, and we have also found that the addition of a relatively large amount of 'foreign' protein, e.g. serum proteins or egg albumin, to an enzyme has no detectable effect on the inhibition-pH curve in the presence of suramin. The use of this method for the determination of isoelectric points of some enzymes has been briefly reported (Wills & Wormall, 1949, 1950) and full experimental details are given elsewhere (Wills, 1950).

DISCUSSION

The results of our investigations on enzyme inhibition by suramin suggest that the enzymes we have studied can be divided into two main groups as follows: Group *A*. This includes the enzymes which, like urease, invertase, peroxidase and catalase, are not inhibited at about pH 7, but are inhibited on the acid side of their isoelectric points. Group *B*. This includes enzymes which are strongly inhibited by the drug in neutral solution; hexokinase, carboxylase, succinic dehydrogenase and choline dehydrogenase are of this type. In these cases the inhibition is largely independent of the pH of the medium. The two types of inhibition appear to show considerable differences and it is believed that two different mechanisms are involved.

We have studied the inhibition of urease more extensively than that of any other enzyme, and urease may be taken to be typical of our group *A* enzymes. For various reasons it seems likely that suramin does not combine directly with the active centre of the enzyme and we suggest that the inactivation involves the combination or association of the two naphthylaminetrisulphonic acid groups of suramin with basic groups of urease lying near and perhaps on opposite sides of the active centre. The resulting combination will form a bridge with the large suramin molecule protecting the active centre of the enzyme, as shown diagrammatically in Fig. 9.

Fig. 9. Schematic illustration of the possible mode of action of suramin on enzymes classified in group *A*. *C*, active centre or centres of the enzymes; *S*, substrate molecule.

The experimental evidence which supports the theory of a 'bridging' mechanism to explain the inhibition of group *A* enzymes by suramin, rather than a direct action of the drug on the enzyme active centres, may be summarized as follows: (i) The effect of a slight change of pH near the isoelectric point of the enzyme has a profound effect on the degree of inhibition caused by suramin, for example, the inhibition of urease by m/3000 suramin is 100, 50 and 10% at pH's 5.0, 5.1 and 5.2, respectively. In the absence of suramin this change of pH has little effect on the activity of the enzyme and the properties of the active centre are apparently not appreciably altered by change of pH from 5.0 to 5.2. A change in the capacity of basic groups to combine with naph-

thylaminetrisulphonic acid groups, as a result of this change of hydrogen-ion concentration, can more readily be envisaged. (ii) The inhibition is not due solely to the sulphonic acid groups of suramin, but it also depends to a very considerable extent on the size of the molecule. Our tests with suramin analogues have shown that a reduction in the length of the chain between the two naphthylaminetrisulphonic acid groups appreciably reduces the capacity of the molecule to inhibit urease. The sulphonic acid groups are essential for this type of inhibition, but if this were the only factor concerned, i.e. if this group combined with the active centre of urease, we should expect many of the suramin analogues to be much more powerful inhibitors of urease than they were found to be by actual experiment. From the results obtained with urease and other enzymes and from our general study of the combination of suramin with proteins, we believe that the shape of the suramin molecule is such that one molecule readily combines with two basic groups of certain protein molecules in such a way as to form a relatively stable complex. When this 'bridge' covers the active centre of an enzyme, the enzymic activity will be inhibited, but inhibition of this type will only occur on the acid side of the isoelectric point of the enzyme.

With the enzymes of group *B* there is not the same dependence on pH, and the enzyme may even be strongly inhibited on the alkaline side of its isoelectric point. It is not possible to obtain reliable values for the isoelectric points of all the enzymes of this group, but in the case of hexokinase and carb-oxylase there is definite inhibition by suramin at pH 7.4 and 6.0 respectively, whereas the generally accepted isoelectric points for these two enzymes are 4.5-4.8 and 5.1 respectively.

Inhibition of group *B* enzymes by suramin thus appears to be due to a mechanism quite different from that suggested for urease and other enzymes of group *A*. With the group *B* enzymes our evidence suggests that suramin combines with the active centre of the enzyme. The spatial configuration of the inhibitor will be of importance in inhibition of this type, but the prime factor will be the presence of a group which can react with the enzyme active centre. The results of our tests with the various analogues of suramin are certainly consistent with this view. Succinic dehydrogenase is inhibited fairly strongly by many sulphonic acid-containing suramin analogues which have only a slight inhibitory effect on urease, and also the size of the inhibitor molecule does not appear to be critical; for example, 1-naphthylamine-4.6:8-trisulphonic acid strongly inhibits succinic oxidase (Table 8) and so do some other suramin analogues which have no appreciable effect on urease (Table 9).

Inhibition by suramin of the activity of enzymes of

group *B* is, of course, most interesting from the pharmacological viewpoint, for this inhibition is effective at pH 7.5. Marked inhibition of these enzymes can be demonstrated with very low concentrations of suramin: e.g. $m/30,000$ gives 75% inhibition of hexokinase and $m/4000$ gives 75% inhibition of succinic dehydrogenase.

Since many enzymes are inhibited by suramin only on the acid side of the isoelectric point of the enzyme and since the inhibition-pH curve in these cases usually falls sharply at about the isoelectric point, we have studied the possible use of this phenomenon for the determination of enzyme isoelectric points. Where the method is applicable the inhibition usually falls sharply from 100 to less than 10% over a range of about 0.2-0.3 of a pH unit, and the middle of this range is taken to be the isoelectric point of the enzyme; where data are available for the pure enzymes the values we have obtained by this method are in remarkably good agreement with the recorded isoelectric points.

SUMMARY

1. Suramin strongly inhibits urease at pH 5.0. $m/3000$ suramin completely inactivates this enzyme in soya bean extracts, and $m/30,000,000$ suramin strongly inhibits the crystalline jack bean enzyme. This inhibition appears to be competitive. Suramin does not inhibit urease at pH 5.3 or at higher pH's.

2. The fermentation of glucose by yeast juice, but not that by whole yeast, is strongly inhibited by $m/15,000$ and occasionally by $m/60,000$ suramin. About half the inhibitory effect of suramin on fermentation by yeast juice is due to inhibition of hexokinase. This enzyme is strongly inhibited by $m/30,000$ suramin at pH 6.0 or 7.5.

3. Suramin is not a general enzyme poison and most of the non-proteolytic enzymes we have tested are unaffected by the drug at pH 7.5 or at the optimum pH of the enzyme concerned. Of the oxidizing enzymes tested only the succinic oxidase system, succinic dehydrogenase and choline dehydrogenase are inhibited by concentrations of suramin equal to those found in the blood after the usual therapeutic doses.

4. Where enzyme inhibition by suramin occurs it is quite specific and the addition of the drug to a rat-liver extract inhibits choline dehydrogenase, but not D-amino-acid oxidase or arginase.

5. Other proteins strongly protect urease and other enzymes against the action of suramin, but no protection is afforded by basic amino-acids or by thiol compounds.

6. Tests with suramin and some suramin analogues suggest that inhibition of succinic oxidase may be due to direct combination of the inhibitor

with the active centre of the enzyme, but in the case of urease the suramin appears to act by 'bridging' across the active centre. In both cases the sulphonic acid groups of suramin are concerned with the reaction with the enzyme.

7. Many enzymes are, like urease, inhibited by suramin in acid solution but not in neutral or alkaline solution. In these cases the inhibition-pH curve usually falls sharply at about the isoelectric point of the enzyme.

The authors are pleased to express their indebtedness to the Medical Research Council and the Central Research Fund of The University of London for grants which have defrayed part of the expenses of these investigations. Thanks are also tendered to Dr M. Dixon, F.R.S., for samples of hexokinase and ATP; to Dr A. Spinks (of Messrs Imperial Chemical Industries Ltd.) for samples of suramin analogues; to Prof. J. B. Sumner of Cornell University, New York, for a generous gift of jack bean meal of high urease activity; and to Miss Barbara Hume, for help with a few of the enzyme tests.

REFERENCES

- Bailey, K. & Webb, E. C. (1948). *Biochem. J.* **42**, 60.
 Ball, E. G. (1939). *J. biol. Chem.* **128**, 51.
 Balls, A. K. & Hoover, S. R. (1937). *J. biol. Chem.* **121**, 737.
 Cori, C. F., Cori, G. T. & Green, A. A. (1943). *J. biol. Chem.* **151**, 39.
 Dewey, H. M. & Wormall, A. (1946). *Biochem. J.* **40**, 119.
 Dixon, M. & Kodama, K. (1926). *Biochem. J.* **20**, 1104.
 Eadie, G. S. (1942). *J. biol. Chem.* **146**, 85.
 Ettori, J. (1949). *Biochem. J.* **44**, 35.
 Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
 Green, D. E., Herbert, D. & Subrahmanyam, V. (1941). *J. biol. Chem.* **138**, 327.
 Hawk, P. B., Oser, B. L. & Summerson, W. H. (1947). *Practical Physiological Chemistry*, 12th ed. London: Churchill.
 Johnson, S. W. & Zilva, S. S. (1937). *Biochem. J.* **31**, 438.
 King, E. J. (1946). *Micro-analysis in Medical Biochemistry*, p. 20. London: Churchill.
 Lineweaver, H. & Burk, D. (1934). *J. Amer. chem. Soc.* **56**, 658.
 Mann, P. J. G. & Quastel, J. H. (1937). *Biochem. J.* **31**, 869.
 Marshall, P. B. (1948). *Brit. J. Pharmacol.* **3**, 8.
 Meyerhof, O. (1945). *J. biol. Chem.* **157**, 105.
 Meyerhof, O. & Junowicz-Kocholaty, R. (1942). *J. biol. Chem.* **145**, 443.
 Nelson, N. (1944). *J. biol. Chem.* **153**, 375.
 Ohlsson, E. (1926). *C.R. Lab. Carlsberg*, **16**, 1.
 Ohlsson, E. (1930). *Hoppe-Seyl. Z.* **189**, 17.
 Quastel, J. H. (1931). *Biochem. J.* **25**, 1121.
 Quastel, J. H. & Yates, E. D. (1936). *Enzymologia*, **1**, 60.
 Straub, F. B. (1939). *Biochem. J.* **33**, 787.
 Sumner, J. B. & Hand, D. B. (1929). *J. Amer. chem. Soc.* **51**, 1255.
 Sumner, J. B. & Somers, G. F. (1947). *Chemistry and Methods of Enzymes*, 2nd ed. New York: Academic Press Inc.
 Town, B. W., Wills, E. D., Wilson, E. J. & Wormall, A. (1950). *Biochem. J.* **47**, 149.
 Town, B. W. & Wormall, A. (1949). *Biochem. J.* **44**, xxxviii.
 Weil, L. & Russell, M. A. (1934). *J. biol. Chem.* **106**, 505.
 Wills, E. D. (1950). *Biochem. J.* (in the Press.)
 Wills, E. D. & Wormall, A. (1949). *Biochem. J.* **44**, xxxix.
 Wills, E. D. & Wormall, A. (1950). *Nature, Lond.*, **165**, 813.

Biochemical Studies of Toxic Agents

2. THE METABOLISM OF 2-NAPHTHYLAMINE AND 2-ACETAMIDONAPHTHALENE

By L. A. MANSON* AND L. YOUNG†
Department of Biochemistry, University of Toronto

(Received 6 March 1950)

Exposure to certain intermediates used in the manufacture of dyestuffs has long been recognized as a factor in the production of bladder tumours in workers in the industry. Among the various compounds used in the synthesis of dyestuffs, only 2-naphthylamine appears to have been shown experimentally to give rise to tumours located predominantly in the bladder. The first well established experimental demonstration of this carcinogenic

action of 2-naphthylamine was that of Hueper, Wiley & Wolfe (1938), who obtained papillomas and carcinomas of the bladder in thirteen out of sixteen female dogs which had received commercial 2-naphthylamine by subcutaneous injection and oral administration for a period of 20–26 months. Later, Bonser (1943) described the production of bladder tumours in dogs following prolonged oral administration of purified 2-naphthylamine. The fact that the tumours to which this amine gives rise in man and in experimental animals occur almost exclusively in the bladder has led to the suggestion that the carcinogenic agent is excreted in the urine and that it is

* Present address: Department of Microbiology, Western Reserve University, Cleveland, Ohio.

† Present address: Department of Biochemistry, St Thomas's Hospital Medical School, London, S.E. 1.