Carotenoid content of Patella depressa. A few samples of *P. depressa* were obtained from Plymouth at the same times as were *P. vulgata*. It has already been shown that the carotenoid constituents are the same in *P. depressa* and *P. vulgata* (Goodwin & Taha, 1950) and the present data (recorded in Table 4) indicate that the quantitative distribution of the pigment is also similar in the two species. This is specially marked in the case of the testes, the rather greater differences between the ovaries of the two species again being probably due to the different stages of spawning in the samples examined. One of the most interesting problems concerning limpet ovaries is the question of the production of brown and green ovaries. It has already been shown (Goodwin & Taha, 1950) that carotenoids play no major part in the production of green ovaries, a chromoprotein containing a non-carotenoid prosthetic group being the most important factor. The increased concentration of carotenoids in brown ovaries is unlikely to cause the change from green to brown; more probably it depends on the observation that the isolated green chromoprotein turns brown on addition of a few drops of dilute acetic acid. The brown and green colours may well be due, therefore, to variations in pH of the ovaries.

SUMMARY

1. The four carotenoids, β -carotene, echinone, cryptoxanthin and zeaxanthin, occur in the gonads of *Patella vulgata* in the ratios 5:2:2:2.

2. There are no significant deviations from this distribution in either the brown or green ovaries or in the testes.

3. A similar distribution of the same pigments is found in *Patella depressa*.

4. Although brown and green ovaries contain the same amounts of carotenoids, the brown have a higher concentration, indicating that brown gonads are generally smaller than the green.

5. The concentrations in the testes and ovaries are very similar, but the content of the testes is greater; this, again, merely reflects the weight relationship of ovaries and testes at the time of sampling.

6. Limpets collected at the same time from different regions of the British Isles contain comparable amounts of carotenoids.

Thanks are due to Prof. J. H. Orton, F.R.S., for supplying the limpets.

REFERENCES

Goodwin, T. W. (1949). *Biochem. J.* **45**, 472.

Goodwin, T. W. & Taha, M. M. (1950). *Biochem. J.* **47**, 244.

The Reaction of Oxidizing Agents with Wool

2. THE SORPTION OF HYDROGEN PEROXIDE

BY P. ALEXANDER, D. CARTER AND C. EARLAND
Research Department, Wolsey Ltd., Leicester, England

(Received 4 February 1950)

Hydrogen peroxide was shown by Harris & Smith (1936*a, b*, 1937*a, b*) to oxidize the cystine in wool partly to cysteic acid and partly to cystine disulphoxide. Rutherford & Harris (1938) found that the reaction was far from specific, and that many other groups besides the disulphide bond were oxidized, and some evidence for preferential attack on the free amino groups of the basic amino-acids was found.

Elöd & Zahn (1940) showed that silk, which contains no cystine, dissolved completely in 3% (w/v) hydrogen peroxide (pH 4) after 3 days at 60°. Wool (Elöd & Zahn, 1942) behaved likewise, and the solution obtained contained no high molecular weight fractions. Hydrogen peroxide is therefore undoubtedly capable of breaking peptide bonds, although in view of the stability of polyglycine to

this reagent it seems probable that the attack is preceded by the oxidation of a reactive side group. The rate of reaction of wool with hydrogen peroxide is slow and at room temperature no oxidation can be detected, except in very alkaline solutions for several hours. All the reactions discussed were carried out at elevated temperatures.

While studying bleaching it was found that there was a small but detectable decrease in the peroxide concentration immediately on immersing the wool in the solution, after which no further change could be detected for several hours. A similar effect was observed with nylon and silk but not with cotton. The peroxide, which very rapidly disappeared, was unambiguously shown to be physically held by the nitrogen-containing fibres and not to have reacted

chemically, since it was completely recovered by successive washings of the fibres with water.

Hydrogen peroxide forms addition compounds with both organic and inorganic substances (Machu, 1937), the crystalline compound with urea, $\text{NH}_2\text{CONH}_2 \cdot \text{H}_2\text{O}_2$, being the best known of these because of its industrial use as a concentrated stable source of peroxide. X-ray diffraction analysis of this compound (Lu, Hughes & Giguère, 1941) has revealed that the molecules are held together by hydrogen bonds.

All organic compounds capable of forming stable complexes with hydrogen peroxide contain either amino or imino groups (Krepelka & Buksa, 1937; Tanatar, 1901). The adducts with compounds containing carboxyl or hydroxyl groups, on the other hand, are very unstable in the few cases in which they are formed (Tanatar, 1901). Maass & Matheson (1929) found in a series of nitrogen bases that the amount of hydrogen peroxide combined increased with increasing basicity. It would appear that hydrogen bonding is not the only factor which plays a part in the formation of the peroxide complexes.

No addition compounds between proteins and hydrogen peroxide have been recorded except for the special case of the enzyme catalase.

EXPERIMENTAL

Materials

The H_2O_2 solutions were prepared by diluting a 90% sample of very high purity (obtained from LaPorte Ltd.). H_2SO_4 (0.01N) was added to stabilize the solutions and the pH was adjusted by the addition of NaHCO_3 or sodium silicate.

The fibres used were in the form of knitted fabric and were purified by Soxhlet extraction with ether, ethanol and water. The wool powder was prepared by oxidizing all the cystine with peracetic acid in a sample of wool which then became soluble in dilute NH_4OH . This solution was filtered and the wool powder precipitated by the addition of 0.1N-HCl (Alexander & Earland, 1950). The acetylated wool was prepared by the I.G. Farben process (Alexander & Whewell, 1946) and from acid-adsorption measurements it was concluded that 77% of the amino groups were blocked.

Measurement of sorption

The amount of H_2O_2 sorbed was determined by measuring the amount of oxidizing agent on the fabric and not from changes in concentration of the solution. This procedure gave more accurate results and also ensured that only true physical adsorption was measured.

Accurately weighed samples of textile material of approx. 2 g. dry weight were thoroughly wetted out in a solution of the same pH as the H_2O_2 . The fabric was then passed through squeeze rollers, put into 100 ml. of the H_2O_2 solution at 20° and thoroughly stirred for 5 min. The material was withdrawn, again passed through squeeze rollers and the weight of solution retained determined by weighing. The fabric was washed five times with 20 ml. of water and the H_2O_2 in the combined washings determined by titration

with acid KMnO_4 . More than 95% of the peroxide was thus removed and that remaining in the material was measured by placing the fabric in 0.5N- H_2SO_4 and titrating with KMnO_4 , taking care not to allow the latter to come into direct contact with the fabric. The strength of the original H_2O_2 solution was determined by titration after removal of the fabric. It was assumed that all the water carried over from the H_2O_2 solution contained H_2O_2 in the same concentration as in the latter, and the amount adsorbed was calculated by subtracting the amount in the water carried over from the total amount of peroxide in the washings. The amount of peroxide adsorbed may be somewhat greater than the value obtained in this way, since some of the water carried over may be physically held by the fibres and thus not function as solvent water. The magnitude of this effect is the subject of controversy (Hailwood & Horrobin, 1946), but the error introduced by neglecting it cannot be serious in these experiments.

With fast stirring the sorption was complete in less than 1 min., and identical results were obtained after 1, 2, 5 and 10 min. The sorption by wool powder was determined by shaking 1.5 g. of the latter with 10 ml. of peroxide solution and measuring the change in titre, a constant value being obtained after 30 sec.

RESULTS

From Table 1 it is seen that only nitrogen-containing fibres sorb hydrogen peroxide from aqueous solution. The adsorption isotherms for wool and silk with hydrogen peroxide at different pH values are shown in Figs. 1-3. The maximum sorption could in no case be determined since the flat part of the isotherm was not reached in the range of hydrogen peroxide which it was possible to study. At concentrations greater than 40% (v/v) the results became erratic and completely unreliable because of instability of the peroxide.

Table 1. *Sorption of hydrogen peroxide from 3% (w/v) aqueous solutions at pH 8*

Fibre	Sorption (mg. H_2O_2 /g. fibre)
Wool	9.8
Acetylated wool	5.5
Nylon	5.5
Silk	4.5
Cotton	0
Glass	0

The amount of sorption by wool is seen from Fig. 1 to be independent of the pH of the peroxide solution in the range pH 2.5-9, but to decrease at both lower and higher values. The sorption by both silk and nylon, on the other hand, is not decreased at low pH values and is the same at pH 6 as at pH 0 (see Table 2 and Fig. 2).

The sorption of hydrogen peroxide by wool was unchanged when the peroxide solution contained normal potassium chloride, both at pH 6 and 9, and it is concluded that electrolytes have no effect.

The powder obtained by precipitation with acid from a solution of wool, all the disulphide bonds of which have been oxidized yielding pairs of sulphonic acid groups, sorbs the same amount of peroxide as untreated wool fibres (see Fig. 3). Two conclusions

hydrogen peroxide combines with the amino-acid and imino groups in wool and silk. A similar parallelism between simple compounds and fibres is shown by sugars and cotton, neither of which combines with hydrogen peroxide.

Fig. 1. Sorption of H_2O_2 by wool at different pH values: $\bullet$ — $\bullet$, pH 0; $\circ$ — $\circ$, pH 1; $\otimes$ — $\otimes$, pH 2.5; $\odot$ — $\odot$, pH 6; ϕ — ϕ , pH 8; $\ominus$ — $\ominus$, pH 9; $\oplus$ — $\oplus$, pH 10.

Fig. 2. Sorption of H_2O_2 by silk at pH 6; *, solution of pH 0.

Fig. 3. Sorption of H_2O_2 by precipitated wool at pH 4.

can be drawn from this experiment. First, the hydrogen peroxide cannot be held by some purely physical capillary effect in the pores of the fibre. Secondly, the formation of sulphones or sulfoxides can play no part and the apparent sorption cannot arise from the reversible formation of such compounds. By analogy with the behaviour of organic nitrogen compounds it seems probable that the

Table 2. Sorption of hydrogen peroxide by nylon from 6% (w/v) aqueous solutions at different pH values

pH	Sorption (mg. H_2O_2 /g. nylon)
0	13.0
1	11.3
2	11.7
3	11.2
4	11.9
5	11.6
6	11.5

No accurate estimate of the maximum sorption of hydrogen peroxide can be made since the flat part of the isotherm cannot be reached experimentally, but inspection of the curve in Fig. 1 indicates a value of 75–100 mg. of hydrogen peroxide per g. of wool (i.e. 2–3 mol. of hydrogen peroxide per kg. of wool). In 1 kg. of wool there are approximately 0.9 mol. of free-amino groups from the basic amino-acids arginine and lysine and 9 mol. of peptide carbimino groups (Astbury, 1942). There are not sufficient amino groups, therefore, to combine with all the hydrogen peroxide and the carbimino groups must also play a part. Two observations indicate that the amino group, however, has the greater affinity for hydrogen peroxide; first, acetylation of wool during which an amino group is converted to a carbimino group leads to a greatly decreased sorption; secondly, silk which contains 12 mol. of carbimino groups but only 0.1 mol. of amino groups per kg. sorbs less peroxide than wool.

The decrease in sorption at low pH values observed in the case of wool might at first sight be explained by the ionization of the carbimino group,

which has been shown to decrease its hydrogen-bonding capacity (Theis, 1946). A similar ionization would, however, also take place in silk and nylon but their sorption of hydrogen peroxide is not decreased at low pH values and this explanation is therefore unlikely to be correct.

The sorption of peroxide by wool from a solution at pH 10 is markedly decreased (see Fig. 1), and this is probably due to the combination of wool with alkali which becomes appreciable only above pH 9. The amino group changes from NH_3^+ to NH_2 and the wool consequently becomes less electropositive. A similar effect was observed by Maass & Matheson (1929), who found that the sorption of peroxide by organic bases is increased with increasing basicity.

SUMMARY

1. Wool, silk and nylon, but not cotton, sorb hydrogen peroxide from aqueous solution and the concentration range 0–45 % has been studied. The sorption has not reached its limit at 45 %, but stronger solutions could not be studied.

2. With wool the sorption is independent of the pH of the peroxide solution in the range

pH 2.5–9, but decreases at both lower and higher pH values.

3. It is concluded that the hydrogen peroxide is held by both the amino and carbimino groups and that the sorption process resembles the formation of complexes with simple organic nitrogen-containing compounds such as urea.

The authors wish to thank Mr R. H. Reed for experimental assistance and the directors of Messrs Wolsey Ltd. for permission to publish this paper.

REFERENCES

- Alexander, P. & Earland, C. (1950). *Textile Res. J.* **20**, 296.
 Alexander, P. & Whewell, C. S. (1946). *B.I.O.S. Report* 1472, p. 91. London: H.M. Stationery Office.
 Astbury, W. T. (1942). *J. chem. Soc.* p. 337.
 Elöd, E. & Zahn, H. (1940). *Kolloidzshr.* **93**, 50.
 Elöd, E. & Zahn, H. (1942). *Melliand Textilber.* **23**, 313.
 Hailwood, A. J. & Horrobin, S. (1946). *Trans. Faraday Soc.* **42B**, 84.
 Harris, M. & Smith, A. L. (1936*a*). *Bur. Stand. J. Res., Wash.*, **16**, 301.
 Harris, M. & Smith, A. L. (1936*b*). *Bur. Stand. J. Res., Wash.*, **17**, 557.
 Harris, M. & Smith, A. L. (1937*a*). *Bur. Stand. J. Res., Wash.*, **18**, 623.
 Harris, M. & Smith, A. L. (1937*b*). *Bur. Stand. J. Res., Wash.*, **19**, 81.
 Krepelka, J. H. & Buksa, R. (1937). *Chem. Listy*, **31**, 447.
 Lu, C.-S., Hughes, E. W. & Giguère, P. A. (1941). *J. Amer. chem. Soc.* **63**, 1507.
 Maass, O. & Matheson, G. L. (1929). *J. Amer. chem. Soc.* **51**, 674.
 Machu, W. (1937). *Wasserstoffperoxyd und die Perverbindungen*. Vienna: Springer.
 Rutherford, H. A. & Harris, M. (1938). *Bur. Stand. J. Res., Wash.*, **20**, 559.
 Tanatar, S. (1901). *Z. anorg. Chem.* **28**, 225.
 Theis, E. R. (1946). *Trans. Faraday Soc.* **42B**, 244.

Acid-soluble Pigments of Molluscan Shells

5. IDENTITY OF SOME SUBSIDIARY FRACTIONS DERIVED FROM *PINCTADA VULGARIS*

By A. COMFORT

Department of Physiology, London Hospital Medical College, London, E. 1

(Received 28 March 1950)

Reference was made in a previous paper (Comfort, 1949) to the violet and blue subsidiary fractions which accompany uroporphyrin in the shell of the pearl oyster *Pinctada vulgaris*. A fuller examination of these substances, isolated by the methods already described, shows that they exhibit reactions characteristic of biladienes.

EXPERIMENTAL

The pigments were obtained as by-products of a large-scale extraction of uroporphyrin I from 5 kg. of broken shell, which yielded about 150 mg. of uroporphyrin, purified as its methyl ester. The blue and violet fractions (3 and 3*a*) from talc chromatograms were pooled, dried, rechromatographed on talc from N-hydrochloric acid, and the pigments eluted with acetone.

The main subsidiary pigment (3) is violet in acid solution, and is separated with difficulty from a sky-

blue fraction (3*a*). Both these fractions are green in alkaline solution, with a strong red fluorescence in ultraviolet light. The spectrophotometric curve of pigment 3 is given in Fig. 1. The alkaline spectrum is greatly intensified by the addition of methanolic zinc acetate, but without displacement of the bands. Pigment 3 is converted to a red material ($\lambda_{\max.} = 494 \text{ m}\mu$.) by concentrated nitric acid. Reduction by zinc dust in N-hydrochloric acid gives a yellow solution with strong blue fluorescence.

Pigment 3 cannot be extracted from aqueous solution at any pH by chloroform, and is very slightly extracted by ether at pH 7.5. Pigment 3*a* appears to be slightly more soluble in ether. Both pigments are soluble in acetone and in methanol.

A portion of pigment 3 was dried in a desiccator and esterified with a saturated solution of hydrogen chloride in methanol for 2 hr. at room temperature. Chloroform (1 ml.) was added, followed by water to