5. 2:4-Dinitrophenol acts like azide, but requires a much higher concentration for 50 % inhibition. Salicylaldoxime and sodium diethyl-dithiocarbamate also act like azide, but are even less effective than 2:4-dinitrophenol.

6. Monoiodoacetate (0.005 M) entirely inhibits acid and ethanol production, whereas fluoroacetate has no appreciable effect.

7. The facts of the acid secretion are well explained by a redox theory, in which it is considered that the H^+ ions are produced in an outer cell region by the ionization of the reduced form of a heavy metal

catalyst, receiving metabolic hydrogen atoms from a dehydrogenase (directly or at one or more removes) and being in turn oxidized within the cell where it receives H^+ ions and transfers hydrogen atoms to an acceptor.

8. This redox theory has an obvious application to secretion of acid by the gastric mucosa, the facts of acid secretion by yeast being now strikingly similar to acid secretion by the parietal cells. Other applications may also be entertained, such as acidification in the nephrons of the kidney or alkalinization by the pancreas.

REFERENCES

- Brady, T. G. (1948). *Biochem. J.* **42**, lxiii.
 Conway, E. J. (1949). *Irish J. med. Sci.* **288**, 787.
 Conway, E. J. & Brady, T. G. (1948). *Nature, Lond.*, **162**, 456.
 Conway, E. J. & Brady, T. G. (1950). *Biochem. J.* **47**, 360.
 Conway, E. J., Brady, T. G. & Carton, E. (1949). *1st Int. Congr. Biochem. Abstr.* p. 311.
 Conway, E. J. & Downey, M. (1950). *Biochem. J.* (in the Press).
 Conway, E. J. & Nolan, B. (1948). Unpublished observations.
 Conway, E. J. & O'Malley, E. (1948). *Biochem. J.* **42**, lxi.
 Conway, E. J. & O'Malley, E. (1949). *1st Int. Congr. Biochem. Abstr.* p. 515.
 Davies, R. E. (1948). *Biochem. J.* **42**, 609.
 Hanke, M. E. (1937). *Science*, **85**, 54.
 Spiegelman, S., Kamen, M. D. & Sussman, M. (1948). *Arch. Biochem.* **18**, 409.

The Use of Radioactive Isotopes in Immunological Investigations

3. FURTHER STUDIES ON THE FATE OF INJECTED ^{32}P -CONTAINING PROTEINS

By T. E. BANKS, G. E. FRANCIS, K. J. FRANKLIN AND A. WORMALL

Departments of Biochemistry, Physics and Physiology, Medical College of St Bartholomew's Hospital, London

(Received 17 February 1950)

In a previous paper (Banks, Bournsnel, Dewey, Francis, Tupper & Wormall, 1948) it was shown that the ^{32}P -containing protein antigens, lipovitellin, vitellin and artificially phosphorylated proteins, when injected intravenously into rabbits, rapidly disappear from the blood stream. With lipovitellin and vitellin, for example, only one-eighth of the injected protein could be detected in the blood 5 min. after the injection. Furthermore, in preliminary investigations on two rabbits injected with a suspension of vitellin it was found that much of the vitellin which had left the general circulation had been removed by the liver and lungs; e.g. about one-third of the injected vitellin was found in these two organs 1 min. after the injection. Vitellin is, however, a very insoluble protein, and it had to be injected as a suspension. It was considered possible that different results might be obtained with more soluble proteins, and we decided, therefore, to use the more soluble lipovitellin for a further study of this problem.

For an investigation of the immediate distribution of the injected protein it was necessary to kill the animal as soon as the injected material had become well distributed throughout the blood stream, i.e. 2-3 min. after the injection, and to do this we used the rapid coagulation technique recently introduced by Franklin & Amoroso (1948), namely the intravenous injection of Russell's viper venom, which causes clotting in 1-4 min. Subsequent immersion of the whole animal in saline containing formaldehyde fixes the protein antigen and prevents its diffusion from the blood or tissue. In the other experiments described here lipovitellin was injected intravenously, and the injected rabbits were killed, by the injection of magnesium sulphate or by stunning, after intervals ranging from 2 to 44 hr.

One object of these investigations was to study the distribution of protein antigens in the body at varying intervals after their injection into the blood stream, since information on this matter might throw light on the problem of the site of antibody

formation. Our experiments are also designed to study the possibility of obtaining a radioactive preparation which is specifically deposited and retained in certain tissues of the body.

EXPERIMENTAL

^{32}P -containing lipovitellin

Preparation A. This was prepared as described previously (Francis & Wormall, 1948) from the yolks of eggs laid by a hen during the period 4–11 days after the hen had received an intramuscular injection of neutralized sodium phosphate solution containing 400 μc . of ^{32}P . The final lipovitellin solution (in approx. 5% NaCl solution) contained 1.21 g. N/100 ml. and 0.169 g. P/100 ml., and these values correspond to an approx. 10% (w/v) solution of lipovitellin.

Preparation B. This preparation was made in the same way as preparation A; the final solution contained 0.185 g. P/100 ml.

Injections

'Venom' series. Rabbits nos. SRB43, SRB44, SRB46 and SRB48 were anaesthetized with nembutal and ether, and the lipovitellin solution (2 ml. of preparation A/kg. for the first three rabbits, and 1 ml./kg. for SRB48) was injected through a cannula into the external jugular vein. Russell's viper venom (Stypven; Burroughs Wellcome Ltd.) was injected 1–2 min. later, and after death, about 2 min. later, the rabbits were immersed in 'formol-NaCl' solution (1 vol. formalin plus 3 vol. 0.9% NaCl) and left for about 9 days. In another set of experiments, rabbits SRB32 and SRB33 were injected with 3.8 and 1.8 ml. of lipovitellin solution (preparation B)/kg., respectively, and were subsequently treated in the same way as were those injected with preparation A.

'Non-venom' series. Lipovitellin solution (1 ml. of preparation A/kg.) was injected into the marginal ear veins of seven rabbits. Two rabbits died about 2 hr. after the injection, two were killed 21 hr. and three 44 hr. after the injections. Three of the rabbits were killed by the intravenous injection of 2–5 ml. of a saturated solution of MgSO_4 , and two by stunning; no essential difference was observed between the radio-P contents of the organs of rabbits killed by MgSO_4 and the corresponding values for the rabbits killed by stunning.

Collection and treatment of blood and tissue samples for ^{32}P determinations

Blood. Blood clots were taken from the hepatic vein and the dorsal aorta of the venom-treated rabbits. In the case of the two rabbits (nos. 524 and 717), which died 2 hr. after the lipovitellin injections, blood was withdrawn from the heart, and with all the other rabbits blood was collected, immediately before the animal was killed, from the ear not used for the injection of lipovitellin. These blood samples (1.5–2.95 g. for the venom-treated rabbits and 2.5–10 ml. for the others) were ashed and treated as described for the tissue samples.

Tissues. The whole liver, kidneys, heart, lungs, spleen and gall bladder, and a sample of muscle from the abdominal wall were removed from each rabbit, and the tissue surface washed; it was then dried on filter paper, weighed,

dried at 100–150° and subsequently ashed in a porcelain basin at 400–800° in an electric muffle furnace. The ash was dissolved as completely as possible in 10.5N-HCl, and the solution or suspension was diluted with 4 vol. of water and boiled for 1 hr. to hydrolyse pyrophosphate. The product, or 20–25% of it in the case of the liver preparations, was treated with magnesia mixture, and the precipitate was collected by filtration on a sintered glass or silica Gooch crucible; it was then dried at 110° and a suitable weighed amount (sufficient to give not more than 20 mg./sq.cm. on the plate) made into a paste with ethanol containing a little collodion and spread evenly on brass 'sample plates'. The specimen was allowed to dry and ^{32}P determinations were then made.

In order to obtain a preliminary guide to the distribution of the radioactive material, thin slices were cut as uniformly as possible from the formalin-treated tissues of some of the rabbits killed by venom; these slices were placed directly under the window of the Geiger-Müller counter and radioactivity measurements were made. The results agreed fairly well with those obtained with the ashed samples, and this method of 'counting' is sometimes useful when an indication of the relative amounts of radioactive material in the tissues is required and the more tedious ashing process cannot be undertaken.

RESULTS

The distribution of lipovitellin in the rabbit about 2 min. after its intravenous injection

The first three rabbits received injections of 2 ml. of our stock radioactive lipovitellin solution (preparation A)/kg. of body weight, but since this amount was found to be toxic, the other animals injected with this preparation were given 1 ml./kg. The results obtained with four rabbits showed that a considerable fraction of the injected material was localized in the lungs, liver and heart (Table 1). Of the lipovitellin injected, 25–38% was found in the lungs and 10–15% in the liver. The amount of lipovitellin found in the hearts of these rabbits was quite large (11–26% of that injected), but perhaps little more than could be attributed to the large amount of blood 'fixed' in that organ under the conditions of these experiments.

It was not possible to determine the exact amount of lipovitellin in the blood of these animals at the time of death. If, however, it is assumed that the injected protein was uniformly distributed throughout the blood during the 2 or 3 min. interval between injection of the protein and the clotting of the blood, and if the blood volume of the rabbit is taken as 50 ml./kg. of body weight, the amount of radio P in the blood when the rabbits died would be about 68 μg ./ml. for nos. SRB43, SRB44 and SRB46, and 34 μg ./ml. for no. SRB48. The values found for the blood clots removed from veins or arteries were 181, 140 and 18.3 μg . of radio P/g. for rabbits SRB43, SRB44 and SRB48, respectively; since two of these values are much greater than the 'theoretical'

Table 1. *Radio-P content of the tissues of rabbits killed with viper venom about 2 min. after the injection of ^{32}P -containing lipovitellin (preparation A)*

(Rabbits nos. SRB43, SRB44 and SRB46 (1.9, 2.0 and 2.45 kg., respectively) received intravenous injections of 2.0 ml. of lipovitellin solution/kg., and rabbit SRB48 (2.75 kg.) received 1.0 ml./kg.)

Rabbit no.	...	...	SRB43	SRB44	SRB46	SRB48
Radio P injected ($\mu\text{g.}$)	...	...	6450	6780	8300	4660
Radio-P* content of organs						
Liver	($\mu\text{g./g.}$)		11.6	9.8	17.7	6.8
	Total ($\mu\text{g.}$)		640	737	899	697
Lungs	($\mu\text{g./g.}$)		210	172	222	158
	Total ($\mu\text{g.}$)		1674	1770	2096	1770
Heart	($\mu\text{g./g.}$)		150	165	—	47.5
	Total ($\mu\text{g.}$)		1218	1777	—	540
Spleen	($\mu\text{g./g.}$)		<16	33	—	<14
	Total ($\mu\text{g.}$)		<10	20	—	<10
Kidneys	($\mu\text{g./g.}$)		3.2	5.5	—	3.0
	Total ($\mu\text{g.}$)		34.7	79.3	—	52.4
Gall bladder (plus contents)	Total ($\mu\text{g.}$)		21.9	<10	—	18.7
Abdominal wall muscle (sample)	($\mu\text{g./g.}$)		<1.4	<1	—	<0.8

* The term 'radio P' is used throughout this paper to denote amounts of radioactive P of the same specific activity as that of the P in the lipovitellin injected into the rabbit in that particular experiment.

Table 2. *Radio-P content of the tissues of rabbits killed with viper-venom about 2 min. after the injection of ^{32}P -containing lipovitellin (preparation B)*

(Rabbit SRB32 received 3.8 ml. lipovitellin solution/kg. and rabbit SRB33 1.8 ml./kg.)

Rabbit no.	...	...	...	SRB33	SRB32
Radio P injected ($\mu\text{g.}$)	...	...	...	7,400	14,800
Radio-P content of organs					
Liver	($\mu\text{g./g.}$)			14.7	22
	Total ($\mu\text{g.}$)			1,181	2,020
Lungs (part)	($\mu\text{g./g.}$)			161.3	400
Heart	($\mu\text{g./g.}$)			18.6	114.5
	Total ($\mu\text{g.}$)			133	694
Spleen	($\mu\text{g./g.}$)			24.5	37.4
	Total ($\mu\text{g.}$)			20	22
Kidney (one)	Cortex ($\mu\text{g./g.}$)			13.3	
	Medulla ($\mu\text{g./g.}$)			9.4	19.5
Gall bladder (plus contents)	Total ($\mu\text{g.}$)			39	15.3
Abdominal wall muscle	($\mu\text{g./g.}$)			1.1	4

values it is concluded that they do not give a true measure of the amount of lipovitellin in the blood present in the organs and tissues examined. It is probable that the clots had removed practically all the lipovitellin from the plasma, and an accurate value for the radio-P content of the blood could only have been obtained by making measurements on the clot plus an appropriate volume of plasma. This was not possible under the conditions of our experiments.

The amount of radio P present in the lungs of the rabbits of this group was very much more than could be attributed to the blood present in the tissue, but the amounts in the liver (6.8–17.7 $\mu\text{g./g.}$) and kidneys (3.0–5.5 $\mu\text{g./g.}$) are probably explained by the lipovitellin in the blood present in these organs.

Rabbit no. 43 was found to be pregnant, and the foetuses were examined for radio P. The amount present was extremely small, being <0.75 $\mu\text{g./g.}$, and the total for the whole of the eight foetuses was <100 $\mu\text{g.}$ Similarly, there was little lipovitellin in the urine of these rabbits, the mixed urines of nos. SRB43 and SRB46 containing <2.5 $\mu\text{g.}$ of radio P/ml.

The results obtained with the rabbits injected with lipovitellin preparation B (Table 2) fully confirmed those recorded in Table 1, in showing that the lungs contained much more lipovitellin per g. of tissue than did the liver and other organs examined. In these experiments, however, no attempt was made to determine the radio-P content of the whole

of all the organs examined, and in certain respects the experiments differed from those described above; e.g. rabbit SRB 32 received a much larger injection of lipovitellin than did SRB 33 and the rabbits injected with preparation A. The distribution of the injected lipovitellin was, however, generally the same in all the venom-treated rabbits.

The distribution of radio P in the body 2, 21 and 44 hr. after intravenous injection of ³²P-containing lipovitellin

In these experiments the rabbits were killed either by the intravenous injection of magnesium sulphate or by stunning, and here it was possible to take blood samples just before the animals were killed. Calculations can thus be made to determine approximately what fraction of the radio P found in any tissue can be attributed to that present in the blood in the tissue.

During the first 21 hr. after the injection of the radioactive lipovitellin there was a large amount of radio P in the lungs and liver (Table 3). The amount of radio P, and presumably of lipovitellin, in the lungs of the two '2 hr.' rabbits was extremely high, representing 58 and 74 % of that injected, and it is probable that deposition of this large amount of lipovitellin in the lung capillaries was the main factor responsible for the death of these two rabbits about 2 hr. after the injection. Smaller but significant amounts of radio P were found in the kidneys and spleen, the amount present in these organs being usually considerably greater per g. of tissue than the amount per ml. of blood.

The lungs and livers of the rabbits killed 44 hr. after the injection similarly contained large amounts of radio P. The amount in the liver was considerably greater, and the amount in the lungs less, than the corresponding values for the rabbits killed 2 hr. after

Table 3. *Radio-P content of the tissues of rabbits 2 or 21 hr. after the injection of ³²P-containing lipovitellin*

(Rabbits nos. 524, 717, 532 and 698 (1.9, 3.3, 2.2 and 2.2 kg., respectively) each received an intravenous injection of 1.0 ml. lipovitellin solution/kg. Rabbits 524 and 717 died 2 hr. after the injection; rabbits 532 and 698 were killed 21 hr. after the injections, the former by intravenous injection of 3-4 ml. of a saturated solution of MgSO₄ and the latter by stunning.)

Rabbit no. ...		2 hr. after injection		21 hr. after injection	
		524	717	532	698
Radio P injected (μg.) ...		3210	5600	3725	3725
Radio-P content of organs					
Liver	(μg./g.)	8.0	6.3	16.1	10.2
	Total (μg.)	487	759	1340	804
Lungs	(μg./g.)	228	320	58.3	64.8
	Total (μg.)	1850	4160	629	700
Spleen	(μg./g.)	41.8	<12.5	32.5	37.5
	Total (μg.)	27.6	<10	24.4	26.6
Kidneys	(μg./g.)	8.2	< 0.7	10.5	8.6
	Total (μg.)	99	<14	158	123
Gall bladder (plus contents)	Total (μg.)	<11	<10	20.7	<12
Blood	(μg./ml.)	<8	11.6	About 2	2.2

Table 4. *Radio-P content of some organs of rabbits 44 hr. after the injection of ³²P-containing lipovitellin*

(Rabbits nos. 534, 740 and 531 (1.65, 3.5 and 2.1 kg. respectively) each received an intravenous injection of 1 ml. of lipovitellin solution/kg. Rabbits 534 and 740 were killed by the intravenous injection of a saturated solution of MgSO₄ (2 and 5 ml. respectively) and no. 531 by stunning.)

Rabbit no.	...	534		740		531	
Radio P injected (μg.)	...	2796		5930		3565	
		Radio-P content of organs					
		(μg.)	(μg./g.)	(μg.)	(μg./g.)	(μg.)	(μg./g.)
Liver		550	10.8	1570	12.0	773	10.1
Lungs		292	40.2	416	32.5	406	37
Spleen		26	43.3	57	75	16*	26*
Kidneys		72	7.4	185	8.8	87	6.2
Gall bladder (plus contents)		<10	—	<11	—	<11	—
Blood		—	1.8†	—	1.85†	—	<1.1†

* Approximate values.

† μg./ml.

the injection. The spleen and kidneys of these '44 hr.' rabbits contained considerably more radio P per g. of tissue than did the blood.

The results of these experiments, including those with most of the rabbits killed by venom, are summarized in Table 5, where the amounts of ^{32}P found

A similar deposition of zinc in the lungs of rabbits after the intravenous injection of a ^{65}Zn -containing zinc diphenylthiocarbazon (dithizone) complex (Banks, Tupper & Wormal, 1950) might be explained in the same way, though the material injected in this case was quite unlike lipovitellin;

Table 5. *Amount of lipovitellin (calculated from the ^{32}P content and expressed as a percentage of that injected) in the organs of rabbits at intervals after the injection of lipovitellin containing ^{32}P*

(Experimental details are given in Tables 1-4.)

Time after injection	Rabbit no.	How killed	Percentage of the injected lipovitellin* found in					
			Liver	Lungs	Heart	Spleen	Kidneys	Gall bladder
2 min.	SRB43	Venom	9.9	26	19	<0.2	0.5	0.3
	SRB44	Venom	10.9	26	26	0.3	1.2	<0.15
	SRB46	Venom	10.8	25	—	—	—	—
	SRB48	Venom	15.0	38	11.6	<0.2	1.1	0.4
	SRB32	Venom	13.6	—	4.7	0.15	—	0.1
	SRB33	Venom	16.0	—	1.8	0.27	—	0.5
2 hr.	524	Died	15.2	58	—	0.9	3.1	<0.4
	717	Died	13.5	74	—	<0.2	<0.3	<0.2
21 hr.	532	MgSO ₄	36	17	—	0.7	4.3	0.6
	698	Stunned	22	19	—	0.7	3.3	<0.4
44 hr.	534	MgSO ₄	20	10.4	—	0.9	2.6	<0.4
	740	MgSO ₄	26	7.0	—	1.0	3.1	<0.2
	531	Stunned	22	11.4	—	0.4	2.4	<0.3

* Rabbits SRB43, SRB44 and SRB46 received an injection of 3.38 mg., SRB32 7.03 mg., SRB33 3.33 mg., and the other rabbits 1.69 mg. of lipovitellin P/kg.

in the various organs are expressed as percentages of the amounts injected. Although a general comparison of the results for the different groups of rabbits may be justified, it must be remembered that all the experiments were not made under exactly the same conditions. For example, the '2 min.' group of rabbits had rather different treatment from that given to the others, for three of the rabbits received twice as much lipovitellin as did the others. Also, for reasons given earlier in this paper, the method used to kill the rabbits in this group differed from that used for the other groups, and so did the site of injection and the time interval between the death of the animal and the removal of the organs.

DISCUSSION

Our results show that intravenously injected lipovitellin is rapidly taken up by the lungs and liver, and it is rather surprising that as much as one-quarter of the injected protein was found in the lungs of the injected rabbits 2 min. after the injection. This rapid deposition of injected protein in the lungs may possibly be related to the insolubility of lipovitellin in solutions containing less than 3-4 % of NaCl; in other words, the lipovitellin may have been precipitated almost immediately it entered the blood stream, and subsequently 'filtered off' when the blood passed through the lung capillaries.

the organic zinc compound was dissolved in triacetin and this solution made into an emulsion with normal rabbit serum before it was injected. However, until further information is available, it would be unwise to attempt to decide whether the deposition of the injected lipovitellin (or zinc dithizone complex) in the lungs is due to a mechanical filtering action of the lung capillaries or to some specific absorptive capacity of lung tissue for the injected compound.

Intravenously injected colloidal particles are not always taken up by the lungs. For example, Jones, Wrobel & Lyons (1944), using ^{32}P as tracer, found that anhydrous colloidal chromic phosphate is rapidly taken up by the liver, 59-81 % of the injected material being found in this organ 5-23 days after the injection; small amounts were found in the spleen (4.3-14.7 %) and the lungs (0.6-6.6 %). Similar localization in the reticuloendothelial cells of the liver and spleen has been observed with radioactive colloids containing yttrium (^{89}Y), zirconium (^{40}Zr) and columbium (^{41}Cb) (Dobson, Gofman, Jones, Kelly & Walker, 1948). In our experiments, however, the amount of injected compound localized in the liver shortly after the injection was relatively small (10-15 %), and although about twice as much was present in the livers of the rabbits killed 44 hr. later, the values were much lower than those obtained by the American workers in their experiments with inorganic colloids. Furthermore, the

amount of lipovitellin found in the spleen of our injected rabbits during the 44 hr. after the injection was never greater than 1% of the amount injected, and, as mentioned above, the lungs took up a large part of the lipovitellin which is removed from the blood shortly after the injection. It should be remembered, however, that the compounds we have injected, lipovitellin in the experiments reported here and a zinc dithizone complex in others (Banks *et al.* 1950), are quite different, chemically and physically, from the inorganic colloids used by Jones *et al.* (1944), Dobson, Kelly, Jones & Gofman (1947) and Dobson *et al.* (1948). It may be significant, in connexion with the similarities between the early fate of injected zinc dithizone complex and lipovitellin, that the former is fat soluble and the latter is a lipoprotein and contains about 25% of phospholipin.

In tracer experiments with injected compounds containing a radioactive isotope, radioactivity determinations merely estimate the amount of the isotopic label present. In the experiments described here, however, the conditions were such that one is probably justified in believing that the ^{32}P determinations give an assessment of the tissue distribution of the injected lipovitellin. The time interval before the rabbits were killed was short, and the ^{32}P -containing lipovitellin, made biologically, had a label which was relatively firmly attached. Any metabolic breakdown of the lipoprotein would liberate ^{32}P -containing compounds which might be transported elsewhere, but it seems probable that much of the lipovitellin (and possibly all the vitellin portion of this lipoprotein) must remain, in its original or some very closely related state, in the tissues where the specific antilipovitellin antibodies are formed. Furthermore, it is most unlikely that 'physical exchange' will occur between the radioactive phosphate groups of either the phospholipin or the vitellin and those of other phosphorus compounds in the body, for such an 'exchange' does not readily occur (cf. the review by Hevesy, 1948).

However, the possibility of exchange of phosphatide between the radioactive lipovitellin and the ordinary blood and tissue lipids should not be overlooked, but as yet we have no information on this point.

Although we have found that intravenously injected lipovitellin is at first localized mainly in the lungs, it does not necessarily follow that all similarly injected antigens are deposited in that tissue. Indeed it has been shown that the arsenic of an injected arsenic-containing azo-protein is mainly localized in the liver and bone marrow (Haurowitz & Breinl, 1932), and that injected ^{32}P -containing tobacco-mosaic virus is largely deposited in the liver (Libby & Madison, 1947). We hope to continue our studies on this problem of the distribution of injected antigens with the aid of iodinated and mustard gas treated proteins labelled with ^{131}I and ^{35}S respectively.

SUMMARY

1. The fate of ^{32}P -containing lipovitellin after its intravenous injection into rabbits was studied by determinations of the distribution of ^{32}P in the tissue at intervals after the injection.

2. A considerable part, at least one-quarter, of the injected lipoprotein was present in the lungs, and 10–15% in the liver, 2 min. after the injection. Over half the injected lipovitellin was present in the lungs of two rabbits which died 2 hr. after the injection.

3. The results for the rabbits killed 21 and 44 hr. after the injections suggested that after the initial deposition of lipovitellin in the lungs, with less in the liver, there was a gradual diminution in the amount held in the lungs and an increase in that deposited in the liver.

We are indebted to the Medical Research Council and the Central Research Fund of the University of London for grants (to A. W.) which have defrayed part of the cost of these investigations. We are also grateful to the Medical Research Council Tracer Elements Sub-committee for supplies of ^{32}P .

REFERENCES

- Banks, T. E., Boursnell, J. C., Dewey, H. M., Francis, G. E., Tupper, R. & Wormall, A. (1948). *Biochem. J.* **43**, 518.
 Banks, T. E., Tupper, R. & Wormall, A. (1950). *Biochem. J.* (in the Press).
 Dobson, E., Gofman, J., Jones, H., Kelly, L. & Walker, L. (1948). *Proc. 4th Int. Cancer Res. Congr.* Cited by Lawrence, J. H. & Hamilton, J. G. (1948).
 Dobson, E., Kelly, L., Jones, H. & Gofman, J. (1947). Communication to the 17th Int. Physiol. Congr. Oxford.
 Francis, G. E. & Wormall, A. (1948). *Biochem. J.* **42**, 469.
 Franklin, K. J. & Amoroso, E. C. (1948). Oral communication to the Physiological Society, 20 March. Unpublished.
 Haurowitz, F. & Breinl, F. (1932). *Hoppe-Seyl. Z.* **205**, 259.
 Hevesy, G. (1948). *Radioactive Indicators*, p. 85. New York: Interscience Publishers.
 Jones, H. B., Wrobel, C. J. & Lyons, W. R. (1944). *J. clin. Invest.* **23**, 783.
 Lawrence, J. H. & Hamilton, J. G. (1948). *Advances in Biological and Medical Physics*, 1. New York: Academic Press.
 Libby, R. L. & Madison, C. R. (1947). *J. Immunol.* **55**, 15.