$C_{14}H_{14}O_6$ requires: C, 64.1; H, 5.4%; mol. wt. 262.) The substance is almost insoluble in water, but dissolves to a yellow solution in hot NaOH, being recovered unchanged on acidification. It is insoluble in aqueous $NaHCO_3$. It is readily soluble in $CHCl_3$ from which it is precipitated in fine crystals by ether or light petroleum. Its ethanolic solution gives no marked colour with ethanolic $FeCl_3$ and no precipitate with Brady's reagent. It gives a clear orange-brown solution in cold conc. H_2SO_4 . It resists acetylation with sodium acetate and acetic anhydride, and on treatment with ethereal diazomethane a yellow oil, which could not be obtained crystalline, was formed. Similarly, ozonolysis in $CHCl_3$ afforded an uncrystallizable red gum. Catalytic reduction of its ethyl acetate solution (0.1 g. in 30 ml.) with Adams's PtO_2 catalyst (0.02 g.) resulted in the uptake in 3.5 hr. of 23 ml. H_2 . ($C_{14}H_{14}O_6$ absorbing 6H requires 25.7 ml. H_2 .) The isolated reduction product was an oil (0.12 g.). Similarly,

catalytic reduction of the ethanolic solution of the colourless ozonolysis product, $C_{14}H_{16}O_4$, with $PdCl_2$ -norite resulted in the slow uptake of almost 2H in 27 hr. and the formation of a stiff oil.

SUMMARY

An account is given of a number of derivatives and degradation products of fuscine, an antibacterial metabolite of the mould *Oidiodendron fuscum* Robak.

One of us, S.E.M., wishes to acknowledge with thanks receipt of a grant from The Therapeutic Research Corporation of Great Britain Ltd. We also desire to thank Mr P. Chaplen of this Department for carrying out the majority of the methoxyl determinations. All other analyses are by Drs Weiler and Strauss of Oxford.

REFERENCES

- Kinney, C. R. & Langlois, D. P. (1931). *J. Amer. chem. Soc.* **53**, 2189.
Michael, S. E. (1948). *Biochem. J.* **43**, 528.
Slotta, K. H. & Müller, J. (1936). *Hoppe-Seyl. Z.* **238**, 14.

Studies in Immunochemistry

9. THE OXIDATION OF THE HUMAN BLOOD-GROUP A SUBSTANCE WITH THE PERIODATE ION

By D. AMINOFF AND W. T. J. MORGAN

Lister Institute of Preventive Medicine, London, S.W. 1

(Received 14 June 1950)

Apart from the methylation studies of Bray, Henry & Stacey (1946) few structural investigations have been carried out on the blood-group A substance. These workers isolated and identified 2:3:4-trimethyl- α -methyl-L-fucoside and 3:4:6-trimethyl- α -methylglucosaminide hydrochloride in the esterified products from acid hydrolysis of a fully methylated A substance derived from pepsin, thus indicating that both fucose and *N*-acetylglucosamine are present as non-reducing end groups in the alkali-stable residues examined. A methylated derivative of D-mannose was also identified, but according to other investigators (Bendich, Kabat & Bezer, 1947; Partridge, 1948*b*; Aminoff, Morgan & Watkins, 1950) mannose is not present in the A substance. The behaviour of the A substance during mild hydrolysis with alkali (Morgan, 1946) and with acid (Kabat, Baer, Bezer & Knaub, 1948; Baer, Dische & Kabat, 1948; Aminoff, Morgan & Watkins, 1948, 1950) has provided additional information regarding the *N*-acetylhexosamine and fucose components and the roles they play in the different serological activities associated with the blood-group substances.

Oxidation of the A substance with periodate offers an independent and more specific approach to the determination of the structure of this important biological material. Periodic acid is a selective oxidizing agent which breaks the C—C linkage in substances containing either two adjacent unsubstituted hydroxyl groups or an unsubstituted hydroxyl group adjacent to a primary or secondary amino group. The products of oxidation are aldehydes, acids and, with hydroxyamino derivatives, ammonia. The reactions, under suitable conditions, are practically quantitative. The application of oxidation with periodate to structural studies in the field of carbohydrate chemistry has been reviewed by Owen (1943), Jackson (1944), and Jones (1946), and further discussion here is unnecessary.

The investigation of the oxidation of the A substance with periodate was approached along the following lines: (1) studies on the lability of the serological properties during oxidation; (2) determination of the amount of periodate reduced and the rate of reduction by the A substance under different conditions; (3) qualitative and quantitative analyses of

the material after oxidation to determine which of the constituent molecules are involved; and (4) analysis of the simple products liberated during the oxidation. A preliminary communication on this subject has already appeared (Aminoff & Morgan, 1949).

EXPERIMENTAL

The preparations and properties of the A substance used in the oxidation studies are described by Aminoff *et al.* (1950). Most of the oxidations were carried out at room temperature, and the periodate content of the reaction mixture was determined by titrating, with standard arsenite, the I_2 liberated on addition of KI (Jackson, 1944).

Rate of inactivation of A substance at different pH values

The A substance was oxidized with HIO_4 at room temperature at pH values between 5 and 10 using acetate, phosphate, bicarbonate and borate buffers. The initial concentration of A substance was 0.1%, of periodate 0.005M and of

Fig. 1. The effect of oxidation of the A substance with periodate at different pH values as determined by the capacity of the A substance to inhibit the iso-agglutination of A cells by anti-A serum. (A) pH 5, acetate; (B) pH 6, acetate; (C) periodic acid alone; (D) pH 8, phosphate; (E) pH 8, borate; (F) pH 9, (a) borate and (b) bicarbonate (G) pH 10, borate.

buffer 0.075–0.1M. The immediate and complete reduction of the excess periodate in 0.5 ml. quantities of the reaction solution, withdrawn at pre-arranged intervals, was accomplished by addition of 0.2 ml. of 20% (v/v) glycerol. The resulting solutions were then diluted with 0.85% NaCl to make a final dilution of 1:5000 of the A substance originally present, and the capacity of the material to inhibit iso-agglutination was determined by the standard method (cf. Morgan & King, 1943). The property of inhibiting the agglutination of A red cells by anti-A serum is rapidly lost, and when this specific activity, expressed as a percentage of the original value, is plotted against time a series of curves is obtained (Fig. 1). It is seen that with one exception the rate

of inactivation falls with rise in pH. The system containing free periodic acid, curve C, in the absence of buffer, behaves as though its pH were between 6 and 7, which is surprising in view of the fact that periodic acid has a pH of 2.5 under the conditions used.

Rate of reduction of periodate by A substance

Reaction at pH 5. A 0.1% solution of A substance was made up to contain 0.005M- HIO_4 and 0.1M-acetate buffer at pH 5.0. At appropriate intervals 2 ml. portions of the solution were withdrawn and added to conical flasks which contained 0.1N-sodium arsenite (0.5 ml.) and 9% (w/v) $NaHCO_3$ (0.5 ml.). The walls of the flask were washed down with 1 ml. of water and 20% (w/v) KI (0.1 ml.) was immediately added. After standing at room temperature for 15 min. the excess arsenite, not oxidized by the periodate, was titrated with 0.01N- I_2 solution using 0.5 ml. of 1% starch solution as indicator.

At the same time as each specimen was removed for titration, 0.4 ml. of the reaction mixture was also withdrawn and added to 1.6 ml. of glycerol-NaCl solution (0.2 ml. 20% (v/v) glycerol with 1.4 ml. 0.85% NaCl) to give a 1:5000 dilution of the A substance originally present, for the determination of serological activity. The solutions were kept at 0° until the experiment was completed and then assayed for the capacity of the material to inhibit (a) the agglutination of A cells, and (b) the haemolysis of sheep cells by anti-A serum.

Owing to the rapidity of reduction of periodate during the initial stages of the reaction, the periodate content at zero time was determined on a solution which contained all the reagents without A substance. This solution was also used to check the pH of the reaction mixture. Fig. 2, obtained by

Fig. 2. The oxidation of the A substance at pH 5. Rate of reduction of periodate by the A substance (A). The resulting inactivation as determined by the capacity of the material to inhibit the haemolysis of sheep cells (B), and the agglutination of A cells (C), by anti-A serum.

plotting the periodate reduced against the time of oxidation, shows that the reaction takes place in two distinct phases. An initial rapid reaction which is essentially complete after 4 hr. and a second phase in which the reduction still continues, but at a much reduced rate. The second phase is not due to instability of the periodate or to the oxidation of acetate present, since the control solution showed no fall in periodate content over a period of 44 hr. A correction for the over-oxidation can be obtained by simple extrapolation to zero time. The analytical results reveal that 1 mg. of A substance reduces 0.235×10^{-5} g.mol. of HIO_4 and hence 427 g. of A substance reduce 1 g.mol. of HIO_4 . A similar value, 370 g. of A substance/1 g.mol. of HIO_4 , was obtained with

preparation III (Aminoff *et al.* 1950). The power of the A substance to inhibit iso-agglutination falls very rapidly and reaches a value of not more than 1% of the original activity in 3 hr. The capacity of the material to inhibit anti-A sheep-cell haemolysins falls less rapidly and oxidation for 22 hr. is required to bring about inactivation to the same extent.

Reaction at pH 7.3. The reaction was carried out under the same conditions as described for the oxidation at pH 5.0. The HIO_4 concentration was, however, doubled as a preliminary experiment indicated that a greater consumption of periodate occurred at pH 7.3. Owing to the low solubility of NaIO_4 in the reaction mixture an opalescence developed and frequent shaking was necessary. A 0.1% solution of the A substance was prepared in a solution of 0.01M periodate and 0.04N- NaHCO_3 at pH 7.3. At intervals 1 ml. samples were withdrawn and titrated against standard I_2 as described above.

Fig. 3. The oxidation of the A substance at pH 7.3. Rate of reduction of periodate by the A substance (A). The resulting inactivation as determined by the capacity of the A substance to inhibit the haemolysis of sheep cells (B), and the agglutination of A cells (C), by an anti-A serum.

Samples (0.4 ml.) were also withdrawn for the serological titrations. The oxidation curve, Fig. 3, shows a more extensive and continuous reduction of periodate than was found for the oxidation at pH 5. There was only a slight indication of the existence of a point of inflexion and this occurred after about 2 hr. The examination of a sample of the reaction mixture at 23 hr. showed the complete reduction of all the periodate present, whereas the titration results with the control solution indicated that periodate was perfectly stable under these conditions. The fall in the serological activities is not as rapid as at pH 5.

Rate of oxidation in the presence of borate. In the presence of boric acid at pH 2.4 the oxidation of A substance was essentially complete in 3 hr. and 1 g. mol. of periodate was reduced by 294 g. of A substance (preparation III). The oxidation was also carried out in NaHCO_3 solution at pH 7.5 and with NaHCO_3 -borate mixture at the same pH. The course of oxidation was identical in these two buffer systems and was not complete after 72 hr.

Re-oxidation at pH 5.0 and 7.5 of A substance oxidized at pH 5. The solution of A substance obtained after oxidation at pH 5 was dialysed in a thoroughly washed cellophan tube. A preliminary experiment had shown that the material obtained after drying the oxidized product from the frozen state was largely insoluble in water. To determine the concentration of the main bulk of the solution only a portion of

the oxidized material was therefore taken to dryness. The concentration of the solution was found to be 0.476% with respect to the oxidized A substance. The recovery was 89%, no allowance being made for the weight of diffusible formic acid and formaldehyde, which together would have been of the order of 4% (see below).

Re-treatment of this solution with more periodate at pH 5.0 caused no further oxidation to occur (Fig. 6, curve C). At pH 7.5, however, there was a steady reduction of the periodate (Fig. 6, curve D). In one particular experiment, after 21 hr. at pH 5, 1 mg. of the oxidized A substance had reduced only 0.05×10^{-5} g.mol. of HIO_4 , whereas at pH 7.3 the corresponding value was 0.895×10^{-5} g.mol. and after 45 hr. 1.25×10^{-5} g.mol. had been reduced.

Re-oxidation at pH 5 and 8 of A substance oxidized at pH 7.5. The product from A substance after oxidation (48 hr., pH 7.5) and dialysis was mixed with 0.008M- HIO_4 in (a) 0.25M-acetate buffer pH 5, and (b) 0.04N- NaHCO_3 , pH 8. The course of oxidation was followed in the usual manner. Since the concentration of the oxidized A substance was not exactly known in this experiment, the results are expressed in terms of molarity of the solution with respect to the periodate.

The curves for the re-oxidation at pH 5 and 8 of material already oxidized at pH 7.5 are very similar (Fig. 4). The reaction in each instance approaches completion in about 24 hr., exclusive of the original 48 hr. oxidation at pH 7.5.

Fig. 4. Comparison of the rate of reduction of periodate by the A substance at (A) pH 5 and (B) pH 8 after initial oxidation with periodate at pH 7.5.

Fig. 5. Comparison of the rate of reduction of periodate by the A substance at (A) pH 5 and (B) pH 8 after an initial treatment with 0.5N- NaHCO_3 .

The effect of initial treatment with NaHCO_3 on the course of oxidation. In view of the known alkali-lability of the A substance (Morgan & King, 1943; Morgan, 1946) an experiment was carried out with undegraded A substance which had been exposed to 0.5N- NaHCO_3 for 24 hr. at about 16° to determine whether the more extensive oxidation at pH 7.5 was due to the dissociating effect of the NaHCO_3 . The results (Fig. 5) indicate that initial treatment with bicarbonate in no way influences the course of the subsequent oxidation at pH 5 (curve A) and pH 8 (curve B) (cf. Figs. 2A and 3A).

In order to determine whether the material oxidized at pH 5.0 is more labile to alkali than the unoxidized A substance some of the material oxidized at pH 5.0 was treated with NaHCO_3 at pH 9.4 for 24 hr. Subsequently, half the solution was adjusted to pH 5 with acetic acid and the HIO_4 added, whilst the addition of HIO_4 to the remainder of the alkali-treated material resulted in a reaction mixture with a

Fig. 6. Course of re-oxidation with periodate (at pH 5.0 or 7.5) of oxidized A substance (preparation III) with and without intermediate treatment with NaHCO_3 (pH 9.4; 24 hr.). A, re-oxidation at pH 5 after treatment with NaHCO_3 at pH 9.4; B, re-oxidation at pH 7.5 after treatment with NaHCO_3 at pH 9.4; C, re-oxidation at pH 5 without the initial treatment with NaHCO_3 at pH 9.4; D, re-oxidation at pH 7.5 without the initial treatment with NaHCO_3 at pH 9.4.

pH value of 7.5. The course of the re-oxidation at pH 5 and 7.5 of (a) material oxidized at pH 5.0 and then treated with NaHCO_3 for 24 hr. (at pH 9.4) and (b) some of the material oxidized at pH 5.0 without treatment with the NaHCO_3 , was followed in the usual manner and the results are shown in Fig. 6. The results indicate that the more extensive oxidation at pH 7.5 is not due to the greater alkali lability of the primary oxidation products of the A substance.

Qualitative and quantitative examination of oxidized A substance

Unless otherwise stated, the tests were carried out on a 0.476% solution of the product from A substance (preparation III) oxidized at pH 5.0, 24 hr., which had been thoroughly dialysed.

Qualitative tests. The oxidized material readily restored the colour to SO_2 -decolorized fuchsin (Schiff's reagent). The unoxidized A substance did not restore the colour. Oxidation, therefore, gives rise to the formation of an indiffusible aldehyde. A Molisch test with the oxidized A substance yields an intense bluish coloured boundary layer, with no trace of pink coloration; whereas, tested under the same conditions, unoxidized A substance gives a bright-red interfacial layer, and A substance previously added to the oxidized material imparts a reddish tint to the blue coloration. It may be concluded, therefore, that galactose and fucose, both of which readily give a positive Molisch reaction, are not present in the oxidized A substance.

Chromatographic analysis of the products of acid hydrolysis of the oxidized A substance. Oxidized and unoxidized A substance (5 mg.) were hydrolysed for 16 hr. with (a) *N*-acetic acid, (b) 0.5*N*-HCl and (c) 6*N*-HCl. The unoxidized A substance gave more humin during the hydrolysis than did the oxidized material and the *N*-acetic acid hydrolysis products of the oxidized A substance dissolved less readily than the other products in water.

The hydrolysis products arising from the unoxidized and oxidized A substance were analysed simultaneously, and the results are strictly comparable. No difference could be detected in the amino-acid distribution of the materials when run in two directions on paper with collidine (54 hr.) and phenol-ammonia-coal gas (24 hr.) as solvent systems. Both hexosamines were present in the oxidized A substance as demonstrated by the ninhydrin, ammoniacal silver and the hexosamine tests. Quantitatively, however, the spot attributed to chondrosamine was less intense after oxidation of the A substance. Fucose was absent from the hydrolysis products of the oxidized A substance. The hydrolysates (0.5*N*-HCl, 16 hr.) of A substance oxidized at pH 5 (24 hr.) and at pH 7.5 (48 hr.), when run in *n*-butanol-water containing 3% acetic acid in the lower aqueous layer (Aminoff *et al.* 1950) were found to contain only a trace of galactose.

The total fucose content of the A substance. The colorimetric method of Dische & Shettles (1948) was employed using a Hilger Uvispek spectrophotometer. This method determines the total fucose content, free and combined, whereas the method involving the oxidation with periodate (Aminoff *et al.* 1950) only measures the fucose residues with the hydroxyl groups on C atoms 4 and 5 unsubstituted, e.g. free fucose as liberated during mild acid hydrolysis. The material was heated at 100° for 10 min. with the H_2SO_4 reagent, and the colour developed was measured 3–4 hr. after the addition of the cysteine hydrochloride (3%). The absorption curves of

Table 1. Total fucose content of the A substance and some of its degradation products as determined by the colorimetric method of Dische & Shettles (1948)

Substance	Percentage of original A substance	Fucose	
		Percentage	Percentage of original A substance
A substance	—	18.5*	—
A substance 16 hr. <i>N</i> -Acetic acid hydrolysate:			
(a) Diffusate	38	28.2	10.9
(b) Indiffusible material	62	12.4	7.7
A substance (preparation III)	—	18.3	—
A substance oxidized with periodate at pH 5 for 24 hr.	—	0.82	—
A substance oxidized with hypiodite	—	16.3	—

* A preparation of A substance obtained from cyst fluid no. 122 gave 18.4% fucose.

various concentrations of L-fucose, the A substance and an artificial mixture of *N*-acetylglucosamine, *N*-acetylchondrosamine, D-galactose and L-fucose, in the proportions believed to be present in the A substance, were plotted over the range 370–460 $m\mu$. The maximum absorption was found to be at 400 $m\mu$. on this instrument and not at 396 $m\mu$. as quoted by Dische & Shettles (1948) (cf. Dische, Shettles & Osnos, 1949). The fucose content was, therefore, calculated on the basis of the difference in the extinctions at 400 and 430 $m\mu$. The results are given in Table 1 and show that the best preparation of A substance contains 18.5% fucose. Preparation III (Aminoff *et al.* 1950) gave a lower value, 18.3%. The results also indicate that all the fucose in the A substance is destroyed by oxidation with periodate, and that the non-carbohydrate constituents in the complex do not interfere with the fucose determination. In contrast, the A substance oxidized with sodium hypiodite retains its fucose residues largely unchanged.

N-Acetylhexosamine and hexosamine colour reactions. A substance (preparation III) gave a colour equivalent to 10.4% *N*-acetylglucosamine (Morgan & Elson, 1934), whereas the material after oxidation for 24 hr. at pH 5 gave a colour equivalent to 9.5% *N*-acetylglucosamine. There is, therefore, little change in this property as a result of oxidation. Nevertheless, after acid hydrolysis, the oxidation products are found to contain less free hexosamine base (Table 2), and the nature of the distribution of the hexosamines in the undegraded A substance and its oxidation products are given below.

Quantitative determination of the distribution of hexosamines. Portions (5 μ l.) of the hydrolysates (6*N*-HCl, 3 hr., 100°) of the A substance before and after oxidation, or of an artificial mixture of glucosamine and chondrosamine, each of which contained about 50 μ g. of total amino-sugar hydrochloride, were placed 4 cm. apart on a straight line 6 cm. from the top edge of a sheet (32 $\times$ 56 cm.) of Whatman no. 1 paper. The sheet was irrigated for 48 hr. with collidine saturated with water. The collidine had been purified according to the technique described by Partridge (1948*a*) and subsequently carefully fractionated through a Fenske column at 15 mm.

pressure to give a colourless product. The paper was then dried overnight at room temperature and cut into strips, each corresponding to the path of the substances derived from a single spot. Alternate strips were sprayed with the hexosamine reagents (Partridge, 1948*a*) to locate the positions of glucosamine and chondrosamine. The corresponding areas in the unsprayed strips were cut out and eluted with

Fig. 7. The glass chamber used for the elution of sugars and hexosamines from the appropriate areas of a paper-strip chromatogram.

water in a covered glass chamber as shown in Fig. 7. Approximately 1.5 ml. of the eluate was collected directly into the test tubes used for the determination of hexosamines and calibrated at the 2 and 10 ml. levels. The funnel and walls of the tube were then washed down with water to yield a final volume of 2 ml. The hexosamine in the eluate was determined by a modification of the procedure of Elson & Morgan (1933).

The results obtained with an artificial mixture of equal amounts of glucosamine and chondrosamine hydrochlorides and hydrolysates of the undegraded A substance and some of its oxidation products are given in Table 2. It is seen that the

Table 2. *Quantitative distribution of the hexosamines in the A substance and its oxidation products*

Material and treatment	Total hexosamine (base) (%)	Chondrosamine (μ g.)	Glucosamine (μ g.)	Recovery (%)	Ratio glucosamine/chondrosamine
a Chondrosamine and glucosamine hydrochlorides (27 μ g. of each)	—	11.5 $\pm$ 6%	13.5 $\pm$ 7.5%	46	1.17
b A substance (preparation III), hydrolysed with 6 <i>N</i> -HCl for 16 hr.	32	10.0 $\pm$ 10%	8.9 $\pm$ 18%	33	0.89
c A substance (preparation III), oxidized at pH 5 for 24 hr. and then hydrolysed with 6 <i>N</i> -HCl for 16 hr.	24	6.4 $\pm$ 6%	12.5 $\pm$ 2%	44	1.95
d A substance hydrolysed with 6 <i>N</i> -HCl for 3 hr.	37	7.9 $\pm$ 8%	7.8 $\pm$ 15%	33	1.01
e A substance oxidized at pH 4.5 and then hydrolysed with 6 <i>N</i> -HCl for 3 hr.	27	3.7 $\pm$ 13%	7.0 $\pm$ 7%	30	1.90
f A substance hydrolysed with 6 <i>N</i> -HCl for 3 hr.	37	7.1	7.6	36	1.07
g A substance oxidized at pH 7.5 and then hydrolysed with 6 <i>N</i> -HCl for 3 hr.	23	3.9 $\pm$ 23%	7.3 $\pm$ 8%	22	1.90

recovery of the hexosamines is of the order of 30–50% only. It is known, however, that the free hexosamine bases are unstable in an alkaline medium and this would account for the low recovery obtained with collidine (pH 9.4–10.0) as the solvent. In an acidic medium such as butanol-acetic acid, the recovery of the total hexosamine from the paper was up to 90%, but unfortunately it is only in basic solvents such as collidine and butanol-ammonia that a useful separation of the two hexosamines is achieved (Aminoff & Morgan, 1948). The recovery of the hexosamines with collidine as solvent can be much improved by running the chromatograms at 0°, but the rate of irrigation is greatly reduced owing to the increased viscosity of the solvent.

Despite the obvious limitations of the method for the determination of the total hexosamine, the similar behaviour of glucosamine and chondrosamine under these conditions makes possible the determination of the relative distribution of the two hexosamines. In the undegraded A substance glucosamine and chondrosamine are present in equimolecular proportions. After oxidation at pH 5, however, the hexosamine content of the hydrolysate of the oxidized material falls to about 75% of that of the undegraded A substance and the glucosamine/chondrosamine ratio is approximately 2, thus indicating that the fall in hexosamine content can be attributed entirely to the oxidation of half of the total chondrosamine. Oxidation of the A substance at pH 7.5, under the conditions used for the determination of formaldehyde (see below) results in a product which contains 23% hexosamine and represents only 64% of the total hexosamine in the undegraded A substance. The ratio glucosamine/chondrosamine in the material oxidized at pH 7.5 is again 2, and therefore oxidation at the higher pH involves some of the glucosamine and rather more than half of the chondrosamine in the A substance.

Analysis of the products liberated during oxidation

Determination and characterization of formaldehyde as the dimedone derivative. Preliminary determinations of the formaldehyde produced during the oxidation of the A substance, by steam distillation and titration of the bound bisulphite, indicated an apparently greater production of formaldehyde than that determined by the dimedone method. In view of the findings of Bell (1948) that the distillate from the oxidation products of some methylated hexoses contained bisulphite-binding compounds other than formaldehyde, it was decided to dispense with the distillation and to precipitate the formaldehyde *in situ* following the procedure of Reeves (1941). In order to avoid hydrolysis of the oxidized A substance with the possible liberation of other aldehydes (e.g. glyoxal), the oxidized material was not heated with the dimedone reagents. Nicolet & Shinn (1941b) recommend 48–72 hr. standing with the dimedone reagent, whereas Reeves (1941) and Rees (1946) find 16–24 hr. sufficient.

A 2% aqueous solution of the A substance, preparation III (1.0 ml.), was added to 2 ml. of 9% (w/v) NaHCO_3 and 1 ml. of 0.5M-HIO₄ contained in a stoppered tube. After 24 hr. oxidation at room temperature, the following reagents were added with vigorous shaking in the order indicated: N-HCl (3 ml.), approx. N-sodium arsenite (2 ml.), M solution of sodium acetate (2 ml.) and 8% dimedone reagent in 95% ethanol (1 ml.). The resulting mixture was well stirred and kept at room temperature for 48 hr. to permit the complete precipitation of the dimedone derivative. The precipitate

which formed was collected on a porosity 3 sintered-glass funnel, washed with cold water and dried to constant weight in an electric oven at 60°. The precipitate (a) weighed 3.82 mg. More prolonged oxidation, 48 hr., gave a precipitate (b) of similar weight 3.72 mg. The reagents alone gave (c) 0.18 mg. of precipitate. The product from A substance, oxidized (pH 5, 24 hr.) and dialysed, was treated with periodate and NaHCO_3 and immediately HCl, sodium arsenite and the other reagents were added. A precipitate (d) slowly developed and increased in amount on long standing. After a week at room temperature the precipitate was voluminous and proved difficult to filter. Drying the material at 60° gave rise to charring and the substance became gummy.

The melting points of the different precipitates were not sharp. After recrystallization from ethanol, in which precipitates a, b and c were readily and d sparingly soluble, a and b gave long needles, m.p. 185° (uncorr.). Precipitate d crystallized very poorly and softened slowly with charring between 178 and 196°. The ready solubility of a, b and c in ethanol in contrast to the low solubility of d indicated that the weight of the true dimedone derivative of formaldehyde could be obtained as the weight of the precipitate that was soluble in ethanol.

By this simple modification in technique it was shown that 32.83 mg. of the purest A substance, on oxidation for 24 hr. and filtration 20 hr. after the addition of dimedone reagents, gave rise to 5.21 mg. of ethanol-soluble formaldehyde-dimedone derivative of which 0.33 mg. was derived from the reagents. 1 g.mol. of formaldehyde is therefore obtained from 1964 g. of A substance. For preparation III a higher value, 2551 g., was obtained.

After removal, by filtration, of the formaldehyde-dimedone derivative from the oxidation products of the purest A substance the filtrate was dialysed and dried from the frozen state. The hexosamine content and the relative distribution of the glucosamine and chondrosamine residues were determined on a hydrolysate (6N-HCl for 3 hr.) of the material and the results obtained are given in Table 2, g.

After oxidation under the same conditions methyl N-acetylglucosaminide gave only a trace of precipitate, whereas N-acetylglucosamine gave 93–95% of the theoretical amount of formaldehyde as dimedone derivative (m.p. 186°, uncorr.) identical with the material formed on the addition of formaldehyde to dimedone.

Determination of the acid produced during oxidation. It is only recently that the estimation of the acid produced during oxidation of substances of large molecular weight with periodate has been investigated as a method of end-group assay. Two methods have been suggested to overcome the difficulty of 'over acid production' (a) the slow oxidation at room temperature with the sparingly soluble KIO₄ (Halsall, Hirst & Jones, 1947), and (b) the more rapid oxidation with NaIO₄ at a lower temperature (Potter & Hassid, 1948).

(a) Employing the method of Halsall *et al.* (1947). The A substance (2 ml. of 2% solution of preparation III) was treated in a 25 ml. flask with 5 ml. of 0.254M-HIO₄, which had been brought to pH 4.5 (methyl red) with NaOH, and 4 ml. of 25% (w/v) KCl solution were added. The solution was then made up to 25 ml. with distilled water. Throughout the period of oxidation the mixture was kept in the dark and at room temperature. It was shaken frequently to disperse the crystalline precipitate of KIO₄. Samples (4 ml.) were withdrawn at intervals over a period of 237 hr. and added to flasks containing 0.75 ml. of ethylene glycol (which had

Table 3. *Rate of reduction of periodate and production of acid on oxidation of the A substance with KIO_4 and NaIO_4*

KIO_4		NaIO_4 at 2-3°		
Period of oxidation (hr.)	Weight of A substance (preparation III) giving 1 g. mol. of acid (g.)	Period of oxidation	Weight of A substance (preparation III)	
			Giving 1 g. mol. of acid (g.)	Reducing 1 g. mol. of periodate (g.)
26	1935	7 min.	11840	304
75	1575	185 min.	—	255
170	1215	345 min.	3480	258
215	1030	27 hr.	2190	250
237	1092	47 hr.	1740	235
		70 hr.	1535	229

previously been distilled over NaOH) and kept in the dark for 15 min. The excess periodate slowly dissolved and the resulting solution was titrated against 0.01 N-Ba(OH)₂ using methyl red as indicator. A reagent blank (4 ml.) was treated similarly and the appropriate corrections applied. The production of acid had reached a maximum in 215 hr. (Table 3), after which time 1030 g. of A substance would have produced 1 g. equiv. of acid.

(b) Oxidation with NaIO_4 (Potter & Hassid, 1948); A substance (2 ml. of 2% solution of preparation III) contained in a 25 ml. flask was treated with 5 ml. 0.207 M-HIO₄ brought to pH 4.5 (methyl red) with NaOH and 3.75 ml. of 10% (w/v) NaCl added and made up to 25 ml. with water. The solution was kept in the dark at 2-3°. At intervals samples were withdrawn, added to ethylene glycol as described above and after standing for 1 hr. in the dark, titrated with 0.01 N-Ba(OH)₂. At the same time portions of the solution were added to alkaline arsenite solution and the course of the periodate reduction followed in the usual manner. The reduction was practically complete in 3 hr., after which time 255 g. of A substance would have reduced 1 g. mol. of periodate. After that period of oxidation 1535 g. A substance would have liberated 1 g. equiv. of acid. Details of the experimental results are given in Table 3. Under the same conditions 1 g. equiv. of acid would have been liberated by 2120 and 1790 g. of the purest specimen of A substance after 26 and 51 hr., respectively.

After neutralization with Ba(OH)₂ the oxidation products of the purest A substance were pooled, thoroughly dialysed and dried from the frozen state. The products of hydrolysis (3 hr., 6 N-HCl at 100°) of this material contained 27% of hexosamine, or 74% of the value (37%) for the undegraded A substance. Quantitative chromatographic analysis of this material gave a glucosamine:chondrosamine ratio of 2:1 (Table 2e).

Maltose and lactose behaved similarly when oxidized under the same conditions. The periodate was rapidly reduced, the reaction being complete in about 1 hr. 6 g. mol. of periodate were reduced per g. mol. of disaccharide, instead of the expected 5 g. mol. Again, the rate of acid production was much slower than the rate of reduction of periodate, and ultimately exceeded the theoretical amount of 3 g. mol./g. mol. of disaccharide. The theoretical amount was, however, obtained for both maltose and lactose after a period of oxidation of 25 hr.

Determination of hydroxyamino nitrogen. The method employed was based on that described by Gordon, Martin & Synge (1943). The oxidations were carried out in Conway units and the NH₃ liberated absorbed in boric acid.

Boric acid (1 ml., 2% solution) which contained a mixed bromocresol green-methyl red indicator (Ma & Zuazaga, 1942) was placed in the inner chamber of a Conway unit and the following solutions were added to the outer chamber in the order given: (a) the substance to be oxidized contained in 0.5 ml. water; (b) 9% (w/v) glycine solution (0.1 ml.); (c) N-NaOH (0.25 ml.); (d) M-HIO₄ (0.2 ml.); and (e) saturated solution of K₂CO₃ (1 ml.). The unit was then immediately sealed, carefully rotated to ensure mixing and kept at room temperature. After the given period of oxidation the contents of the inner chamber were titrated with 0.01 N-HCl to a faint-pink colour. The results are given in Table 4 which also includes the results obtained with several substances that were used for control purposes. It will be seen that ethanolamine gives 94-98% recovery, depending on the time of oxidation. Threonine yields its N quantitatively as NH₃, whereas glucosamine hydrochloride gives only about 70% of its N in this form. N-acetylglucosamine reacts slowly and liberates rather less than half of its N as NH₃ in 24 hr. Undegraded A substance gives no NH₃ after 3 hr. oxidation and a trace only after 24 hr., and there is, therefore, no possibility of liberated NH₃ interfering with the determination of the acid produced during the oxidation of the A substance. An acetic acid hydrolysate of A substance gives only 7-8% of the total N as hydroxyamino N. This could arise from (1) N-acetylhexosamine, (2) hexosamine or (3) the hydroxyamino-acids, serine or threonine. The acetic acid hydrolysis products of A substance which had previously been oxidized for 24 hr. at pH 5 give more than twice as much NH₃ as the hydrolysate of unoxidized A substance. This result could be attributed to the greater lability of the oxidized A substance to acid hydrolysis, and suggests that some of the more labile linkages involve the hexosamines and/or hydroxyamino-acids. The value obtained for the 6 N-HCl hydrolysis products (Table 4) represents the total hydroxyamino N and amide N. The hydrolysate contained 37% hexosamine (corrected for the high values obtained with 6 N-HCl) and, assuming 68% conversion of the hexosamine N into NH₃, would account for 34% of the 47% N appearing as NH₃ on oxidation. Thus 13% of the total N is due to hydroxyamino acids and amide N. This represents quite a high proportion (34%) of the

Table 4. *Hydroxyamino-nitrogen content of miscellaneous substances as determined by the liberation of ammonia on oxidation with periodate*

Substance	Treatment	Period of oxidation (hr.)	Hydroxyamino N as percentage of total N
Ethanolamine	—	2	94.5
Ethanolamine	—	4	95.0
Ethanolamine	—	24	98.0
Threonine	—	2.5	99.0
Threonine	—	21	100.0
Glucosamine hydrochloride	—	3	68.5
Glucosamine hydrochloride	—	24	73.0
N-Acetylglucosamine	—	3	27.0
N-Acetylglucosamine	—	24	42.0
A substance (preparation III)	—	3	0
A substance (preparation III)	—	24	0.075
A substance (preparation III)	Hydrolysed with N-acetic acid for 16 hr.	4	6.9
A substance (preparation III)	Hydrolysed with N-acetic acid for 16 hr.	24	7.9
A substance (preparation III)	Oxidized at pH 5 for 24 hr. and then hydrolysed with N-acetic acid for 16 hr.	24	19.5
A substance	Hydrolysed with 6N-HCl for 2 hr.	3.5	47.3

Table 5. *Apparent fucose content of A substance and of some of its degradation products as determined by the acetaldehyde liberated on oxidation with periodate*

Substance	Treatment	Fucose (%)
Fucose	—	88, 92
A substance	—	0.30
A substance (preparation III)	—	0
A substance (preparation III)	Oxidized for 24 hr. at pH 5 and then hydrolysed at 100° with N-acetic acid for 16 hr.	0.98
A substance (preparation III)	N-Acetic acid, 16 hr.	11.6
A substance (preparation III)	N-Acetic acid, 48 hr.	12.3
A substance	N-Acetic acid, 48 hr.	12.9
A substance	N-Acetic acid, 16 hr. indiffusible material	0.56*

* In terms of the original A substance this represents 0.35% fucose.

maximum amino-acid N which is 38% of the total N of the A substance as determined by the ninhydrin method (Van Slyke, Dillon, MacFadyen & Hamilton, 1941).

Determination and characterization of the acetaldehyde formed on oxidation. Fucose and threonine give rise to acetaldehyde on oxidation with periodate under the same conditions (Nicolet & Shinn, 1941a; Shinn & Nicolet, 1941). The successful determination of threonine in Conway units (Winnick, 1942) suggested the adoption of the method for the determination of fucose and, although no serious attempt was made to develop an absolutely quantitative technique, yields of acetaldehyde equivalent to about 90% of the fucose taken were readily obtained.

NaHSO₃ (1.5 ml. of 0.25M solution) was pipetted into the central chamber of a Conway unit. The substance to be analysed contained in a volume of 0.5 ml. was placed in the outer chamber and 1 ml. of 0.1M-K₃PO₄ added. The concentration of K₃PO₄ was adjusted so that 1 ml. exactly neutralized, to pH 7.0, the 0.5 ml. of 0.2M-HIO₄ which was finally added. The greased lid was then slipped into position, the unit was rotated in the usual manner, and kept at room temperature for the period of oxidation (2–5 hr., depending on temperature). A drop of 1% starch solution was then added to the inner chamber and the excess bisulphite oxidized with I₂ until a faint-blue tint was obtained. The aldehyde-bisulphite compound was decomposed with powdered Na₂HPO₄ or NaHCO₃ and titrated with 0.005N-I₂. The results

obtained with two preparations of A substance and some of their degradation products are given in Table 5. Fucose, included with each set of determinations, gives rather variable results; the average recovery of acetaldehyde was 90% of the theoretical. The values for the fucose content of the hydrolysis products are, therefore, corrected for a low recovery of this order. Threonine behaves more consistently and gives 95% of the theoretical yield of acetaldehyde. A substance does not liberate any appreciable amount of acetaldehyde during oxidation, hence acetaldehyde is not involved in the formaldehyde-dimedone determinations. The acetic acid hydrolysis products of A substance oxidized for 24 hr. at pH 5 give little acetaldehyde compared with the corresponding hydrolysate of the unoxidized A substance. The acetic acid hydrolysate of two preparations of A substance gave acetaldehyde equivalents to 12.3 and 12.9% fucose respectively. This is the maximum amount of free fucose obtained after mild acid hydrolysis and will be designated as acid-labile fucose. It represents about three-quarters of the total fucose, 18.5%, present in the A substance (Table 1). The indiffusible fraction of the A substance after a partial hydrolysis (16 hr., N-acetic acid) (Aminoff *et al.* 1948) has a total fucose content of 12.4% which, expressed in terms of the original A substance before hydrolysis, represents a fucose content of 7.7% (Table 1) and therefore still contains 2.2% of the original A substance as acid-labile fucose residues. This material yields a negligible amount of

acetaldehyde on oxidation with periodate, equivalent to a fucose content of 0.56% (i.e. 0.35% of the original A substance used for the hydrolysis).

The acetaldehyde obtained on the oxidation of the products of mild acid hydrolysis of A substance did not arise from threonine since chromatographic analysis shows that this amino-acid is not present in the free state in the acetic acid hydrolysates. Furthermore, the oxidation of the A substance with periodate involves all the fucose residues but not threonine, and, although an acetic acid hydrolysate of both the oxidized and the unoxidized A substance gave a little NH_3 on subsequent oxidation (Table 4), the former material gave very little acetaldehyde. The amount of acetaldehyde arising from threonine would, therefore, appear to be negligible in the mild acid hydrolysis products examined.

The acetaldehyde formed during the oxidation was characterized as the dimedone derivative by absorbing the volatile vapours produced in the outer chamber of the Conway unit into a solution of dimedone at pH 4.5 (acetate buffer), contained in the inner chamber. At the end of 24 hr. several large needle-shaped crystals were formed. After recrystallization from ethanol, the crystals melted sharply at 138–139° (cf. Weinberger, 1931). A mixture of the crystals obtained from the oxidation products of fucose and from an acetic acid hydrolysate of A substance showed no depression of melting point.

DISCUSSION

Course of oxidation. The rate of serological inactivation of A substance during oxidation with periodate, at pH values between 5 and 10, was most rapid at pH 5 (acetate buffer) and decreased with rise in pH, with no inactivation after 3 hr. at pH 10 (borate buffer). The stability at the higher pH is remarkable in view of the known lability of the blood-group substances to alkali. Although the presence of borate has little effect on the rate of reduction of periodate at pH 2.4 and 7.5 the slower rate of inactivation of the A substance at the higher pH values might well be due to the formation of chelation complexes with the *cis*-hydroxyl groups. Both serological properties are lost on oxidation, but the haemolysis-inhibiting (Forssman) activity is destroyed less rapidly than the capacity to inhibit iso-agglutination.

The pH values chosen to follow the course of oxidation were those optimal for the determination of the acid and aldehyde produced, namely pH 5.0 and 7.5 respectively. Although the rate of serological inactivation is less rapid at the higher pH, the oxidation is more extensive. Oxidation at pH 5 is essentially complete in about 6 hr., while after 3 days at pH 7.5 substantial reduction of the periodate is still occurring. A further marked difference in behaviour of the A substance is observed after the initial oxidation at pH 5 or 7.5. Thus re-oxidation of material previously oxidized at pH 5 results in further reduction of periodate only at pH 7.5, whereas the material initially oxidized at pH 7.5 reduces more periodate at both pH 5 and 7.5. This suggests

that the initial oxidation at pH 7.5 exposes additional structures susceptible to oxidation. A similar instance of more extensive oxidation in an alkaline medium has been observed by Head (1947) during the oxidation of β -methylglucoside and β -methylcellobioside. The cause of this overoxidation was attributed by this author to the sensitivity of the primary oxidation products to hydrolysis by alkali with the production of structures susceptible to further oxidation. This interpretation is not applicable to our observations, since treatment of the material oxidized at pH 5 with sodium bicarbonate at pH 9.4 for 24 hr. and subsequent retreatment with periodate at pH 5 and 7.5 shows no difference in the behaviour from that of material oxidized at pH 5 without subsequent treatment with sodium bicarbonate. It must be concluded, therefore, that acid and alkaline periodate differ markedly in their oxidizing capacity.

Oxidation of the A substance reaches completion in an acid medium, but the amount of periodate reduced depends on the conditions used. Thus, in acetate buffer (pH 5) 1 g.mol. of periodate is reduced by 430 g. of A substance after 4 hr. and 364 g. after 24 hr., whereas the corresponding figure, the amount of A substance reducing 1 g.mol. of periodate, obtained when determining the formic acid produced on oxidation with sodium periodate at pH 4.5 is 260 g. after 26 hr. These values are a measure of the total reduction of periodate due to (a) specific oxidation of the sugar molecules containing adjacent, unsubstituted hydroxyl groups and (b) the nonspecific 'overoxidation' of the primary oxidation products.

Constituents involved in the oxidation. Removal, by dialysis, of excess periodate and buffer after oxidation and subsequent qualitative chromatographic analysis of the hydrolysates of the oxidation products reveals no change in the amino-acid composition of the A substance, but indicates the absence of galactose and fucose. Quantitative chromatographic analysis of the undegraded A substance shows the presence of glucosamine and chondrosamine in equimolecular proportions, whereas the corresponding hydrolysate of the oxidized material contains about three-quarters of the hexosamine originally present in the A substance and the ratio of glucosamine to chondrosamine is 2:1. Thus oxidation with periodate at pH 5 involves the fucose, galactose and half the *N*-acetylchondrosamine residues.

Simple substances liberated during oxidation. Formaldehyde and formic acid, but no acetaldehyde or ammonia, are produced during the oxidation of the A substance and the determination of the former pair of substances is not subject to interference by the latter.

1 g.mol. of formaldehyde is produced by the oxidation of 1964 g. of A substance. The formaldehyde

could arise from the serine, galactose or *N*-acetylhexosamine residues. No ammonia is liberated during the oxidation and hence serine is not involved.

The measurement of the acid produced presents no difficulty, the problem, however, devolves on the interpretation of the results obtained. Halsall *et al.* (1947) showed that the time required for the production of the theoretical amount of acid varied for different sugars and with both potassium periodate (Halsall *et al.* 1947) and sodium periodate (Potter & Hassid, 1948) the period of oxidation of a large molecular substance was empirically fixed by the time required to produce the theoretical amount of acid from a disaccharide which possessed the structural unit responsible for the acid production in the material under investigation. Thus β -methyl maltoside and maltose were chosen as the standard substances when determining the chain length of the glycogens and starches. The acid formed during the oxidation of the A substance could arise from the galactose, fucose or the reducing terminal *N*-acetylhexosamine residues, but in the absence of suitable simple derivatives of the latter two sugars, lactose and maltose, were used as standard substances to establish a suitable period for oxidation. Maclay, Hann & Hudson (1939) found that α - and β -methyl-L-fucosides reduced two molecules of periodate within 1 hr., but the acid formed was not determined. Lactose and maltose on oxidation with sodium periodate produce the expected quantity of formic acid after 25 hr. and about 2000 g. of A substance produce 1 g. equiv. of an acid after oxidation with either potassium or sodium periodate for the same length of time. More prolonged oxidation of lactose, maltose and the A substance by either method gives an increase in acid production.

The reducing end group. The blood-group A substance readily gives a colour with Ehrlich's reagent, under the conditions used for the determination of *N*-acetylglucosamine (Morgan & Elson, 1934) and the intensity of the colour is equivalent to approximately a quarter of the total calculated *N*-acetylhexosamine content (Aminoff *et al.* 1950). *N*-acetyl- and *N*-benzoyl-methylglucosaminides, on the other hand, do not give this reaction, and one might therefore conclude that the reactive *N*-acetylhexosamine residues in the A substance form the reducing end groups (I, where R or R' is a glycoside or a peptide residue).

The yields of formaldehyde obtained from our purest material and from a partially purified preparation of A substance (preparation III) are proportional to the *N*-acetylhexosamine colour given by these materials and the origin of the formaldehyde could therefore be attributed to the reducing *N*-acetylhexosamine residues which on oxidation with periodate would result in cleavage of the C-5—C-6 linkage (I). This conclusion does not conflict with the observation that the A substance after oxidation develops the same amount of colour as does the original A substance when treated with mild alkali and Ehrlich's reagent, since the colour reaction only involves the groups at C atoms 1 and 2 of the *N*-acetylhexosamine molecule. That an *N*-acetylhexosamine residue is involved in oxidation is further substantiated by the observation that the hydrolysis products of the oxidized A substance show a decrease in hexosamine content, the value of which is again equivalent to a quarter of the total hexosamine. On the basis of quantitative paper-chromatographic analysis of the A substance oxidized at pH 5.0 and 7.5 the decrease in hexosamine content is found to be due principally to a loss in chondrosamine which would suggest that the terminal *N*-acetylhexosamine residue involved in the first stages of oxidation is *N*-acetylchondrosamine. It has been found, however, that *N*-acetylchondrosamine gives considerably less colour than *N*-acetylglucosamine and that the earlier report (Morgan & Elson, 1934), that equal weights of the two *N*-acetylhexosamines give the same amount of colour, is incorrect. This observation will be considered in greater detail in a further paper. As *N*-acetylchondrosamine gives about one-fifth of the colour given by *N*-acetylglucosamine one must conclude that *N*-acetylglucosamine also contributes to the total colour given by the A substance. The contribution of *N*-acetylglucosamine to the total colour can arise in two ways. First, by an *N*-acetylglucosamine residue also being present as a reducing end group. In that case the hydroxyl group on C atom 6 of the proposed *N*-acetylglucosamine end groups must be substituted to account for the low yield of formaldehyde which is equivalent to only one *N*-acetylhexosamine unit present as a reducing end group. A second possibility, as formulated in an earlier contribution (Aminoff & Morgan, 1949), is that the terminal *N*-acetylchondrosamine residue is linked

through C atom 1 of an *N*-acetylglucosamine residue, and that the mild alkaline treatment, under the conditions used for the determination of *N*-acetylhexosamine, breaks this glycoside linkage and thus transforms the *N*-acetylglucosamine into an end group and renders it reactive. These changes would satisfactorily account for the slower development of the maximum colour given by the A substance on heating with the alkali (14 min.) as compared with the time required (4 min.) for the free *N*-acetylhexosamines. Further evidence for differentiating between these two structural possibilities is now being sought.

The mode of combination of fucose residues. The evidence for the form and sequence of fucose residues in the A substance is deduced from the following observations: (1) The total fucose content of the A substance is 18.5%. (2) Undegraded A substance gives no acetaldehyde on oxidation with periodate.

(3) Most of the fucose is readily liberated in an uncombined state by mild acid hydrolysis, i.e. *N*-acetic acid. (4) The maximum amount of acid-labile fucose is obtained by hydrolysis with *N*-acetic acid at 100° for 48 hr., as determined by the acetaldehyde liberated on oxidation with periodate, and represents 13% of the A substance (Aminoff *et al.* 1950), i.e. three-quarters of the total fucose. The remaining 5.5% of the fucose in the A substance represents the acid-stable fucose residues. (5) The indiffusible material obtained by dialysis of an incomplete (16 hr.) acetic acid hydrolysate of the A substance (Aminoff *et al.* 1948), a material which still contains some acid-labile fucose, gives no acetaldehyde on oxidation with periodate. (6) Oxidation of the A substance with periodate at pH 5 results in the destruction of all the fucose, i.e. both the acid-labile and acid-stable fucose residues. (7) The mild acid hydrolysis products of the A substance after

oxidation at pH 5 for 24 hr., gives a negligible amount of acetaldehyde on re-oxidation with periodate. (8) The fucose content is little changed after oxidation of the A substance with sodium hypiodite.

From the observations given under (2) and (8) above it is most probable that the fucose residues are not present as reducing end groups but form either non-reducing terminal residues or are situated in the main carbohydrate chain. Since three-quarters of the total fucose in the A substance is readily liberated on complete mild acid hydrolysis (48 hr. *N*-acetic acid), in an uncombined state and without extensive hydrolysis of the other carbohydrate constituents, it would appear that these fucose residues are terminal and are therefore present either as non-reducing end groups or are closely linked at the end of the chain. It might be concluded from the ease of hydrolysis of the glycoside bonds that the terminal fucose units possess a furanose structure (II). The

oxidation of the A substance for 24 hr. at pH 5.0 with periodate would then presumably involve the hydroxyl groups at C atoms 2 and 3 and mild acid hydrolysis of the oxidation products will yield a material which on further oxidation should give rise to the same amount of acetaldehyde as is obtained from the undegraded A substance after acid hydrolysis and oxidation alone, viz. an amount equivalent to 13% of fucose. It has been found, however, that A substance after oxidation at pH 5 for 24 hr., followed by hydrolysis with acetic acid, and re-oxidation with periodate yields acetaldehyde equivalent to a content of less than 1% fucose. The results obtained, therefore, indicate that the hydroxyl groups at C atoms 3 and 4, at least, of the acid-labile fucose units, are involved in the initial oxidation and must therefore be unsubstituted. The acid-labile fucose units, in spite of their acid lability, would therefore appear to be present as pyranosides.

The acid produced during the oxidation of the A substance with periodate can therefore be attributed to the terminal fucose residues, with the hydroxyl groups on C atoms 2, 3 and 4 unsubstituted, as shown in III. The amount of acid produced on oxidation of the A substance for 25 hr., however, corresponds to only two-thirds of the acid-labile fucose, and therefore to a half of the total fucose content of the A substance.

Since no acetaldehyde is formed on oxidation of the mild acid hydrolysis products of the A substance oxidized at pH 5 (Table 5) the hydroxyl groups at C atoms 3 and 4 of all the acid-labile fucose residues are involved in the initial oxidation. The non-terminal acid labile fucose residues (IV), representing a quarter of the total fucose, are presumably substituted at C atoms 1 and 2 in the undegraded A substance. (R is a glycoside or peptide residue.)

Assuming, as seems most probable, that the terminal and non-terminal acid-labile fucose residues are closely linked, two modes of combination are possible (V or VI).

After hydrolysis of A substance for 16 hr. with acetic acid (Aminoff *et al.* 1948) 38% by weight diffuses through a cellophan membrane. The indiffusible residue contains all the acid-stable fucose

(about a quarter of the total fucose originally present in the A substance) together with about 14% of the acid-labile fucose (three-quarters of the total fucose) which has not been liberated owing to the hydrolysis being incomplete after 16 hr. No significant amount of acetaldehyde was obtained on oxidation of this

indiffusible material (Table 5) and thus C atom 1 of the non-terminal acid-labile fucose residue in this partial acid hydrolysis product is still glycosidically bound, and the two fucose residues are therefore presumably linked as shown in V. The configuration of the glycoside linkages is not known.

The acid-stable fucose residues, which represent the remaining quarter of the total fucose of the A substance, are also, like the acid-labile fucose residues, involved in the oxidation with periodate as

No acetaldehyde

Partial mild hydrolysis with
acetic acid

Oxidation of
indiffusible moiety

Acetaldehyde

demonstrated by the negligible fucose content of the A substance after oxidation at pH 5.0 for 24 hr. (Table 1). The hydroxyl group on C atom 3 of the acid-stable fucose residues must therefore be unsubstituted. The other unsubstituted hydroxyl group is most probably on C atom 2, in contrast to the acid-labile non-terminal fucose residue (IV) which has a substituted hydroxyl group at C atom 2. As such this acid-stable fucose residue would be apparently less sensitive to acid hydrolysis since both the linkages at C atoms 1 and 4 must be broken before acetaldehyde is liberated on oxidation with periodate; whereas the liberation of acetaldehyde from a fucose residue linked at C atoms 1 and 2 (IV) depends on the fission of only the linkage involving C atom 1.

reducing terminal residues. The evidence for the mode of combination of the galactose and *N*-acetylglucosamine residues in the A substance is at present only circumstantial and is derived from the study of the mild acid hydrolysis products of this material. The undegraded A substance shows no pneumococcal type XIV or anthrax C polysaccharide specificity, but both these immunological characters, and an enhancement in the pre-existing heterophile (or Forssman) property, develop as a result of mild hydrolysis with acid (Ivanovics, 1940*b*; Kabat, Baer, Bezer & Knaub, 1948; Aminoff *et al.* 1948). It is well known that the immunologically active polysaccharides obtained from the Forssman antigen, the pneumococcal type XIV and the anthrax C antigens contain *N*-acetylglucosamine and a hexose which is

The galactose and N-acetylglucosamine residues. The production of formaldehyde, formic acid and acetaldehyde under different conditions has given some indication of the position of linkages joining the various fucose residues, and also the *N*-acetylchondrosamine end group in the A substance. The yield of formaldehyde and formic acid is used to determine the chain length; the formaldehyde liberated is a measure of the number of reducing end groups and the formic acid of the number of non-

most probably galactose (Brunius, 1936; Chase & Landsteiner, 1939; Goebel, Beeson & Hoagland, 1939; Ivanovics, 1940*a*). It seems probable, therefore, that the *N*-acetylglucosamine and galactose units are closely associated in the A substance and in its mild acid hydrolysis products, and account for the cross-reactivity observed.

The yield of formaldehyde on oxidation of the A substance is fully accounted for by the reducing *N*-acetylchondrosamine residue, and hence the oxida-

tion of galactose presumably involves rupture between C atoms 2 and 3 or possibly 3 and 4.

The amino-acid moiety. According to Nicolet & Shinn (1939) and Posternak & Pollaczek (1941) the acetylated hydroxyamino-acids give no ammonia when treated with periodate. *N*-Acetylglucosamine, however, gives an appreciable amount of ammonia. Neuburger (1941) also detected ammonia in the oxidation products of ethyl *N*-benzoylglucosamate. In these instances, therefore, the acyl group offers little or no protection to the amino group. Undegraded A substance gives no ammonia and hence the hydroxyl groups attached to carbon atom 3 in the *N*-acetylhexosamine residues are most probably substituted. In spite of the high content of the hydroxyamino-acids in the amino-acid moiety of the A substance (34% of the maximum amino-acid nitrogen is present in the form of hydroxyamino N), neither serine nor threonine can be situated at the end of a peptide chain (VII) or be attached by ester linkages involving the carboxyl group to the primary (VIII) or secondary (IX) alcohol groups in the sugar residues.

Certain of the hydroxyl groups of fucose, galactose and part of the hexosamine units are unsubstituted, to permit oxidation, and thus the points of attachment of the amino-acid residues are limited to the hydroxyl groups that have not been involved in oxidation.

Conclusions. On the basis of the analytical results available at present, the carbohydrate constituents of the A substance, galactose, fucose, *N*-acetylglucosamine and *N*-acetylchondrosamine, are present in equimolecular proportions and correspond to a unit with a minimum molecular weight of 968

(Aminoff *et al.* 1950). The oxidation results, however, suggest that the smallest unit with a reducing end group and a non-reducing terminal residue and containing three different forms of fucose is about 3880 and would thus contain four galactose, four fucose, four *N*-acetylglucosamine and four *N*-acetylchondrosamine residues. This represents an 'average' and not necessarily a repeating structural unit.

The results of periodate oxidation taken in conjunction with those of mild acid and alkaline hydrolysis have suggested the form and sequence of certain of the sugar molecules in the carbohydrate moiety of the A substance. Further investigations to establish the structure of this important biological material are in progress.

SUMMARY

1. Fucose, galactose and a half of the total *N*-acetylchondrosamine residues of the A substance are oxidized by periodate at pH 5.

2. Formaldehyde and formic acid, but no acetaldehyde or ammonia, are liberated during the oxidation.

3. The formaldehyde and formic acid (25 hr. oxidation) are produced in approximately equimolecular proportions and correspond to a minimum molecular weight of the order of 2000.

4. The hydroxyamino-acids, serine and threonine, are not present as terminal residues in the amino-acid-containing moiety and are not involved as single molecules in simple esterification with the hydroxyl groups of the sugar units.

5. Certain aspects of the structure of the A substance are discussed in the light of the results obtained.

REFERENCES

- Aminoff, D. & Morgan, W. T. J. (1948). *Nature, Lond.*, **162**, 579.
- Aminoff, D. & Morgan, W. T. J. (1949). *Biochem. J.* **44**, xxi.
- Aminoff, D., Morgan, W. T. J. & Watkins, W. M. (1948). *Biochem. J.* **43**, xxxvi.
- Aminoff, D., Morgan, W. T. J. & Watkins, W. M. (1950). *Biochem. J.* **46**, 426.
- Baer, H., Dische, Z. & Kabat, E. A. (1948). *J. exp. Med.* **88**, 59.
- Bell, D. J. (1948). *J. chem. Soc.* p. 992.
- Bendich, A., Kabat, E. A. & Bezer, A. E. (1947). *J. Amer. chem. Soc.* **69**, 2163.
- Bray, H. G., Henry, H. & Stacey, M. (1946). *Biochem. J.* **40**, 124.
- Brunius, F. E. (1936). *Chemical Studies on the True Forssman Hapten, the corresponding Antibody, and Their Interaction*. Stockholm: Fahlcrantz.
- Chase, M. W. & Landsteiner, K. (1939). *Ann. Rev. Biochem.* **8**, 579.
- Dische, Z. & Shettles, L. B. (1948). *J. biol. Chem.* **175**, 595.
- Dische, Z., Shettles, L. B. & Osnos, M. (1949). *Arch. Biochem.* **22**, 169.
- Elson, L. A. & Morgan, W. T. J. (1933). *Biochem. J.* **27**, 1824.
- Goebel, W. F., Beeson, P. B. & Hoagland, C. L. (1939). *J. biol. Chem.* **129**, 455.
- Gordon, A. H., Martin, A. J. P. & Synge, R. L. M. (1943). *Biochem. J.* **37**, 538.
- Halsall, T. G., Hirst, E. L. & Jones, J. K. N. (1947). *J. chem. Soc.* pp. 1399, 1427.
- Head, F. S. H. (1947). *J. Text. Inst., Manchr.* **38**, T. 389.
- Ivanovics, G. (1940a). *Z. Immunforsch.* **97**, 402.
- Ivanovics, G. (1940b). *Z. Immunforsch.* **98**, 373.
- Jackson, E. L. (1944). *Organic Reactions*, vol. II, p. 341. New York: J. Wiley.
- Jones, J. K. N. (1946). *Rep. Progr. Chem.* **43**, 180.
- Kabat, E. A., Baer, H., Bezer, A. E. & Knaub, V. (1948). *J. exp. Med.* **88**, 43.
- Kabat, E. A., Bendich, A., Bezer, A. E. & Knaub, V. (1948). *J. exp. Med.* **87**, 295.
- Ma, T. S. & Zuazaga, G. (1942). *Industr. Engng Chem. (Anal. ed.)*, **14**, 280.
- Maclay, W. D., Hann, R. M. & Hudson, C. S. (1939). *J. Amer. chem. Soc.* **61**, 1660.
- Morgan, W. T. J. (1946). *Biochem. J.* **40**, xv.

- Morgan, W. T. J. & Elson, L. A. (1934). *Biochem. J.* **28**, 988.
Morgan, W. T. J. & King, H. K. (1943). *Biochem. J.* **37**, 640.
Neuberger, A. (1941). *J. chem. Soc.* p. 47.
Nicolet, B. H. & Shinn, L. A. (1939). *J. Amer. chem. Soc.* **61**, 1615.
Nicolet, B. H. & Shinn, L. A. (1941*a*). *J. Amer. chem. Soc.* **63**, 1456.
Nicolet, B. H. & Shinn, L. A. (1941*b*). *J. biol. Chem.* **139**, 687.
Owen, L. N. (1943). *Rep. Progr. Chem.* **40**, 115.
Partridge, S. M. (1948*a*). *Biochem. J.* **42**, 238.
Partridge, S. M. (1948*b*). *Biochem. J.* **42**, 251.
Posternak, T. & Pollaczek, H. (1941). *Helv. chim. Acta*, **24**, 1190.
Potter, A. L. & Hassid, W. Z. (1948). *J. Amer. chem. Soc.* **70**, 3488.
Rees, M. W. (1946). *Biochem. J.* **40**, 632.
Reeves, R. E. (1941). *J. Amer. chem. Soc.* **63**, 1476.
Shinn, L. A. & Nicolet, B. H. (1941). *J. biol. Chem.* **138**, 91.
Van Slyke, D. D., Dillon, R. T., Macfadyen, D. A. & Hamilton, P. (1941). *J. biol. Chem.* **141**, 627.
Weinberger, W. (1931). *Industr. Engng Chem.* (Anal. ed.), **3**, 365.
Winnick, T. (1942). *J. biol. Chem.* **142**, 461.

Kinetic Studies of the Metabolism of Foreign Organic Compounds

1. THE FORMATION OF BENZOIC ACID FROM BENZAMIDE, TOLUENE, BENZYL ALCOHOL AND BENZALDEHYDE AND ITS CONJUGATION WITH GLYCINE AND GLUCURONIC ACID IN THE RABBIT

By H. G. BRAY, W. V. THORPE AND K. WHITE
Physiology Department, Medical School, University of Birmingham

(Received 16 June 1950)

Investigations carried out in this laboratory in recent years (summarized by Thorpe, 1950) have dealt with the factors determining the metabolic pathways followed by foreign aromatic compounds when several such pathways are available. While it is to be expected that all possible reactions will occur simultaneously, the proportions of the dose which actually follow the various routes differ greatly. Many factors may theoretically contribute to these differences, e.g. the efficiency of the necessary enzyme systems, the availability of conjugating molecules (e.g. glycine) and the availability of free energy for synthetic reactions. All such factors, taken together, determine the velocity of each individual reaction in the intact animal, so that knowledge of the relevant velocity constants is likely to prove of value in considering the detailed metabolism of foreign organic compounds. Since such information does not appear to be available in the literature we have planned a series of investigations in which we hope to compare the velocities of typical metabolic reactions which foreign organic compounds may undergo in the rabbit. In this first investigation the formation of benzoic acid from various precursors and its conjugation with glycine and glucuronic acid have been studied. The method consisted essentially of analysis of the individual urine samples passed by rabbits which had received the compounds under investigation. The metabolites estimated were ether-soluble acid (in some experiments fractionated into benzoic and hippuric acids) and ester glucuronide.

When a compound is administered to a rabbit it may undergo several consecutive processes, e.g.

absorption, conversion of potential centre (see Bray, Ryman & Thorpe, 1948), conjugation and excretion. Preliminary experiments showed that absorption appeared to be virtually complete within about 2.5 hr., so that if measurements were made only on those portions of the excretion curves corresponding to times after this, the effect of absorption could be neglected. For the rest it was assumed that the curves obtained would represent the slowest process of the series. Thus if the amount of the compound administered is the only limiting factor then it would be expected that the reaction being studied would be of the first order.

METHODS

Animals, diet and dosage. Rabbits of not more than 2 kg. weight were used, since larger animals were liable to pass urine samples greater in volume than the capacity of the receivers used (140 ml.). They were maintained on the constant diet of rabbit pellets and water customary in this laboratory (Bray, Ryman & Thorpe, 1947). The compounds were administered by stomach tube as solutions or suspensions in water.

Collection of urine samples. For this purpose a fraction collector was constructed. It consisted of a brass turntable which carried round its periphery forty-eight receiving tubes each of capacity 140 ml. Its movement was weight-activated and controlled by an escapement operated by an electromagnet activated by impulses from a timing clock. In this way the receiver could be changed automatically at predetermined time intervals ranging from 1 min. to 1 hr. The rabbit was housed in a metabolism cage fitted into a tinned-copper collecting funnel of the usual type, the spout of which was connected to a glass tube delivering into the receivers beneath.