3. Some modifications of the technique of estimating amino-acids by paper chromatography were used.

4. The behaviour of glutathione on paper chromatograms is described.

We are indebted to Dr P. György for the sample of brewer's

yeast, to Sir Charles Harington, F.R.S., for a specimen of synthetic glutathione and to Dr R. Consden for thiazolidine carboxylic acid. We acknowledge technical assistance from Mr R. Denman and Mr F. Grover. One of us (O.L.) is in receipt of a Research Grant from the Medical Research Council.

REFERENCES

- Astrup, T. & Øhlenschläger, Y. (1948). *Biochem. J.* **42**, 211.
 Awapara, J. (1948). *Arch. Biochem.* **19**, 172.
 Barton-Wright, E. C. (1949). *Biochim. Biophys. Acta*, **3**, 679.
 Block, R. J. (1948). *Science*, **108**, 608.
 Block, R. J. & Bolling, D. (1945). *Arch. Biochem.* **7**, 313.
 Consden, R., Gordon, A. H. & Martin, A. J. P. (1944). *Biochem. J.* **38**, 224.
 Consden, R., Gordon, A. H. & Martin, A. J. P. (1946). *Biochem. J.* **40**, 580.
 Crumpler, H. R. & Dent, C. E. (1949). *Nature, Lond.*, **164**, 441.
 Dent, C. E. (1947). *Biochem. J.* **41**, 314.
 Dent, C. E. (1948). *Biochem. J.* **43**, 169.
 Dohan, J. S. & Woodward, G. E. (1939). *J. biol. Chem.* **129**, 393.
 Edwards, L. E., Sealock, R. R., O'Donnell, W. W., Partlett, G. R., Barclay, M. B., Tully, R., Tybout, R. H., Box, J. & Murlin, J. R. (1946). *J. Nutrit.* **32**, 597.
 Fisher, R. B., Parsons, D. S. & Morrison, G. A. (1948). *Nature, Lond.*, **161**, 764.
 György, P. (1948). *Trans. 3rd Conf. on Biological Anti-oxidants. Josiah Macy, Jun. Foundation*, p. 71.
 György, P. & Goldblatt, H. (1949). *J. exp. Med.* **89**, 245.
 Himsworth, H. P. & Glynn, L. E. (1944a). *Clin. Sci.* **5**, 93.
 Himsworth, H. P. & Glynn, L. E. (1944b). *Clin. Sci.* **5**, 133.
 Himsworth, H. P. & Lindan, O. (1949). *Nature, Lond.*, **163**, 30.
 Hock, A. & Fink, H. (1943a). *Hoppe-Seyl. Z.* **278**, 136.
 Hock, A. & Fink, H. (1943b). *Hoppe-Seyl. Z.* **279**, 187.
 Landua, A. J. & Awapara, J. (1949). *Science*, **109**, 385.
 Lindan, O. & Work, E. (1951). *Biochem. J.* **48**, 344.
 Lugg, J. W. H. (1938). *Biochem. J.* **32**, 114.
 Martin, A. J. P. (1949). *Ann. Rep. Chem. Soc.* **45**, 267.
 Martin, A. J. P. & Mittelman, R. (1948). *Biochem. J.* **43**, 353.
 Masters, M. (1939). *Biochem. J.* **33**, 1313.
 Matet, A., Matet, J. & Fridenson, O. (1947). *J. Physiol., Paris*, **39**, 381.
 Naftalin, L. (1948). *Nature, Lond.*, **161**, 763.
 Pratt, J. J. & Auclair, J. L. (1948). *Science*, **108**, 213.
 Polson, A. (1948). *Biochim. Biophys. Acta*, **2**, 575.
 Schwarz, K. (1944). *Hoppe-Seyl. Z.* **281**, 101, 109.
 Weichselbaum, T. E. (1935). *Quart. J. exp. Physiol.* **25**, 363.
 Woiwod, A. J. (1948). *Nature, Lond.*, **161**, 169.
 Work, E. (1949). *Biochim. Biophys. Acta*, **3**, 400.
 Work, E. (1950). *Nature, Lond.*, **165**, 74.

The Nutritive Properties of Two Yeasts used to Produce Massive Dietetic Liver Necrosis in Rats

BY O. LINDAN AND ELIZABETH WORK

Medical Unit and Department of Chemical Pathology, University College Hospital Medical School, London, W.C. 1

(Received 7 July 1950)

Dietetic liver necrosis in rats has been produced by feeding diets deficient both in protein and in tocopherol (Himsworth, 1950). Such diets containing suboptimal amounts of casein, soya-bean meal or yeast, produce liver necrosis within periods varying from a few weeks to several months; when extra cystine or methionine is added to the diets the lesion is prevented (Weichselbaum, 1935; Hock & Fink, 1943a; Himsworth & Glynn, 1944b; Matet, Matet & Fridenson, 1947; György & Goldblatt, 1949). A deficiency in the sulphur-containing amino-acids relative to other amino-acids in the diet thus appears to be one of the factors responsible for the development of liver necrosis. This theory is

supported by experiments of Glynn, Himsworth & Neuberger (1945) in which dietetic liver necrosis was produced in rats by substituting the dietary protein by a synthetic mixture containing all the essential amino-acids except methionine and cystine. Hock & Fink (1943b), following an investigation of the necrogenic properties of diets in which the major part of the protein was provided by different types of yeasts, associated the ease of development of liver necrosis with the relative amounts of sulphur and nitrogen in the diet. When the sulphur:nitrogen ratio was 1:30 the incidence of necrosis was high, but there was no necrosis with a ratio of 1:13. It is obvious that the authors thought that this ratio

might be a rough measure of the amounts of the sulphur-containing amino-acids present in the dietary protein.

In this laboratory, diets containing 18 % yeast as the sole source of protein (approx. 7 % protein) have been used to produce liver necrosis (Himsworth & Glynn, 1944*a*), but comparison of the results with those obtained by György & Goldblatt (private communication and 1949) on their yeast diets showed that their incidence and severity of necrosis was much lower. The yeast used in this laboratory was baker's yeast, while György used a brewer's yeast. Since it was probable that the difference in types of yeast accounted for the dissimilar results in the two laboratories, samples of yeasts were exchanged in order to compare their nutritional properties for the same strain of rat and also their chemical composition. György's results showed no great difference between the two yeasts with respect to their total S:N ratios (1:10.5 for our baker's and 1:9.1 for his brewer's yeast), but there was a marked difference in their necrogenic properties; thirteen out of fourteen rats on our baker's yeast diet died of acute liver necrosis, while none of fourteen rats fed György's brewer's yeast developed any lesion (György, 1948).

Determination of the amino-acid compositions of the two yeasts is reported in the preceding paper (Lindan & Work, 1951); this paper describes a comparison of their dietetic properties. The necrogenic properties of yeast diets were compared using young tocopherol-depleted rats and the biological values of the yeast proteins were studied by growth experiments on young normal rats. Following the observation of Leaf & Neuberger (1947) that a deficiency of sulphur-containing amino-acids lowered the level of liver glutathione, the effect of the two yeast diets on liver glutathione was also investigated. A connexion was sought between the nutritional differences shown by the two yeasts and their content of amino-acids, particularly the sulphur-containing amino-acids. No investigations on the tocopherol contents of the yeasts were carried out, since tocopherol is supposed to be absent from all types of yeast (György, 1948, and private communication).

EXPERIMENTAL

General. The yeast used in this laboratory to produce rat liver necrosis was a dried baker's yeast (D.C.L. no. G.B. 161) marketed by Distiller's Co. Ltd. London. The sample provided by Dr P. György, was a dried brewer's yeast (Fleischmann's type 50B) marketed by Standard Brands, Inc., U.S.A. All our nutritional experiments were carried out with the same bulk samples of each yeast; these were also used for the analyses reported in the previous paper (Lindan & Work, 1951).

The composition of the diets (Himsworth & Glynn, 1944*b*) was as follows: dried yeast (baker's or brewer's), 18%; corn

starch, 73%; salt mixture (Glaxo), 3%; olive oil (B.P.), 5%; cod liver oil, 1%. (The dried yeast contained about 6% moisture.)

The olive oil and cod liver oil were mixed daily into the diets, as were the following vitamins, the figures given being the amount allowed for each rat: aneurin, 20 μ g.; riboflavin, 25 μ g.; pyridoxine, 20 μ g.; calcium pantothenate, 100 μ g.

Newly weaned albino rats (wt. 34–46 g.) were used as experimental animals. They were of the same strain (Wistar) as used in other experiments on dietetic liver necrosis in this laboratory (Lindan & Himsworth, 1951) and were kept three to five together in cages with raised wire-mesh bases.

Experiment on necrogenic properties of yeasts. Tocopherol-depleted animals were used for this experiment in order to eliminate the protective action of stored tocopherol against liver necrosis (Himsworth & Lindan, 1949). The young rats and their mothers were fed a diet low in tocopherol (approx. 1.1 mg. tocopherol/100 g. diet) from the tenth day of lactation until weaning at 21 days (Mason & Bryan, 1938; Lindan & Himsworth, 1951). Eighteen such weanling rats from three litters were divided into two groups, the first group (five rats) receiving baker's, the second group (thirteen rats) brewer's yeast diet. The diets were fed *ad lib.* for 64 days; after that the seven surviving rats, which all belonged to the second group, were subdivided into two groups and given either baker's or brewer's yeast diet restricted to 8 g./rat/day. This restriction of food intake was found by Glynn & Himsworth (1944) to be necessary to produce necrosis in rats large enough to consume more than this amount of food when food consumption was unrestricted.

Experiment on growth-promoting properties of yeasts. The weanling rats used were reared on normal stock diet. In order to avoid early deaths from necrosis they were not tocopherol-depleted. Nine litter mates were divided into three groups, three rats to a cage, and the food consumption of each group was recorded daily. Baker's and brewer's yeast diets were fed *ad lib.* to groups 1 and 2 respectively. Because more brewer's yeast diet was consumed than baker's, the third group was fed brewer's yeast diet in amount restricted to that consumed by group 1 receiving baker's yeast. The experiment was continued for 33 days, after which the rats were killed and their livers were analysed in order to determine the influence of diet on liver composition. The values for reduced glutathione (iodometric method of Leaf & Neuberger, 1947), total N and water contents were compared with those of normal rats.

RESULTS

Necrogenic properties of yeasts

Growth curves of eighteen tocopherol-depleted rats fed the diets containing either baker's or brewer's yeast as the sole source of protein are shown in Fig. 1. The occurrence of each death from acute liver necrosis is also indicated.

The five rats fed baker's yeast diet *ad lib.* showed no gain in weight and they all died of acute massive liver necrosis after 24–30 days (mean 26 days) on the diet. The general condition of the animals and the histological picture of their livers corresponded with those found in other experiments on liver necrosis from this laboratory (Glynn & Himsworth, 1944; Lindan & Himsworth, 1951).

The brewer's yeast diet fed *ad lib.* for 64 days supported a fair rate of growth (1.2 g./day) and only six out of thirteen rats died of necrosis between 28 and 45 days (mean 35 days). The rats which died had not shown any difference in growth rate compared with those which survived for this period. The pathological changes in their livers (acute massive necrosis) were similar to those found in the necrotic

Fig. 1. Average growth curves and occurrence of liver necrosis in tocopherol-depleted rats fed yeast diets. —, Baker's yeast diet *ad lib.*; ---, brewer's yeast diet *ad lib.*; - · - ·, baker's yeast diet 8 g./rat/day; · · · ·, brewer's yeast diet 8 g./rat/day; +, death from acute liver necrosis.

rats fed baker's yeast diet, but the animals' general condition was far better. The remaining seven animals continued to grow at the same rate until the 64th day. After that, three rats were continued on brewer's yeast diet, but in amounts reduced to 8 g./rat/day. These animals survived another 96 days in fairly good condition; when killed they showed neither liver necrosis nor post-necrotic scarring. The diet of the other four rats was changed to one containing baker's yeast and their food intake was restricted to 8 g./day; all died of liver necrosis within 34–74 days. Even after restriction of the food intake, the superior nutritive properties of the brewer's yeast were revealed by the growth curves.

Growth-promoting properties of yeasts

The results of feeding yeast diets to rats not depleted of tocopherol are shown in Fig. 2, where mean growth curves and food consumption are recorded. There were no deaths from necrosis during the experimental period of 33 days. During the whole experiment, the rats fed brewer's yeast diet *ad lib.* consumed 48 % more food than did the ones receiving baker's yeast. After about 12 days, the food consumption of both groups decreased, by 12 % in the case of brewer's yeast (group 2), and 31 % in group 1 on baker's yeast. The growth curves

Fig. 2. Effect of yeast diets on growth of weanling rats (mean initial wt. 40 g.). Each curve represents mean values for three rats. —, Group 1, baker's yeast diet *ad lib.*; · · · ·, group 2, brewer's yeast diet *ad lib.*; ---, group 3, brewer's yeast diet restricted to consumption of group 1. Figures against curves indicate average daily food consumption (g./rat) for the periods indicated by arrows.

of groups 1 and 2 fed the two diets *ad lib.* were similar to those of corresponding groups of tocopherol-depleted rats shown in Fig. 1. Group 3, receiving the brewer's yeast diet restricted in amount to that consumed by group 1 on baker's yeast, showed a growth rate of 0.5 g./day which was higher than that of group 1, confirming the higher biological value (weight for weight) of brewer's yeast.

The influence of the diets on the glutathione, total nitrogen and water contents of the livers is shown in Table 1. The livers of all the rats fed the yeast diets showed only half the glutathione content of normal control livers, but there was no difference between

Table 1. *Effect of yeast diets on glutathione, nitrogen and water contents of liver*

(Figures represent mean values for three rats.)

Diets and rats used	Body wt. (g.)	Reduced glutathione		Nitrogen (g./100 g. liver)	Water (g./100 g. liver)
		(mg./100 g. liver)	(mg./100 g. liver N)		
Stock diet:					
Weanlings	38	197	68	2.9	72
33 days after weaning	109	207	66	3.1	71
Yeast diets:					
Baker's <i>ad lib.</i>	45	93	38	2.4	66
Brewer's <i>ad lib.</i>	75	96	35	2.8	66
Brewer's restricted	59	97	41	2.4	67

the three groups of rats on the yeast diets. Both the nitrogen and water contents of the livers of rats fed the yeast diets were slightly lower than those of the controls.

DISCUSSION

It is evident that baker's and brewer's yeasts when used as sources of dietary protein for rats, differ in their nutritive qualities as shown both by their tendency to produce liver necrosis and by their growth-promoting properties. The experiment in which liver necrosis was produced between the 24th and 30th day in the whole of the small group of tocopherol-depleted weanling rats fed diets containing 18% baker's yeast has been confirmed by further data (Lindan & Himsworth, 1951) on 124 rats, all of which developed necrosis within a mean period of 28 days (S.D. ± 10.1). Only half the animals fed brewer's yeast diet developed necrosis, and the time of onset was delayed to between the 28th and 45th day. The relative difference in the necrogenic properties of the two yeasts is similar to the results reported by György (1948), although György found the onset of necrosis with our baker's yeast to be very much later (100th to 144th day, one rat surviving 200 days); his sample of brewer's yeast produced no necrosis within this period. The differences between György's results and ours could be attributed to his using older rats with no pre-experimental depletion of tocopherol. Older rats have been shown to be more resistant to necrosis, and the onset of necrosis may be considerably delayed by feeding diets high in tocopherol in the pre-experimental period (Lindan & Himsworth, 1951). On the other hand, the difference in results might be due to variations in composition of the particular batches of yeast used in the two laboratories; thus the values reported by György for both nitrogen and sulphur differ from our results, particularly in the case of sulphur, which in the baker's yeast is almost double our figure (see Table 5, Lindan & Work, 1951). However, the general conclusion which can be drawn from these two sets of nutritional experiments is that both the baker's yeast samples were much more necrogenic than the brewer's yeast samples.

Comparison of the biological values of the two yeast proteins, judged by growth-promoting properties, agreed with the results of our necrosis experiment. With the same food intake, brewer's yeast as protein produced a slight gain in weight (17 g. in 33 days), while the weights of rats fed the baker's yeast diet remained stationary. With unrestricted feeding appetite was much better in the brewer's yeast group, but food consumption dropped during the experiment, the decrease in the case of baker's yeast diet being more marked. The assumption of Hock & Fink (1943b) that the total sulphur:

nitrogen ratio in low-protein diets bears a relation to the incidence of liver necrosis should be considered in the light of the following comments. As noted (Lindan & Work, 1951, Table 5), analytical figures for different samples of the same strain of yeast are very variable, ranging from 1:10 to 1:21 for our baker's yeast and from 1:9 to 1:15 for György's brewer's yeast, but the baker's yeast was always more necrogenic than the brewer's. The total sulphur: total nitrogen ratio can give little indication of the relative biological values of the yeasts because the figures for neither sulphur nor nitrogen can be directly related to protein content. Variable proportions of total sulphur consist of sulphate sulphur. This sulphate sulphur probably bears a direct relationship to the sulphur content of the original growth media of the yeasts (Fink & Just, 1939), which may vary from batch to batch.

Considering the relationship between organic sulphur and total nitrogen, the ratios for the two yeasts were found to be very similar (1:21 for baker's and 1:19 for brewer's) and so bore no relation to the necrogenic properties of the yeasts. It is interesting to note that these organic sulphur:nitrogen ratios are of the same order as those (1:18 to 1:20) reported by Fink & Hock (1947) for their yeast diets of variable necrogenic properties.

The total nitrogen figures on which the sulphur:nitrogen ratios are based do not reflect the nutritive values of the yeast proteins, since the amount of non-protein, non-amino-acid nitrogen is not considered. The usual estimation of non-protein nitrogen by trichloroacetic acid precipitation has no significance nutritionally, since a considerable proportion of the non-protein nitrogen must be due to amino-acids or peptides. We found that the proportion of total nitrogen of the yeasts soluble in 75% ethanol was 13% for baker's and 18% for brewer's yeast; of this amount about 63% was represented by amino-acid nitrogen, as determined chromatographically (Tables 2 and 3, Lindan & Work, 1951). It appears then that about 5 and 7% of the total nitrogen was extractable non-amino-acid nitrogen in the baker's and brewer's yeast respectively. The amount of ethanol-insoluble non-protein nitrogen, also calculated by chromatography, was about 10 and 6% of the total nitrogen respectively. The total non-protein non-amino-acid nitrogen can therefore be assumed to be about 14% of the total nitrogen of both yeasts. As brewer's yeast contained 8.40% total nitrogen, compared with 7.35% for baker's yeast, we can conclude that, weight for weight, the former yeast had more protein.

Our diets containing 18% of baker's and brewer's yeast were intentionally very poor in protein (about 7 and 8% respectively, calculated as protein nitrogen $\times 6.25$). Table 2 shows the minimal requirements of the essential amino-acids for normal growth in

Table 2. *Essential amino-acid composition of necrogenic yeast diets compared with minimal requirement for normal growth of rats, after Rose (1937)*

(Columns 4-6 represent total average amino-acid intake per rat for 33 days found in the growth experiment for the three groups.)

	Amino-acid, mg. in					
	100 g. diet			123 g. diet		180 g. diet
	Rose's minimal amount for normal growth	Baker's yeast	Brewer's yeast	Baker's yeast (group 1)	Brewer's yeast (group 3)	Brewer's yeast (group 2)
	(1)	(2)	(3)	(4)	(5)	(6)
Arginine	200	190	440	230	540	790
Histidine	400	320	370	390	450	660
Isoleucine	500	350	400	430	490	720
Leucine	800	580	670	710	820	1210
Lysine	1000	520	590	640	730	1060
Methionine	600	110	120	135	150	220
Cystine	—	130	180	160	220	320
Phenylalanine	700	440	170	540	210	310
Threonine	500	430	720	530	880	1300
Tryptophane	200	—	—	—	—	—
Valine	700	730	830	900	1020	1490

rats, as given by Rose (1937), compared with the essential amino-acid composition of the yeast diets used in the present work and based on figures reported (Lindan & Work, 1951; Table 3). Besides a marked deficiency in sulphur-containing amino-acids in both diets, particularly in baker's yeast diet, the table shows that other amino-acids, notably phenylalanine, lysine and leucine were below the minimum figures suggested by Rose. Although addition of extra cystine or methionine to yeast diets is known to afford complete protection against liver necrosis, we do not know whether the imbalance among the other amino-acids is also playing a role in the production of necrosis. The total average consumption per rat of each essential amino-acid during the whole period of the growth experiment (Fig. 2) is also given in Table 2. All the essential amino-acids except phenylalanine were consumed in larger amounts by the rats given brewer's yeast diet either *ad lib.* or in amounts restricted to the consumption of the baker's yeast diet. This could easily explain the higher growth rate observed with the brewer's yeast diet. Supplementation of baker's yeast diet with cystine up to 0.63 g./100 g. of diet did not improve growth of young rats, although it protected the animals against liver necrosis (Lindan & Hims-worth, 1951). The failure of baker's yeast diet to support growth could therefore be due to multiple amino-acid deficiencies.

The two yeasts showed a difference in their contents of ethanol-extractable amino-acids, 0.44 g. of amino-acid nitrogen being extractable from 100 g. of baker's yeast and 0.92 g. from 100 g. of brewer's yeast. The amino-acid make-up of the extracts differed in the case of only a few amino-acids. The

nutritional significance of these extractable amino-acids was not assessed in our experiments, but the higher total glutathione content of brewer's yeast might be of importance.

The brewer's yeast was a non-fermenting sample, while the baker's yeast was capable of fermentation. Experiments (Hims-worth & Lindan, unpublished observations) with autoclaved (20 min. at 115°) non-fermenting baker's yeast showed that it had not only retained, but even increased its necrogenic properties; twelve animals died within a mean time of 16 days from the beginning of the experiment (s.d. ± 2.9). Thus the difference in the necrogenic properties of the two yeasts was not related to their viability.

Since the liver seems to be the organ which is most affected by the deficiency of cystine and methionine, it was of interest to investigate the effect of yeast diets on the distribution of sulphur in the liver. Lee & Lewis (1934) found that the liver protein composition was normal in rats fed cystine-deficient diets, while Dent (1947) reported no change in the cystine and methionine contents of the liver proteins even when the rats had developed necrosis on baker's yeast diet. The only fraction of liver sulphur which varies appears to be the non-protein fraction and may be attributed to variation in glutathione (Leaf & Neuberger, 1947). Despite differences in the necrogenic and growth-promoting properties of the baker's and brewer's yeast, and in their cystine content, we found that they both produced a lowering of the liver glutathione content from 200 to 95 mg./100 g. liver when fed for 33 days to rats which were not tocopherol-depleted during the lactation period. We have found (unpublished work)

that necrotic livers from tocopherol-depleted rats fed baker's yeast diet had an even lower glutathione content of about 25 mg./100 g. liver.

In conclusion, the nutritive properties of baker's and brewer's yeasts differ in respect to both growth-promoting and necrogenic properties. The amino-acid contents of the two yeasts could explain the difference in growth-promoting properties, the diet containing baker's yeast being the more deficient in several amino-acids. We cannot, however, say whether the small differences in the sulphur-containing amino-acid content account entirely for their different necrogenic properties. Further experiments are needed to establish this point.

SUMMARY

1. A comparison was made between the dietetic properties of British baker's yeast and American brewer's yeast, both of which were used as the sole source of protein in the production of dietetic liver necrosis in rats. Diets containing 7-8% of yeast protein were used.

2. When fed to tocopherol-depleted weanling rats the baker's yeast diet proved to be more necrogenic than the brewer's yeast diet. Growth experiments on normal weanlings also showed the inferior biological value of baker's yeast.

3. Rats fed the yeast diets for 33 days had liver glutathione contents which were 50% lower than those of normal rats. There was no difference between the composition of livers of rats fed either type of yeast.

4. Different dietetic deficiencies exhibited by the two yeasts are discussed in relation to their contents of sulphur, nitrogen and amino-acids. Brewer's yeast contains 38% more cystine and glutathione than baker's yeast. Comparison with Rose's (1937) figures for minimum essential amino-acid requirements for normal growth showed that both yeast diets were also deficient in other essential amino-acids.

We are indebted to Dr P. György for the sample of brewer's yeast. We acknowledge technical assistance from Miss A. Moxham and Miss D. Rixom. One of us (O.L.) is in receipt of a Research Grant from the Medical Research Council.

REFERENCES

- Dent, C. E. (1947). *Biochem. J.* **41**, 314.
 Fink, H. & Hock, A. (1947). *Z. Naturforsch.* **2b**, 187.
 Fink, H. & Just, F. (1939). *Biochem. Z.* **303**, 234.
 Glynn, L. E. & Himsworth, H. P. (1944). *J. Path. Bact.* **56**, 297.
 Glynn, L. E., Himsworth, H. P. & Neuberger, A. (1945). *Brit. J. exp. Path.* **26**, 326.
 György, P. (1948). *Biological Antioxidants. Trans. 3rd Conf. Josiah Macy, Jun. Foundation*, p. 71.
 György, P. & Goldblatt, H. (1949). *J. exp. Med.* **89**, 245.
 Himsworth, H. P. (1950). *The Liver and its Diseases*, 2nd ed. Oxford: Blackwell.
 Himsworth, H. P. & Glynn, L. E. (1944a). *Clin. Sci.* **5**, 93.
 Himsworth, H. P. & Glynn, L. E. (1944b). *Clin. Sci.* **5**, 133.
 Himsworth, H. P. & Lindan, O. (1949). *Nature, Lond.*, **163**, 30.
 Hock, A. & Fink, H. (1943a). *Hoppe-Seyl. Z.* **278**, 136.
 Hock, A. & Fink, H. (1943b). *Hoppe-Seyl. Z.* **279**, 187.
 Leaf, G. & Neuberger, A. (1947). *Biochem. J.* **41**, 280.
 Lee, W. C. & Lewis, H. B. (1934). *J. biol. Chem.* **107**, 649.
 Lindan, O. & Himsworth, H. P. (1951). *Brit. J. exp. Path.* **31**, 651.
 Lindan, O. & Work, E. (1951). *Biochem. J.* **48**, 337.
 Mason, K. E. & Bryan, W. L. (1938). *Biochem. J.* **32**, 1785.
 Matet, A., Matet, J. & Fridenson, O. (1947). *J. Physiologie*, **39**, 381.
 Rose, W. C. (1937). *Science*, **86**, 298.
 Weichselbaum, T. E. (1935). *Quart. J. exp. Physiol.* **25**, 363.

The Use of 'CO₂ Buffers' in Manometric Measurements of Cell Metabolism

By H. A. KREBS

Medical Research Council Unit for Research in Cell Metabolism, Department of Biochemistry, University of Sheffield

(Received 22 July 1950)

Pardee (1949) has recently devised a simple manometric method for the measurement of the oxygen absorption of cells respiring in the presence of carbon dioxide. The method is generally applicable to systems where carbon dioxide and a second gas are formed or used and where the pressure changes of the second gas are to be measured. It is based on the use, in a centre well of the manometric vessel, of a

'CO₂ buffer' capable of maintaining a virtually constant pressure of carbon dioxide in the gas phase. A suitable 'CO₂ buffer' is an aqueous solution of diethanolamine which binds carbon dioxide reversibly, mainly according to the equation