Physicochemical Investigations on Two Unusual Sialic Acid-Rich Mucoids from the Intestinal Tract of Patients with Fibrocystic Disease of the Pancreas

By PATRICIA G. JOHANSEN

Gastroenterological Research Unit, Royal Children's Hospital Research Foundation, Melbourne, N. 3, Australia

(Received 28 May 1962)

Fibrocystic disease of the pancreas in human beings is an hereditary condition, characterized by the excessive viscosity of mucous secretions throughout the body. Elevated concentrations of Na^+ and Cl^- ions are present in the sweat, tears and saliva, and there is a deficiency of pancreatic enzymes and an exceptional capacity for *Staphylococcus aureus* to thrive in the bronchial tree. Despite intensive investigations, the primary underlying metabolic defect has not yet been identified. It is not known whether the excessive viscosity of the secretions is due to chemical differences in the mucoid constituents, to different relative proportions of chemically 'normal' mucosubstances, or to a more basic disturbance in, e.g., the ion balance of the cells and tissues which form mucus. The present paper describes the examination of two unusual samples of mucus obtained from two children suffering from this disease.

MATERIALS AND METHODS

Materials

L-Fucose, D-mannose, D-galactosamine hydrochloride and D-glucosamine hydrochloride were chromatographically pure samples, obtained from the Mann Research Laboratories, New York, U.S.A. Glucose-free D-galactose and glucuronic acid were obtained from the Sigma Chemical Co., St Louis, Mo., U.S.A. Authentic crystalline N-acetylneuraminic acid was kindly supplied by Mr B. Graham of the Commonwealth Serum Laboratories, Melbourne. Crystalline salt-free trypsin, α -chymotrypsin, pepsin, papain and egg-white lysozyme were obtained from the Sigma Chemical Co. Crude papain was obtained from E. Merck A.-G., Darmstadt, Germany. Crystalline vibriocholerae neuraminidase (RDE) was kindly provided by Dr G. L. Ada, The Walter and Eliza Hall Institute, Melbourne. All other chemicals were the best available commercially. Organic liquids were redistilled where necessary. Cellulose casings for dialysis were obtained from Visking Corporation, Chicago, U.S.A.

Mucoid samples. Sample 1 was obtained from a boy 2 years old, after a large bowel washout; it is believed to have formed in the terminal small bowel during a temporary blockage due to an ileo-colic intussusception. Sample 2 was passed *per rectum* by a baby 7 days old, 24 hr. after a successful operation for meconium ileus; it

probably arose in the large bowel. Both samples were obtained as firm plugs of ropey mucus and were relatively uncontaminated by extraneous material such as faeces, urine or blood. Clinical investigations indicated that probably neither had been lodged in the gut for more than 24 hr.

Preparation of samples for examination. After preliminary washings with cold 0.9% NaCl to remove any loose material, the mucus plugs were washed with cold 0.9% NaCl containing merthiolate (0.001%) for several days at 2°, the saline solution being changed every day. After several further washings in merthiolate-free 0.9% NaCl, followed by cold water, the material was dehydrated in redistilled ethanol at 2°, dried *in vacuo* over P_2O_5 and anhydrous CaCl_2 , ground to a fine powder in an all-glass pestle and mortar and finally stored *in vacuo* over P_2O_5 .

Analytical procedures

Solutions of the mucus samples were prepared for analysis and examination by dissolving the dried materials in 0.1N-NaOH at room temperature, and subsequent neutralization and dilution of the solutions. This treatment was necessary because of the insolubility of the original native mucus or the dried materials in water, aqueous buffers or other suitable solvents.

Unless otherwise stated, all buffer solutions were prepared according to the directions of Gomori (1955). A Beckman model DU spectrophotometer was used for reading the absorption in all colorimetric estimations. At least two sets of duplicate estimations were carried out on separately weighed portions of each sample for every analysis.

Determination of nitrogen. Nitrogen was determined by a microdiffusion Kjeldahl method (Conway, 1957; Johansen, 1958).

Determination of fucose. Fucose was determined by the method of Dische & Shettles (1948) as modified by Gibbons (1955), with a 10 min. heating period. Fucose standards (5–50 μg .) containing appropriate concentrations of the other carbohydrate constituents were included in each determination. Extinctions were read at 396 and 427 m μ and the differences in the readings plotted against the fucose content.

Determination of hexosamine. Total hexosamine was determined by the method of Cessi & Piliego (1960), as modified by Johansen, Marshall & Neuberger (1960), after liberation of the amino sugar from the mucus samples by hydrolysis in 4N-HCl at 100° for 8 hr. in sealed ampoules; under these conditions destruction of hexosamine is

probably negligible (Johansen *et al.* 1960). In this method, galactosamine gives about 97% of the colour yield of an equal weight of glucosamine.

Determination of total neutral sugar and galactose. Total neutral sugar was determined by the orcinol- H_2SO_4 method as described by Winzler (1955), the absorption being read at 510 $\text{m}\mu$. Fucose and galactose standards (30–120 μg .) were included in each series of estimations. At 510 $\text{m}\mu$, fucose gives about 65% and mannose about 90% of the extinction value of an equal weight of galactose. The galactose content was determined by subtracting the value for the extinction at 510 $\text{m}\mu$ due to fucose from the total value. Fucose was determined independently by the Dische-Gibbons method (see above). This procedure is probably not entirely valid, since it is not possible to determine the extent of interaction of the various constituents of protein-carbohydrate complexes in the presence of strong sulphuric acid (Johansen, 1958). The individual sugars were also determined by a modification of the quantitative paper-chromatographic method of Wilson (1959), after hydrolysis of the samples in $\text{N-H}_2\text{SO}_4$ at 100° for 1–2 hr.

Determination of sialic acid. Total sialic acid was determined as *N*-acetylneuraminic acid by the resorcinol method (Svennerholm, 1957, 1958); the extinction was read at 450 and 580 $\text{m}\mu$ to eliminate interference due to other sugar constituents. Free sialic acid was determined by the thiobarbituric acid method (Warren, 1959).

Determination of uronic acid. The carbazole method of Dische (1950) was used.

Amino acid analysis. The mucoids were hydrolysed in constant-boiling HCl at 115° for 18 hr., and the amino acids in the hydrolysate were separated and determined on two ion-exchange columns according to the method of Moore & Stein (1954a, b).

Paper chromatography. The one-dimensional descending technique was used, with the following solvents and spray reagents: (i) for neutral sugars, the solvent was butan-1-ol-pyridine-water (9:5:8, by vol.) (Glegg & Eidinger, 1954) and the spray reagent aniline hydrogen phthalate (Partridge, 1949); (ii) for sialic acids, the solvent was butan-1-ol-propan-1-ol-0.1 N-HCl (1:2:1, by vol.) and the spray resorcinol reagent (Svennerholm & Svennerholm, 1958).

Determination of ash content. The ash content of sample 1 was determined. A portion of the undried mucus was washed exhaustively with 0.001% merthiolate in 0.9% NaCl ; it was dialysed against glass-distilled water at 2° for several days, then against 0.5% EDTA , adjusted to pH 6.0 with NaOH , for 2 days, and finally against several changes of glass-distilled water. Half of the material was dehydrated in ethanol, the remainder was freeze-dried, and the finely powdered samples were ignited to constant weight in an electric furnace at 600°. One sample was dissolved in hot conc. HCl , and the calcium content was estimated by titration against EDTA by a modification of the micro-method of Diehl & Ellingboe (1956), with a calcein-thymolphthalein indicator. The solution was also examined for the presence of sodium and potassium by flame photometry.

Determination of inhibition of viral haemagglutination. The virus-inhibition titre was tested against two types of influenza-virus strains: type A (MEL) and type B (LEE). Indicator viruses were prepared by the heat-inactivation method of Stone (1949). Serial twofold dilutions of ap-

proximately 0.1% solutions of the two mucus samples were made up in 0.9% NaCl . Samples (0.25 ml.) of each dilution were mixed with 0.025 ml. of the indicator virus containing the equivalent of four haemagglutinating doses. The mixtures were allowed to stand for 20 min. at 20° before the addition of 0.025 ml. of a 5% (v/v) suspension of fowl red cells. These were allowed to settle at 20° for 30 min. and the degree of inhibition was read by the pattern test (Stone, 1949).

Treatment with enzymes. Portions of the undried mucus samples were incubated under suitable conditions with the following enzymes (in each case, about 0.02 g. of the undried material was added to 1 ml. of the digestion medium): trypsin plus chymotrypsin (1.25 mg. of each enzyme/ml. of 0.05 M -phosphate buffer, pH 7 and 8; 37°); pepsin (0.1 mg./ml. of 0.05 M - KCl-HCl buffer, pH 1.8; 37°); lysozyme (0.2 mg./ml. of 0.06 M -phosphate buffer, pH 6.2; 25°); crystalline and crude papain (0.01 and 0.2 mg. respectively/ml. of 0.04 M -citrate buffer, pH 5.5, 4 mm with respect to cysteine and 1 mm with respect to EDTA ; 37°). The papain was suitably activated with cysteine- EDTA , and the crude enzyme suspension was also filtered, immediately before use. The 24-hr. papain digests were dialysed exhaustively against water at 2°, and the diffusate and solution of non-diffusible-material were freeze-dried and analysed for fucose and sialic acid. Samples of the diffusate were also hydrolysed (5.7 N-HCl for 18 hr. at 115°) and the liberated α -amino groups plus amide ammonia were determined by the photometric ninhydrin method of Moore & Stein (1954b). Toluene was added to all digests as a preservative.

The action of neuraminidase on the mucoids was also investigated. Solutions of the samples of dried mucus were dialysed against 0.05 M -tris-maleate buffer, pH 5.5, 1 mm with respect to CaCl_2 . Samples were removed for the determination of total *N*-acetylneuraminic acid by the resorcinol method (Svennerholm, 1957, 1958), and crystalline neuraminidase dissolved in the same buffer was added to the remaining solutions, giving a final concentration of 5000 units of enzyme/mg. dry wt. of mucoid in 2 ml. of buffer (Ada, French & Lind, 1961). The solutions were incubated at 37° and the liberated *N*-acetylneuraminic acid was determined at intervals by the thiobarbituric acid method (Warren, 1959).

Physical examination

Absorption spectra. The ultraviolet-absorption spectra were measured in 0.1 N-NaOH in a 1 cm. cell.

Electrophoresis. Both samples were examined by the microimmunoelectrophoretic method of Scheidegger (1955) in 1.5% agar (Oxoid Ion Agar no. 2), with a veronal buffer (pH 8.35, 1 0.025); the separated samples were stained for protein with (0.01%) nigrosine in aq. 10% (v/v) glycerol-2% (v/v) acetic acid and for glycoprotein (Grabar, 1959), or diffused against a potent goat antiserum to whole human serum. Electrophoresis was also carried out on cellulose acetate strips (Oxoid Ltd., London) in 0.05 M -borate buffer, pH 9.3, by using Kohn's (1958) micromethod; strips were stained for protein with 0.005% nigrosine in aq. 2% (v/v) acetic acid and for glycoprotein (Aronsson, Grönwall & Lausung, 1958).

Ultracentrifuging. Sample 1 was examined in the Spinco analytical ultracentrifuge (model E) in 0.05 M -sodium

Fig. 1. Mucus plug from the lower small intestine of a boy 2 years old suffering from fibrocystic disease of the pancreas (sample 1). The material had been washed for 2 days in 0.001% merthiolate in 0.9% NaCl at 2°. It was readily separated into tough ropes and fibres of opaque gelatinous material.

Fig. 2. Sedimentation diagrams of fibrocystic intestinal mucoid sample 1 (upper pattern) and of human α_1 -glycoprotein (lower pattern) 104 min. after reaching the maximum speed of 42 040 rev./min. The concentrations of mucoid and of protein were approx. 1% (w/v) in borate buffer, pH 8.6, I 0.14. The temperature was 20°.

borate-boric acid buffer, pH 8.6, I 0.14, and in 0.05M-veronal-HCl, pH 7.3, I 0.08, at 20°. The dried material was dissolved in 0.1N-NaOH, neutralized to pH 8.6 with boric acid, and diluted to give a 1% (w/v) solution in 0.05M-borate, pH 8.6. After examination at this pH, the solution was dialysed against several changes of the veronal buffer and examined at the lower pH.

A sample of human α_1 -glycoprotein 430VI-1b was treated in an identical fashion with 0.1N-NaOH, neutralized with boric acid and examined in the ultracentrifuge at the same time, to ascertain whether the alkaline treatment had any marked detrimental effect on a well-characterized mucoid of fairly similar composition.

EXPERIMENTAL AND RESULTS

General properties of the mucus samples. The mucoids were obtained as firm mucus plugs, far more rigid in consistency than any similar material from other human sources, including cervical mucus of pregnancy. They were readily separated into coarse opaque ropes of gelatinous material of remarkable tensile strength (Plate 1: Fig. 1), and retained this form when washed with cold 0.9% sodium chloride solution, but when transferred to water they swelled rapidly and the mucus strands became very fragile.

Apart from admixed serum proteins, both the native gels and the dried materials appeared to be completely insoluble at room temperatures in a variety of solvent systems, including water, urea (3, 6 and 9M, for 6 days), formamide, saturated calcium chloride, saturated calcium chloride-ethanol (9:1, v/v) (Gibbons, 1959), 0.1M-cysteine, various aqueous buffers and salt solutions (including M-potassium iodide, M-potassium thiocyanate and 2mM-sodium chloride), and in dilute mineral acids. The native gels swelled considerably in urea and lost their cohesive properties, but they retained their compact stranded form in the other solvents. However, both the dried and the native mucoids dissolved in 0.1N-sodium hydroxide within a few hours at room temperature.

The mucoids initially contained considerable amounts of serum proteins, but after prolonged washing with 0.9% sodium chloride solution none could be detected by immunoelectrophoresis.

Physical properties. The purified mucoids migrated towards the anode on electrophoresis at pH 8.35 and at pH 9.3. This examination was not entirely satisfactory as the materials formed slightly elongated streaks on the supporting media used (agar gel and cellulose acetate strips), presumably due to adsorptive effects. However, protein and glycoprotein stains gave patterns of corresponding size and intensity with each of the samples.

The ultraviolet-absorption spectra showed no specific absorption peaks between 220 and 350 $m\mu$

apart from a small inflexion in the region of 280 $m\mu$ and again at 250 $m\mu$ in sample 2; the absorption increased rapidly below 260 $m\mu$.

On examination in the ultracentrifuge at pH 7.3 and 8.6, sample 1 was seen as a single main component which formed a hypersharp peak (Plate 1: Fig. 2), similar to that observed with other macromolecular materials, e.g. ovine submaxillary mucin (Gottschalk & McKenzie, 1961). The material appeared to be essentially homogeneous, although somewhat polydisperse. The rate of sedimentation was similar to that of the human α_1 -glycoprotein, which was examined simultaneously in the pH 8.6 buffer; however, the latter gave a much broader peak, indicating its less viscous nature (Plate 1: Fig. 2). Sample 1 gave a $S_{20,w}$ of 5.1s.

Chemical properties. Analytical results for the principal constituents of the two mucoids are given in Table 1. Both samples contained galactose, fucose, sialic acid and hexosamine, as well as amino acids. After hydrolysis in N-sulphuric acid for 1 or 2 hr. at 100°, fucose and galactose were identified in both mucoids by paper chromatography. Extensive formation of humin occurred under these conditions and it is not surprising that the recoveries of the individual sugars, determined for sample 1 by quantitative paper chromatography (Wilson, 1959), were low (see Table 2). Sample 1 also contained traces of mannose, but no other neutral sugars were detected in sample 2 even after hydrolysis under more vigorous conditions (2N-sulphuric acid for 3 hr. at 100°). Glucosamine was the only amino sugar identified in the samples hydrolysed in 5.7N-hydrochloric acid for 18 hr. at 115° and chromatographed on the Moore & Stein (1954a) columns. Neither mucoid contained detectable amounts of uronic acid (less than 0.1%) and there was no evidence of any DNA or RNA.

The sialic acid was completely split from both mucus samples by crystalline neuraminidase at 37° in less than 24 hr. (Fig. 3a) and by 0.1N-sulphuric acid at 80° in less than 1.5 hr. (Fig. 3b). N-Acetylneuraminic acid was the only free sugar that could be detected by paper chromatography in the diffusate of the acid hydrolysate, although quantitative analysis showed that a considerable amount of fucose was also rendered diffusible by this acid treatment (Table 3). In both cases sample 1 was degraded more rapidly than sample 2. The sialic acid was estimated as N-acetylneuraminic acid, although it may be present in the native mucoid as the NO-diacetyl derivative; however, under the alkaline conditions used to prepare the mucoid solutions for examination, it is very probable that the extremely labile O-acetyl groups will be split off.

Virus-inhibitor activity of the mucoids. The titres of the influenza-virus-haemagglutination inhibition

Table 1. *Analytical results for washed preparations of the fibrocystic intestinal mucoids*

The values are uncorrected for ash and are quoted for the dry wt. of mucoid. The values for the amino acids are not corrected for losses during hydrolysis (constant-boiling HCl for 18 hr. at 115°). Details of the analytical methods are given in the text.

	In sample 1		In sample 2	
	g./100 g.	μmole/mg.	g./100 g.	μmole/mg.
Fucose	6.4	0.40	1.8	0.12
Sialic acid	12.6	0.40	10.2	0.33
Hexosamine	16.0	0.90	10.4	0.58
Galactose	9.0	0.50	6.2	0.34
Total nitrogen	7.9	—	10.1	—
Ash	4.2	—	—	—
Amino acid constituents				
Threonine	5.47	0.44	4.32	0.36
Phenylalanine	5.45	0.33	4.79	0.29
Glutamic acid	4.41	0.30	5.69	0.39
Proline	3.34	0.29	2.94	0.26
Leucine	3.41	0.26	5.06	0.39
Aspartic acid	3.46	0.26	4.20	0.32
Glycine	1.80	0.24	2.56	0.34
Valine	2.69	0.23	2.96	0.25
Serine	2.31	0.22	2.50	0.24
Alanine	1.69	0.19	2.41	0.27
Isoleucine	1.97	0.15	2.44	0.19
Lysine	1.90	0.13	2.99	0.21
Arginine	1.76	0.10	3.67	0.21
Tyrosine	1.81	0.10	2.21	0.12
Cystine (half)	0.89	0.07	0.97	0.10
Histidine	1.09	0.07	1.54	0.10
Methionine sulphoxide	0.26	0.02	Trace	Trace
Methionine	0.15	0.01	1.12	0.08
Alloisoleucine	0.03	0.002	—	0.01

Table 2. *Neutral-sugar content of fibrocystic intestinal mucoid sample 1*

The values for the unhydrolysed material were obtained by the direct colorimetric methods for hexose and fucose as described in the text. Acid hydrolysates of the same material (1 mg. of mucoid/ml. of $N-H_2SO_4$ for 1 or 2 hr. at 100°) were separated by paper chromatography with butan-1-ol-pyridine-water (9:5:8, by vol.) as solvent (Glegg & Eidinger, 1954) and analysed by the method of Wilson (1959). The values are expressed as g./100 g. dry wt. of mucoid.

	Unhydrolysed material	Acid hydrolysates	
		1 hr. of hydrolysis	2 hr. of hydrolysis
Galactose	9.0	6.5	5.9
Mannose		Negligible	0.8
Fucose		5.7	4.2

by samples 1 and 2, with type A (MEL) and type B (LEE) strains of influenza virus as indicators, are given in Table 4. Values for bovine and ovine submaxillary mucin, whose activities were determined by Graham & Gottschalk (1960) by the same method, are included for comparison. Both mucoids inhibited the strains of influenza virus selected, the potency of sample 1 being considerably greater than that of sample 2. Compared with

bovine submaxillary mucin, sample 1 was equally potent when measured against an influenza virus A strain, and about ten times as active against the same B strain. Sample 1 was also tested against active influenza virus A (MEL), but there was no inhibition of the active virus. Pusztai & Morgan (1961) have also found a sialic acid-rich mucopolysaccharide isolated from ovarian-cyst fluid to be a potent inhibitor of the haemagglutination of heated type A and type B influenza viruses, but to cause no inhibition of the active viruses.

Action of proteolytic enzymes and lysozyme on the mucus samples. Trypsin and chymotrypsin had no visible action on the native mucoids at pH 8, and at pH 7 no digestion was apparent until after 48 hr. Pepsin also attacked the materials relatively slowly, marked dispersion being observed only after 24 hr. at pH 1.8 and at 37°, conditions under which a portion of the sialic acid may be split off by acid hydrolysis. Lysozyme had no observable effect on the samples at all. However, both crude and crystalline papain attacked the native materials in a dramatic fashion. The crystalline enzyme completely dispersed sample 1 in 3 hr. and sample 2 in 12 hr.; after 24 hr. none of the carbohydrate was diffusible, although a considerable proportion of the amino acid nitrogen (approx. 75

Table 3. *Acid hydrolysis of fibrocystic intestinal mucoid sample 1*

Sample 1 was hydrolysed in 0.1 N-H₂SO₄ for 1.5 hr. at 80° (0.5 mg. of mucoid/ml.). The hydrolysate was exhaustively dialysed against water and the distribution of fucose and *N*-acetylneuraminic acid in the diffusate and non-diffusible fractions determined by specific colorimetric methods. Details of the analytical methods are given in the text.

	Unhydrolysed material (g./100 g. dry wt. of mucoid)	Acid hydrolysate			
		Non-diffusible material		Diffusible material	
		g./100 g.	Recovery (%)	g./100 g.	Recovery (%)
Fucose	6.4	4.3	67	2.0	31
Total <i>N</i> -acetylneuraminic acid	12.6	0.9	7.1	9.9	79

Table 4. *Inhibition of the haemagglutination of influenza virus by the fibrocystic intestinal mucoids*

Experimental details are given in the text. The values for the activity of ovine and bovine submaxillary mucins, determined by Graham & Gottschalk (1960) by the same method, are included for comparison. The values are expressed as the minimum amount (in µg.) giving inhibition.

	Influenza virus A (MEL-i)	Influenza virus A (PR 8-i)	Influenza virus B (LEE-i)
Sample 1	0.02 _a	—	0.18
Sample 2	0.29	—	0.77
Bovine submaxillary mucin	—	0.021	2.00
Ovine submaxillary mucin	—	0.0009	0.0077

Fig. 3. Rate of release of sialic acid from fibrocystic intestinal mucoids. (a) Enzymic removal by crystalline neuraminidase. The incubating medium was 0.05 M-tris-maleate buffer, pH 5.5, 1 mM with respect to CaCl₂; 5000 units of enzyme/mg. dry wt. of mucoid in 2 ml. of buffer were used. The temperature was 37°. (b) Acid hydrolysis. The hydrolysis was carried out in 0.1 N-H₂SO₄ at 80°; 0.5 mg. dry wt. of mucoid/ml. was used. The total sialic acid was estimated as *N*-acetylneuraminic acid by the resorcinol method. Released sialic acid was estimated as *N*-acetylneuraminic acid by the thiobarbituric acid method. ▼, Sample 1; ●, sample 2.

and 95% for samples 1 and 2 respectively) was recovered in the diffusate of the enzymic digests. However, after the incubation of sample 1 with crude papain under similar conditions, all the fucose and sialic acid became diffusible, which suggests that the crude preparation may contain carbohydrate-splitting enzymes which can attack the polysaccharide chains in this mucoid.

DISCUSSION

The homogeneity of the washed preparations of these two fibrocystic intestinal mucoids could not be examined critically because of the small amount of material available; however, the behaviour of sample 1 in the ultracentrifuge indicates that it was probably homogeneous, although somewhat polydisperse. Further, no contaminating serum proteins could be detected in either of the purified samples by immunological methods.

The native mucoids showed many similarities, notably in their lack of solubility in a wide variety of solvents and in their resistance to attack by various enzymes other than papain. A remarkable feature was the tendency of the materials to swell in water, when the gelatinous rope-like masses became very fragile; however, when suspended in salt solutions of a wide range of concentrations they retained their compact stranded appearance. This behaviour suggested that electrolytes may play a significant role in maintaining the original form of the mucus plugs; subsequently a thoroughly dialysed preparation of sample 1 was found to

contain 4.2 g. of total ash/100 g. dry wt. of mucoid, 0.24 g. of which was calcium; sodium was the principal cationic constituent and there was little or no potassium present. There is very little comparable information available in the literature dealing with mucosubstances. Gibbons (1959) has reported values of 3.1 and 3.7 g./100 g. dry wt. of mucoid for the ash content of bovine-oestrus and bovine-pregnancy cervical mucoids, which appear to resemble sample 1 in many of their properties; Odin (1959) found that, in washed preparations of pseudomyxomatous gels, only about 1% of the dry weight was ash. Chernick (1961) and Chernick, Barbero & Parkins (1961) reported an increase in the calcium:sodium ratio in certain fibrocystic mucous secretions, and the molar ratio of these ions in sample 1 is much the same as the value quoted for fibrocystic submaxillary mucin, namely approx. 1:10 (Chernick *et al.* 1961). However, the relationship between the ionic composition and the abnormal properties of mucous secretions in fibrocystic disease is not understood. The behaviour of the mucoids in urea suggests that hydrogen bonds are probably also important in maintaining the structure of the native materials.

These two samples of fibrocystic intestinal mucus resemble typical mucoids found in mammalian epithelial secretions, and according to current terminology (Bettelheim-Jevons, 1958; Pusztai & Morgan, 1961) sample 1 may be described as a 'sialic acid-rich fucomucin'; sample 2, whose composition is similar to that of urinary mucosubstances and pseudomyxomatous gels, may be classified as a 'sialomucin'. Both samples contained galactose, fucose, sialic acid and hexosamine, as well as amino acids, and the major chemical difference between the two preparations was the content of fucose relative to the other carbohydrate constituents. The sialic acid:fucose ratio was 1 and 3 for samples 1 and 2 respectively although the relative proportions of the other sugars were similar in both preparations (Table 5). Sialic acid appears to occupy a terminal position in the carbohydrate chains in both mucoids, since it was all rapidly split off by the neuraminidase and by dilute acid; further, both samples were potent inhibitors of influenza-virus haemagglutination, a reaction in which the essential functional groups are believed to be terminal sialic acid residues (Pusztai & Morgan, 1961).

The distribution of amino acids was fairly similar in both samples and somewhat resembled that found in other epithelial mucosubstances, but there were marked differences in the relative proportions of the amino acids. The most striking feature was the relatively large phenylalanine content of the fibrocystic mucoids, since this amino acid is usually low or absent in comparable mucosubstances. The relative proportions of the basic amino acids were also significantly higher in both samples (Table 6).

Apart from the terminal location of the sialic acid residues in the carbohydrate chains, no conclusions may be drawn about the detailed structure of these two mucoids; however, it seems probable that they consist of relatively large polysaccharide chains associated with polypeptide, since none of the carbohydrate but a high proportion of the con-

Table 5. *Analysis of fibrocystic intestinal mucoids: the molar proportions and molar ratios of the carbohydrate constituents of mucus samples 1 and 2 (cf. Table 1)*

Sialic acid is expressed as *N*-acetylneuraminic acid, and hexosamine as glucosamine base.

	Sample 1	Sample 2
Molar proportions		
Fucose	4	1
Sialic acid	4	3
Hexosamine	9	5
Galactose	5	3
Molar ratios		
Hexosamine:sialic acid	2.2	1.8
Galactose:sialic acid	1.2	1.0
Sialic acid:fucose	1	3
Hexosamine:galactose	1.8	1.7

Table 6. *Distribution of amino acids in epithelial mucoids*

Values quoted as μ mole/mg. dry wt. of mucoid. The values for the bovine mucoids were reported by Gibbons (1959), those for the human salivary substance by Carsten & Kabat (1956) and those for the sialomucopolysaccharide by Pusztai & Morgan (1961).

	Fibrocystic intestinal mucoids		Bovine cervical mucoids		Blood-group substances	
					Human salivary substance	Sialomucopolysaccharide
	Sample 1	Sample 2	Oestrus	Pregnancy		
Threonine + serine	0.66	0.60	0.84	0.69	0.74	0.57
Glutamic + aspartic acid	0.56	0.71	0.32	0.17	0.15	0.10
Proline	0.29	0.26	0.35	0.21	0.18	0.18
Arginine + lysine + histidine	0.30	0.52	0.13	0.15	0.11	0.06
Phenylalanine	0.33	0.29	0.07	0.07	0	0.02

stituent amino acids was diffusible after the material had been digested with crystalline papain.

Histological studies indicate that a variety of distinctive types of mucus-forming cells may be present and active in certain mucous epithelia, whose secretions may therefore contain a mixture of mucosubstances (Spicer, 1960). The physico-chemical properties of these secretions are probably determined by several factors, in particular the chemical composition of the individual mucoids and their relative proportions in the secretion, the total concentration of mucosubstances and the electrolyte content; other organic constituents, such as serum proteins, which probably arise by transudation of plasma through the epithelial wall, may also be of importance.

It has been suggested that the external environment, especially the ionic content of the extracellular fluid, may greatly influence the development and differentiation of mucous epithelia and thus the nature and activity of the cells which form mucus. Consequently the composition of the secretion may be determined by the mechanism responsible for salt-equilibration and which is required to ensure the ionic stability of the tissue (Willmer, 1960). This may explain the relationship between the electrolyte abnormalities found in serous secretions and the extreme viscosity of mucus which characterizes fibrocystic disease. There is no really conclusive evidence for chemical differences in the mucoid constituents of comparable mucus samples from 'fibrocystics' and 'non-fibrocystics', but the electrolyte composition appears to be disturbed in certain mucous secretions from fibrocystic cases (Chernick, 1961; Chernick *et al.* 1961). The production of abnormal mucus in fibrocystic disease may therefore merely reflect alterations in the physiological conditions, especially the ionic environment, of the tissues which form mucus, and may be one result of a generalized metabolic disturbance throughout the body which may possibly be associated with the metabolism of acetylcholine (Roberts, 1959; P. G. Johansen & C. M. Anderson, unpublished work).

SUMMARY

1. Two samples of mucus have been obtained from the lower intestinal tract of patients with fibrocystic disease of the pancreas.

2. The mucoids were originally excreted as tough rope-like plugs of gelatinous material. They were relatively uncontaminated apart from considerable amounts of admixed serum proteins, which were completely removed by washing with 0.9% sodium chloride solution. Sample 1 appeared to be essentially homogeneous, although somewhat polydisperse, when examined in the ultracentrifuge.

3. The mucoids showed many similarities in their physical and chemical properties, and were remarkable for their lack of solubility and retention of their original form under a wide variety of conditions. They resembled typical epithelial mucosubstances in their chemical composition: fucose, sialic acid, hexosamine, galactose and amino acids were identified and determined. Sample 2 contained less total carbohydrate than sample 1 and a considerably lower proportion of fucose, the sialic acid:fucose ratios being 1 and 3 for samples 1 and 2 respectively; however, the relative proportions of the other sugars were similar in both mucoids.

4. Neuraminidase and 0.1N-sulphuric acid at 80° rapidly split off all the sialic acid in both samples. The mucoids were also potent inhibitors of haemagglutination by heated type A and B influenza viruses. Thus the sialic acid residues occur as end groups in the polysaccharide portion of the mucosubstances.

5. Apart from an unusually high phenylalanine content, the distribution of amino acids in the mucoids resembles that found in typical epithelial mucosubstances.

6. Inorganic ions appear to be important in stabilizing the structure of the mucoids.

7. The relationship between the abnormalities in mucous and serous secretions which characterize fibrocystic disease is discussed briefly.

The author thanks Dr C. M. Anderson for her interest and encouragement, and Miss M. Gilbo and Mr G. Wilson of the Commonwealth Serum Laboratories for undertaking the amino acid analyses and the examination of the mucoids in the analytical ultracentrifuge. This study was supported by a grant from the National Cystic Fibrosis Research Foundation, U.S.A.

REFERENCES

- Ada, G. L., French, E. L. & Lind, P. E. (1961). *J. gen. Microbiol.* **24**, 409.
- Arronsson, T., Grönwall, A. & Lausing, E. (1958). In *Protides of the Biological Fluids*, p. 233. Ed. by Peeters, H. Amsterdam: Elsevier Publishing Co.
- Bettelheim-Jevons, F. R. (1958). *Advanc. Protein Chem.* **13**, 35.
- Carsten, M. E. & Kabat, E. A. (1956). *J. Amer. chem. Soc.* **78**, 3083.
- Cessi, C. & Piliego, F. (1960). *Biochem. J.* **77**, 508.
- Chernick, W. S. (1961). *Post Grad. med. J.* **37**, 596.
- Chernick, W. S., Barbero, G. J. & Parkins, F. M. (1961). *J. Pediat.* **59**, 890.
- Conway, E. J. (1957). *Microdiffusion Analysis and Volumetric Error*, 4th ed. p. 152. London: Crosby Lockwood and Son, Ltd.
- Diehl, H. & Ellingboe, J. L. (1956). *Analyt. Chem.* **28**, 882.
- Dische, Z. (1950). *J. biol. Chem.* **183**, 489.
- Dische, Z. & Shettles, L. B. (1948). *J. biol. Chem.* **175**, 595.
- Gibbons, M. N. (1955). *Analyst*, **80**, 268.
- Gibbons, R. A. (1959). *Biochem. J.* **73**, 209, 217.
- Glegg, R. E. & Eiding, D. (1954). *Analyt. Chem.* **26**, 1365.

- Gomori, G. (1955). In *Methods in Enzymology*, vol. 1, p. 138. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
- Gottschalk, A. & McKenzie, H. A. (1961). *Biochim. biophys. Acta*, **54**, 226.
- Grabar, P. (1959). *Meth. biochem. Anal.* **7**, 1.
- Graham, E. R. B. & Gottschalk, A. (1960). *Biochim. biophys. Acta*, **38**, 513.
- Johansen, P. G. (1958). Ph.D. Thesis: University of London.
- Johansen, P. G., Marshall, R. D. & Neuberger, A. (1960). *Biochem. J.* **77**, 239.
- Kohn, J. (1958). *Clin. chim. Acta*, **3**, 450.
- Moore, S. & Stein, W. H. (1954a). *J. biol. Chem.* **211**, 893.
- Moore, S. & Stein, W. H. (1954b). *J. biol. Chem.* **211**, 907.
- Odin, L. (1959). *Acta Soc. med. upsalien.* **64**, 25.
- Partridge, S. M. (1949). *Nature, Lond.*, **164**, 443.
- Pusztai, A. & Morgan, W. T. J. (1961). *Biochem. J.* **78**, 135.
- Roberts, G. B. S. (1959). *Lancet*, ii, 964.
- Scheidegger, J. J. (1955). *Int. Arch. Allergy, N.Y.*, **7**, 103.
- Spicer, S. S. (1960). *J. Histochem. Biochem.* **8**, 18.
- Stone, J. D. (1949). *Aust. J. exp. Biol. med. Sci.* **27**, 337.
- Svennerholm, E. & Svennerholm, L. (1958). *Nature, Lond.*, **181**, 1154.
- Svennerholm, L. (1957). *Biochim. biophys. Acta*, **24**, 604.
- Svennerholm, L. (1958). *Acta chem. scand.* **12**, 547.
- Warren, L. (1959). *J. biol. Chem.* **234**, 1971.
- Willmer, E. N. (1960). *Cytology and Evolution*, pp. 189-200. New York and London: Academic Press Inc.
- Wilson, C. M. (1959). *Analyt. Chem.* **31**, 1199.
- Winzler, R. J. (1955). *Meth. biochem. Anal.* **2**, 279.

Biochem. J. (1963) **87**, 70

Studies on the Chemical Basis of the Antigenicity of Proteins

6. ANTIGENIC SPECIFICITY OF SOME SYNTHETIC POLYPEPTIDES CONTAINING TYROSINE*

BY SARA FUCHS AND M. SELA

Department of Biophysics, The Weizmann Institute of Science, Rehovoth, Israel

(Received 29 August 1962)

The main purpose of the investigation of synthetic polypeptide antigens is to determine the minimal structural requirements for immunogenicity, as well as to elucidate their antigenic specificity relationships. The synthesis, characterization and immunogenicity in rabbits of several linear and multichain polypeptides have been described by Sela, Fuchs & Arnon (1962). It was concluded that the immunogenically important area of the molecule must be accessible to the site of the biosynthesis of the antibody, that the overall shape of the molecule does not seem to be an important factor in determining immunogenicity, and that synthetic materials with molecular weights as small as 4000 may be good immunogens.

The present paper is concerned with studies on the specificity of some of the synthetic antigens described previously (Sela & Arnon, 1960a; Sela, 1962; Sela *et al.* 1962). In order to elucidate the chemical nature of the determinants of the immune specificity of several synthetic polypeptide antigens, the antisera investigated were allowed to react with a variety of linear and multichain copolymers of amino acids with compositions similar to those of the homologous immunogens as well as with unrelated amino acid compositions. Some proteins and polypeptidyl proteins were also allowed to react

with the antisera. Cross-precipitation experiments as well as inhibitions of the homologous reactions were used for this purpose. Similar studies on polypeptidyl derivatives of gelatin have shown that antibodies to polypeptide determinants possess rather narrow specificities (Arnon & Sela, 1960). The present findings show that the synthetic polypeptide antigens investigated contain determinants of well-defined specificity. Thus the antibodies formed against one antigen react well with chemically closely related substances and not at all with unrelated materials.

EXPERIMENTAL

Materials

The samples of gelatin, ovalbumin, poly-L-tyrosyl ovalbumin, edestin, poly-L-tyrosyl edestin, copolymers of L-tyrosine and L-glutamic acid in residue molar ratios of 1:1.1 (number-average degree of polymerization, n , 31), 1:4.0 (n , 31) and 1:9.0 (n , 88), a copolymer of L-tyrosine and DL-alanine in a residue molar ratio of 1:10.1 (n , 35), as well as a copolymer of L-tyrosine, L-leucine and L-glutamic acid in residue molar proportions of 1:1:1 (n , 39), were described by Sela & Arnon (1960b). The polytyrosyl gelatins, pTyrGel A and pTyrGel B, and copoly-(L-glutamyl-, L-tyrosyl) gelatin, pGluTyrGel, were described by Arnon & Sela (1960).

The synthesis, characterization and nomenclature of the linear and multichain polymers of α -amino acids listed

* Part 5: Sela, Fuchs & Arnon (1962).