Identification of the pentapeptide constituting a dominant epitope common to all eukaryotic heat shock protein 90 molecular chaperones

Jun Kishimoto,^{1,2} Yutaka Fukuma,² Akio Mizuno,¹ and Takayuki K. Nemoto²

¹Division of Oral and Maxillofacial Surgery, Department of Developmental and Reconstructive Medicine, Course of Medical and Dental Sciences, Nagasaki University Graduate School of Biomedical Sciences, 1-7-1 Sakamoto, Nagasaki 852-8588, Japan

²Division of Oral Molecular Biology, Department of Developmental and Reconstructive Medicine, Course of Medical and Dental Sciences, Nagasaki University Graduate School of Biomedical Sciences, 1-7-1 Sakamoto, Nagasaki 852-8588, Japan

Abstract We previously reported that, in human heat shock protein (Hsp) 90 (hHsp90), there are 4 highly immunogenic sites, designated sites Ia, Ib, Ic, and II. This study was performed to further characterize their epitopes and to identify the epitope that is potentially common to all members of the Hsp90 family. Panning of a bacterial library carrying randomized dodecapeptides revealed that Glu₂₅₁-Ser-X-Asp₂₅₄ constituted site Ia and Pro₂₉₅-Ile-Trp-Thr-Arg₂₉₉, site Ic. Site II (Asp₇₀₁-Pro₇₁₇) was composed of several epitopes. When 19 anti-hHsp90 monoclonal antibodies (mAbs) were subjected to immunoblotting against recombinant forms of 7 Hsp90-family members, 2 mAbs (K41110 and K41116C) that recognized site Ic bound to yeast Hsp90 with affinity identical to that for hHsp90, and 1 mAb (K3729) that recognized Glu₂₂₂-Ala₂₃₁ of hHsp90 β could bind to human 94-kDa glucose-regulated protein (Grp94), an endoplasmic reticulum paralog of Hsp90. Among the 5 amino acids constituting site Ic, Trp₂₉₇ and Pro₂₉₅ were essential for recognition by all anti-site-Ic mAbs, and Arg₂₉₉ was important for most of them. The necessity of Ile₂₉₆, Thr₂₉₈, and Arg₂₉₉, which are replaced by Leu, Met/Leu, and Lys, respectively, in some eukaryotic Hsp90, was dependent on the mAbs, and K41110 and K41116C could react with Hsp90s carrying these substitutions. From these data taken together, we propose that the pentapeptide Pro₂₉₅-Ile-Trp-Thr-Arg₂₉₉ of hHsp90 functions as an immunodominant epitope common to all eukaryotic Hsp90.

INTRODUCTION

Heat shock proteins (Hsps) exist essentially in all organisms. Reflecting their important and general roles as molecular chaperones, their amino acid sequences are highly conserved in every Hsp family even among eukaryotic and prokaryotic organisms. For instance, the amino acid sequences of Hsp60 (Venner et al 1990), Hsp70 (Hunt and Morimoto 1985), and Hsp90 (Rebbe et al 1987; Hickey et al 1989) of human cells are 48.2%, 47.2%, and 35.6% identical to their respective *Escherichia coli* orthologs, GroEL (Hemmingsen et al 1988), DnaK (Bardwell and Craig 1984), and HtpG (Bardwell and Craig 1987).

The sequence homology of Hsps led to the hypothesis

that the immune system could be triggered by bacterial antigens: Bacterial GroEL, for instance, shares a high degree of homology with host ortholog, which may result in an aberrant immune response and chronicity of inflammation (Ranney 1982; Loesche et al 1985; Tabeta et al 2000). However, it remains unknown whether or not the sequence homology of the Hsp family proteins is sufficient for all members of the family to share the same immunogenicity.

Hsp90 is composed of 3 domains, ie, N-terminal, middle, and C-terminal domains. The N-terminal domain (amino acids 1–220 or 230) carries an adenosine triphosphate (ATP) and geldanamycin-binding site (Prodromou et al 1997a; Stebbins et al 1997). The N-terminal domain of human (Stebbins et al 1997) and yeast Hsp90 (Prodromou et al 1997a) and the middle domain of yeast Hsp90 (Meyer et al 2003) have been reported. On the basis of

Correspondence to: Takayuki K. Nemoto, Tel: 81 95 849 7640; Fax: 81 95 849 7642; E-mail: tnemoto@net.nagasaki-u.ac.jp.

Received 15 March 2005; Revised 1 July 2005; Accepted 5 July 2005.

the results of crystallographic studies and the similarity of the structure, a molecular clamp model on the client binding has been proposed (Prodromou et al 1997b, 2000; Meyer et al 2003). The C-terminal domain (ca 200 residues) is responsible for the dimerization (Minami et al 1994; Nemoto et al 1995). The 3-dimensional structure of the C-terminal domain of Hsp90 remains unknown. Only the C-terminal half (amino acids 511–624) of the C-terminal domain of HtpG, a bacterial ortholog of mammalian Hsp90, has been reported (Harris et al 2004). Recent studies indicate that the middle domain carries the client-binding site (Meyer et al 2003), although an additional binding site might be present in the C-terminal domain (Young et al 1997; Scheibel et al 1998; Minami et al 2001; Harris et al 2004).

We previously produced 33 monoclonal antibodies (mAbs) against 2 isoforms of human Hsp90, ie, Hsp90 α (hHsp90 α) and Hsp90 β (hHsp90 β) (Nemoto et al 1997). Epitope-mapping analysis using an enzyme-linked immunosorbent assay (ELISA) with deletion mutants and an octapeptide multipin library demonstrated the existence of 4 highly immunogenic sites, designated sites Ia (amino acids 247–257), Ib (amino acids 263–270), Ic (amino acids 291–304), and II (amino acids 702–716). Sites Ia and Ib were localized within the highly charged region (amino acids 223–289) of hHsp90 α ; site Ic, C-terminally adjacent to the charged region; and site II, near the C-terminus. The amino acid sequences constituting sites Ia, Ib, and II vary between hHsp90 α and hHsp90 β . The highly charged region and the C-terminal one are dispensable for the growth of budding yeast (Borkovich et al 1989) and are deleted in HtpG, the bacterial ortholog. In contrast, the amino acid sequence of site Ic is highly conserved in eukaryotic Hsp90 and is relatively conserved in all organisms.

We recently compared the immunogenic property of the Hsp90 molecular chaperone with polyclonal antibodies (pAbs) raised against various species of Hsp90-family member and mAbs raised against human Hsp90 (hHsp90) (Kawano et al 2004). Among 33 mAbs developed against hHsp90 α and hHsp90 β , several mAbs that strongly reacted with yeast Hsc82 (yHsc82) recognized Asn₂₉₁-Ile₃₀₄, a conserved region of the family protein. The pAb raised against bacterial ortholog HtpG did not cross-react with hHsp90 α and hHsp90 β , human Grp94 (hGrp94) and human Trap1 (hTrap1), and vice versa. Hence, although mammalian Hsp90 shares some immunological reactivity with yHsc82, it is considerably distinct from its bacterial ortholog, HtpG and that from *E. coli* (EcHtpG) and *Porphyromonas gingivalis* (PgHtpG) (Kawano et al 2004).

In this study, we further characterized the epitopes of the anti-Hsp90 mAbs by a panning technique that selects the 12-amino acid sequences specifically recognized by

the anti-Hsp90 mAbs. We found that 2 mAbs that recognized site Ic reacted with yHsc82 and all eukaryotic Hsp90s examined.

MATERIALS AND METHODS

Materials

The materials used and their sources were as follows: expression vector pQE9 and plasmid pREP4, from Qiagen Inc. (Chatsworth, CA, USA); expression vector pTrcHis TOPO and TOP10 competent cells, from Invitrogen (Carlsbad, CA, USA); CNBr-activated Sepharose 4B and low-molecular weight markers, from Amersham Biosciences (Piscataway, NJ, USA); Wizard genomic deoxyribonucleic acid (DNA) purification kit, from Promega Corp. (Madison, WI, USA); kaleidoscope prestained standards, from Bio-Rad (Richmond, CA, USA); restriction enzymes and DNA-modifying enzymes, from Nippon Gene (Tokyo, Japan); Talon metal affinity resin, from Clontech Laboratories Inc. (Palo Alto, CA, USA); anti-Hsp90 mAb AC88 (SPA-830) from Stressgen Biotech Corp. (Victoria, British Columbia, Canada); alkaline phosphatase-conjugated rabbit anti-mouse Ig(G+A+M)IgG, from Zymed Laboratories Inc. (San Francisco, CA, USA).

Construction of bacterial expression vectors

Construction of plasmids carrying the full-length form of hHsp90 α , hHsp90 β , hGrp94, EcHtpG, and PgHtpG and plasmids carrying the N domain of EcHtpG (residues 1–336, designated EcHtpG-N) and hHsp90 α (residues 1–400, hHsp90 α -N) tagged with a histidine hexamer encoded by pQE9 were described previously (Nemoto et al 1996, 2001; Roi 1998; Kawano et al 2004). Y1090[pREP4] was transformed with the plasmids, and the transformants were selected on Luria broth agar plates containing 50 μ g/mL of ampicillin and 25 μ g/mL of kanamycin.

Plasmids encoding full-length forms of yHsc82 (Borkovich et al 1989) and hTrap1 (Song et al 1995; Chen et al 1996) were generously provided by Drs Y. Kimura (Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan) and W.-H. Lee (University of Texas, San Antonio, TX, USA), respectively. For construction of yHsc82 and hTrap1, DNA fragments encoding the full-length forms were amplified by polymerase chain reaction (PCR) and then directly inserted into the pTrcHis TOPO expression vector in frame according to the manufacturer's recommendation. TOP10 cells were transformed with the plasmids and selected on Luria broth agar plates containing 50 μ g/mL of ampicillin. The insertion and orientation of the DNA fragments were confirmed by conducting Hot Star PCR (Qiagen Inc.). The constructed plasmids were designated pTrcHis-yHsc82 and pTrcHis-hTrap1, respectively.

In vitro mutagenesis

Nucleotides ATC encoding Ile₂₉₆, TGG encoding Trp₂₉₇, ACC encoding Thr₂₉₈, and AGA encoding Arg₂₉₉ of hHsp90α-N were substituted with CTC encoding Leu₂₉₆, GGG encoding Gly₂₉₇, ATG encoding Met₂₉₈, and GGA encoding Gly₂₉₉, respectively, by PCR-based site-directed mutagenesis in combination with *DpnI* cleavage of a template. pQE9-hHsp90α-N (residues 1–400) was this template. As a result, pQE9-hHsp90α-N Trp297Gly, Thr298Met, Arg299Gly, and a double mutant Ile296Leu and Thr298Met were constructed. Mutagenesis was confirmed by DNA sequencing.

Expression and purification of recombinant proteins

After overnight cultivation of transformed bacteria at 37°C, recombinant proteins were expressed at 30°C for 4 hours in the presence of 0.2 mM isopropyl-β-D-thiogalactopyranoside. They were purified by affinity chromatography with a Talon affinity column according to the manufacturer's protocol except that 10 mM imidazole and 10 μg/mL of leupeptin were added to the lysis buffer (20 mM Tris-HCl, pH 8.0, containing 0.1 M NaCl and 0.5 mg/mL lysozyme). After extensive washing with washing buffer (20 mM Tris-HCl, pH 8.0, 0.1 M NaCl containing 10 mM imidazole) to remove nonadsorbed proteins, the bound proteins were eluted with 0.1 M imidazole (pH 8.0) containing 10% (vol/vol) glycerol. Purified proteins were used immediately or stored at –80°C until used.

Production of pAb

Affinity-purified recombinant EcHtpG-N (amino acids 1–336) was further purified by electrophoresis on a 7.5% polyacrylamide gel of 0.5-cm thickness under nondaturing conditions (Nemoto et al 1995). A separated protein was excised after observing with a Gelcode silver snap kit (Pierce, Rockford, IL, USA). A piece of a polyacrylamide gel containing the recombinant protein was homogenized and then injected into rabbits (Japan White) 5 times at 3-week intervals. The immunogen (approximately 1 mg) emulsified with Freund's complete adjuvant was used for primary immunization, and subsequent booster injections were given with the antigen in Freund's incomplete adjuvant. The pAbs were purified using EcHtpG-N-conjugated CNBr-activated Sepharose (Amersham Biosciences) prepared according to the manufacturer's recommendation. Antisera and pH-neutralized, purified pAb were divided into small aliquots and stored in the presence of 0.03% NaN₃ at –80°C. Animal care and experimental procedures were conducted in accordance with the Guidelines for Animal Experimentation of Nagasaki University with the approval of the Institutional Animal Care and Use Committee.

Fig. 1. Epitope map of the mAbs developed against hHsp90β - Recognition sites of 33 anti-hHsp90 mAbs are represented along with the 732 amino acid hHsp90α. N, M, and C represent N-terminal, middle, and C-terminal domains, respectively (Nemoto et al 1997). The hatched box (residues 223–289) represents highly charged region localized in the N-terminal domain. Isoform specificity of mAbs except for K3725A is indicated by their letters as follows: bold, hHsp90α specific; not modified, hHsp90α preferential; bold italic, hHsp90β specific; italic, hHsp90β preferential; and underlined, bound to both isoforms equivalently. The recognition sites of mAbs were identified as epitopes (indicated by boxes) or regions carrying the epitopes (indicated by bars). Highly immunogenic regions, sites Ia, Ib, Ic and II correspond to residues 247–257, 263–270, 291–304, and 702–716, respectively. Thirty mAbs were tested for panning as described in "MATERIALS AND METHODS": Amino acids responsible for the recognition were identified for 14 mAbs (indicated by *); Nine mAbs did not yield positive clones on immunoblotting (–); Seven mAbs yielded clones positive on immunoblotting, but the sequences they recognized either resembled each other nor matched the sequence of hHsp90s (±). Two mAbs without marks were not subjected to the panning. *precisely defined by the panning experiment in this study (see Tables 5–8). ±determined by phage display system in a previous study (Nemoto et al 1997 and see Table 4).

Anti-Hsp90 monoclonal antibodies

K3700 and K41000 number series represent the mAbs produced from hybridoma cells of mice immunized with hHsp90β and hHsp90α, respectively (Nemoto et al 1997). By setting 1% as the threshold for the binding affinity to 1 hHsp90 isoform, we classified 10 (Fig 1, bold letters) and 3 mAbs (bold italic letters) to be specific for hHsp90α and hHsp90β, respectively, and 19 mAbs reactive with both Hsp90 isoforms (nonbold letters). The mAb K3725A remained to be characterized because of its unusual behavior. The recognition sites of the 33 mAbs were described previously, which comprised 4 highly immunogenic sites, designated sites Ia, Ib, Ic, and II (Fig 1; Nemoto et al 1997).

Enzyme-linked immunosorbent assay

The binding affinity of the antibodies for the Hsp90-family proteins was determined by ELISA. Purified proteins

(1 µg) coated on each well of a Maxisorp C96 titer plate (Nalge Nunc Int, Rochester, NY, USA) were incubated with various concentrations of purified mAbs in 0.1 mL of the blocking buffer (0.17 M H₃BO₄, pH 8.5, containing 0.12 M NaCl, 0.05% [vol/vol] Tween 20, 1 mM ethylenediaminetetraacetic acid, and 0.05% [wt/vol] NaN₃ and 0.25% [wt/vol] bovine serum albumin) for 2 hours at room temperature as reported previously (Hornbeck et al 1994). After a 1-hour incubation with the anti-mouse Ig(G+A+M) and IgE conjugated to alkaline phosphatase (1/4000 dilution) and washing, the alkaline phosphatase activity was determined by measuring the absorbance at 655 nm after a 1-hour incubation with Bluephos micro-well phosphatase substrate (Kirkegaard and Perry Lab, Gaithersburg, MD, USA) at 30°C.

Panning of the FliTrix peptide library

Amino acid sequences recognized by the antibodies were determined using the FliTrix peptide library system (Invitrogen), according to the manufacturer's recommendation. In brief, an aliquot of the amplified FliTrix peptide library (2×10^{10} colony forming units) carrying randomized 12-amino acid sequences was screened by panning with anti-hHsp90 mAbs (20 µg/mL in 6-cm dish; Nalge Nunc Int, Roskilde, Denmark) or affinity-purified anti-EcHtpG-N pAb (40 µg/mL). After 5 cycles of panning, selected bacteria were plated on RMG plates [$1 \times$ M9 salts, 2% (wt/vol) casamino acids, 0.5% (wt/vol) glucose, 1mM MgC12, and 1.5% (wt/vol) agar, pH 7.4] containing 0.1 mg/mL ampicillin. After induction of protein expression of the bacteria in IMC medium [$1 \times$ M9 salts, 0.2% (wt/vol) casamino acids, 0.5% (wt/vol) glucose, 1mM MgC12, and 0.1 mg/ml ampicillin, pH 7.4] containing 0.1 mg/mL tryptophan for 6 hours at 37°C, positive clones were selected by immunoblotting of bacterial lysates with respective antibodies used for panning. Finally, the 12-amino acid sequences of the clones that were reactive with the antibodies were deduced by DNA sequencing.

Extraction of Hsp90-family proteins from tissues and cells

Tissue extracts of male Wistar rat liver and leaves of rice *Oryza sativa* var. Taichung65 at seedling stage, an extract of *Paramecium caudatum* strain RB1 prepared as reported previously (Hirabayashi et al 1983), and cell pellets of budding yeast *Saccharomyces cerevisiae* W303-1A and fission yeast *Schizosaccharomyces pombe* 972h⁻ were each mixed with 3 volumes of sodium dodecyl sulfate (SDS)-sample buffer containing 8 M urea. After boiling for 5 minutes, aliquots (10 µL) were subjected to immunoblotting.

Sodium dodecyl sulfate–polyacrylamide gel electrophoresis

Electrophoresis was performed in the presence of 0.1% (wt/vol) SDS at a polyacrylamide concentration of 11% or 12.5%. Generally, 0.25–0.5 µg of purified proteins or 5 µL of lysates was loaded onto an Sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) gel. Separated proteins were stained with Coomassie brilliant blue or subjected to immunoblotting as reported previously (Nemoto et al 1997). Low-molecular weight markers (Amersham Biosciences) and kaleidoscope prestained standards (Bio-Rad) were used as references for proteins stained with Coomassie brilliant blue and observed by immunoblotting.

Immunoblotting analysis

Purified proteins or bacterial lysates containing recombinant proteins separated on SDS-PAGE gels were subjected to immunoblotting analysis as described previously (Nemoto et al 1997). Tissue and cellular extracts of rat liver, budding yeast, fission yeast, rice, and *Paramecium* were prepared as reported previously (Hirabayashi et al 1983). First antibodies were used at 1–2 µg/mL for blotting of Hsp90 and at 4–8 µg/mL for panning clones in 0.1 M Tris-HCl, pH 7.5, containing 0.15 M NaCl, 0.05% (vol/vol) each of Triton X-100 and Tween 20 (TBS-TX). Alkaline phosphatase-conjugated rabbit anti-mouse Ig(G+A+M) IgG was used as the secondary antibody at 0.2 µg/mL in 10 mL TBS-TX. Blots were finally observed by incubation with 5-bromo-4-chloro-3-indolyl phosphate and nitro blue tetrazolium (Nakarai, Kyoto, Japan) at room temperature for 5–20 minutes.

Protein concentration

Protein concentrations were determined by the bicinchoninic acid method, using bovine serum albumin as a standard (Pierce).

RESULTS

Reactivity of anti-hHsp90 mAbs with the members of the Hsp90 family

Recently, the specificity of the anti-Hsp90 mAbs, which we developed against either hHsp90α or hHsp90β, has been characterized systematically with 7 recombinant Hsp90-family proteins, ie, hHsp90α, hHsp90β, hGrp94, hTrap1, yHsc82, EcHtpG, and PgHtpG (Kawano et al 2004). On the basis of the binding profiles, we classified the 19 mAbs that could react with both hHsp90α and hHsp90β into 4 categories: 4 mAbs that could blot hHsp90α, hHsp90β, and yHsc82, apparently with similar

Fig. 2. ELISA of mAbs that could interact with yHsc82 or hGrp94 - (a) Bindings of the 4 mAbs that could blot yHsc82 as intensively-stained bands were quantified by ELISA. (b) The bindings of K3729 was quantified by ELISA. Symbols for the 7 Hsp90-family proteins are the following: hHsp90α (open circle); hHsp90β (closed circle); hGrp94 (open square); hTrap1 (closed square); yHsc82 (open triangle); EcHtpG (closed triangle); and PgHtpG (closed inverted triangle).

intensities; 5 mAbs that reacted with hHsp90s intensively and yHsc82 with a lesser intensity; 9 mAbs that reacted with hHsp90α and hHsp90β but scarcely blotted other members; and finally, K3729, which could react with hGrp94 as well as with hHsp90α and hHsp90β (Kawano et al 2004).

Among the 9 mAbs that could blot yHsc82, 5 of them recognized site Ic, and all the 4 mAbs that blotted yHsc82 and hHsp90 with similar intensities recognized site Ic (mAbs with underlines in Fig 1). To determine their binding affinities quantitatively, we performed ELISA with purified proteins as antigens (Fig 2a). The bindings of K41110 and K41116C to yHsc82 were identical to those for hHsp90s (Fig 2a). The binding affinities of K41122B and K41220 for yHsc82 were 1/30 and 1/3, respectively, of those for hHsp90s. Thus, the 8 anti-site-Ic mAbs were

Table 1 Epitope mapping with anti-site-Ia mAbs^a

hHsp90α	242	E	E	E	K	E	K	E	E	K	E	S	E	D	K	P	E	257
hHsp90β	236	E	K	E	E	E	D	K	D	D	E	E	K	P				249
K41102 #1			G	P	M	<u>E</u>	I	D	R	S	G	D	V	D				
#2				G	G	V	S	A	<u>E</u>	S	L	D	H	A	W			
#3				R	Q	<u>T</u>	<u>E</u>	V	D	S	G	R	D	L	<u>E</u>			
K41116A #1				R	Q	M	<u>V</u>	<u>E</u>	K	<u>E</u>	S	L	D	Q	A			
#2				R	N	T	W	D	Y	W	<u>E</u>	S	L	D	V			
K41122A #1				C	G	K	R	A	L	R	<u>E</u>	S	L	N	R			
K41320 #1				R	P	L	G	L	Q	<u>E</u>	<u>E</u>	S	L	D	G			
#2				G	V	D	G	L	D	L	P	L	<u>E</u>	Q	L			

mAbs, monoclonal antibodies; hHsp, human heat shock protein.
^a Site Ia is underlined. Amino acids identical to hHsp90α are indicated by bold letters.

tentatively classified into 3 groups with respect to their binding to yHsc82: group 1, 2 mAbs (K41110 and K41116C) bound to yHsc82 with the affinity identical to that for hHsp90s; group 2, 3 mAbs (K41122B, K41220, and K41322) bound to yHsc82 with affinity lower than that for hHsp90s; and group 3, 3 mAbs (K3720, K41331, and K41338) that did not bind to yHsc82. On the other hand, there were 4 mAbs (K3738, K41315, K41016, and K41002) that could interact with yHsc82 and recognized the sites distinct from site Ic (Fig 1). They recognized the sites distinct from each other, and therefore, they were not further characterized.

Considering the sequence similarity in site Ic between yHsc82 and hHsp90 (see Table 8), the reactivity of the anti-site-Ic mAbs to yHsc82 was not surprising. Inversely, it is curious that the 8 anti-site-Ic mAbs bore a variety of reactivity toward yHsc82 from the potent binding of group-1 mAbs to no binding of group-3 mAbs. We will further investigate the reason for this later.

We also performed ELISA with K3729, which was the mAb solely reacted with hGrp94. It preferentially bound to hHsp90β over hHsp90α (Fig 2b). Its binding affinity for hGrp94 was one-third of that for hHsp90β but was still 3-fold higher than that for hHsp90α. Consistent with the results of the immunoblotting, K3729 did not react with other Hsp90-family members (Fig 2b).

Essential amino acids constituting site Ia

We performed panning experiments using the FliTrix peptide library and anti-hHsp90 mAbs. We subjected 30 mAbs to panning (indicated by marks attached to the names of mAbs in Fig 1). As a result, amino acids important for recognition were determined for 14 mAbs (indicated by asterisks); immunologically positive clones were not obtained for 9 mAbs (indicated by “-”); and immunologically positive clones were obtained for 7 mAbs, but they did not possess consensus sequences (indicated by “±”) (see Tables 1, 2, 4–7). The epitope of 1 mAb (K41007) determined by a phage display system was reported previously (Nemoto et al 1997).

Table 2 Epitope mapping with anti-site-Ic mAbs^a

hHsp90α	291	N	K	T	K	P	I	W	T	R	N	P	D	D	I	304
hHsp90β	283	N	K	T	K	P	I	W	T	R	N	P	D	D	I	296
Group 1																
K41110 #1				G	K	L	A	P	T	W	T	R	Q	S	A	
#2				G	D	V	R	F	S	R	V	W	M	R	G	
#3				F	V	P	A	E	R	Y	P	*	W	V	D	Q
#4				L	V	G	Q	F	H	T	P	A	W	A	E	
#5								T	M	R	P	I	W	A	R	S
K41116C #1				A	V	L	M	L	P	V	W	A	C	L	S	
#2				G	R	N	G	K	Q	T	M	W	S	G	G	
Group 3																
K3720 #1				A	A	S	S	R	V	P	I	W	S	R	R	
#2										V	W	T	R	A	E	L
#3								R	G	N	V	P	V	W	T	Q
K41331 #1				Y	M	R	F	N	T	P	I	W	A	S	T	
#2						V	G	A	F	G	P	W	W	S	D	H
#3				V	E	S	V	R	A	Y	P	I	W	A	R	
#4								V	T	T	P	P	W	T	A	P
K41338 #1				G	G	G	G	D	I	V	P	I	W	A	R	
#2								G	R	G	G	A	I	W	D	R
#3				F	M	G	F	S	L	P	V	W	L	D	P	

mAbs, monoclonal antibodies; hHsp, human heat shock protein.

^a Site Ic is underlined. Amino acids identical to hHsp90 are indicated by bold letters.

* Deletion.

We first determined the amino acids essential for the immunogenicity of the anti-site-Ia mAbs, ie, K41102, K41116A, K41122A, and K41320 (Table 1). Because K41000-series mAbs were derived from mice immunized with hHsp90α, the epitope sequences were homologous to the sequence of hHsp90α. In particular, Glu₂₅₁-Ser-X-Asp₂₅₄ appeared to be important for the recognition. Even if amino acids were substituted, the changes were restricted: Glu₂₅₁ was replaced by charged amino acids (Arg with K41102, #1 and Asp with K41102, #3); Ser₂₅₂ was replaced by Gln in 1 clone (K41320, #2); and Asp₂₅₄ was replaced by amino acids with similar properties, ie, charged amino acid (Arg) and Asn.

Essential amino acids constituting site Ib

Five mAbs, K41241, K41315, K41020, K41107, and K41233, recognized the immunogenic site Ib, Ser₂₆₃-Asp-Glu-Glu-Glu-Glu-Lys-Lys₂₇₀ (Nemoto et al 1997). The panning with K41020, K41107, and K41233 produced positive clones on immunoblotting, but amino acid sequences were not homologous to each other or to the sequence of site Ib (data not shown). The panning with the mAbs K41241 and K41315 failed to result in positive clones on immunoblotting.

Essential amino acids constituting site Ic

The presence of 8 anti-site-Ic mAbs categorized as having a variety of bindings to yHsc82 (Fig 2a and data not shown) strongly indicated the differential requirement of amino acids for their bindings. To evaluate this, we per-

Table 3 Amino acids that appeared at the 5 positions of site Ic in selected clones

Amino acid	Frequency		
	Total (%)	Group 1 (%)	Group 3 (%)
Pro ₂₉₅	13 (76)	5 (71)	8 (80)
Ile ₂₉₆	6 (35)	1 (14)	5 (50)
Ile/Val/Trp/Ala/Met ₂₉₆	14 (82)	5 (71)	9 (90)
Trp ₂₉₇	17 (100)	7 (100)	10 (100)
Thr ₂₉₈	4 (24)	1 (14)	3 (30)
Thr/Ala/Ser ₂₉₈	13 (76)	5 (71)	8 (80)
Arg ₂₉₉	8 (47)	3 (43)	5 (50)
Total	17 (100)	7 (100)	10 (100)

formed the panning of the peptide library using the mAbs with 2 typical binding characteristics to site Ic, ie, K41110 and K41116C (group-1 mAbs), which bound to site Ic with high affinity, and K3720, K41331, and K41338 (group 3), which did not bind to it. As a result, we obtained 17 clones that interacted with 5 anti-site-Ic mAbs (Table 2). Among them, 15 dodecapeptides were aligned along with the sequence of site Ic without ambiguity. One clone (K41110, #3) could be reasonably aligned if a deletion was introduced. The alignment of the clone (K41116C, #2) was less confirmative because there was little homology. However, the alignment seems to be reasonable if we consider the chemical characteristics of the 5 amino acids, ie, Thr-Met-Trp-Ser-Gly (Table 2, K41116C, #2) to refer to Pro₂₉₅-Ile-Trp-Thr-Arg₂₉₉ (site Ic), as discussed in the following paragraphs.

Table 2 shows that the region of site Ic defined in the previous study (Asn₂₉₁-Ile₃₀₄) should be narrowed into 5 amino acids, ie, Pro₂₉₅-Arg₂₉₉. The frequencies of respective amino acids appearing in the positive clones are summarized in Table 3. Trp₂₉₇ was the sole amino acid appearing in all clones. Thereafter, Pro₂₉₅ (76%) and Arg₂₉₉ (47%) were observed; Ile₂₉₆ was observed at 35% frequency and finally Thr₂₉₈ at 24%. Among the selected clones, Pro₂₉₅, Trp₂₉₇, and Arg₂₉₉ existed simultaneously in 5 clones (29%). The combination of Pro₂₉₅ with Trp₂₉₇ was found at 47% and that of Arg₂₉₉ with Trp₂₉₇ was at 18%. Taken together, Trp₂₉₇ with Pro₂₉₅ or Arg₂₉₉ (or both) was present at 94%.

Table 3 further indicates the amino acids permissible at positions of Ile₂₉₆ and Thr₂₉₈. At the position of Thr₂₉₈, Ala or Ser with properties similar to those of Thr was occasionally found. Thus, the percentage when Thr, Ala, or Ser was located at the position was 76% (Table 3). Similarly, in 7 clones (35%), hydrophobic amino acids, such as Val (5 clones), Ala (1 clone), Trp (1 clone), and Met (1 clone), were found at position 296. Hence, the percentage when Ile or other hydrophobic amino acids were located at the position of Ile₂₉₆ was 82%.

Finally, Table 3 highlights the difference in amino acid preference between group-1 and group-3 mAbs. The fre-

Table 4 Epitope mapping with anti-site-II mAbs^a

hHsp90α	701	D	P	T	A	D	D	T	S	A	A	V	T	E	E	M	P	P	L	E	G	D	D	D	724	
hHsp90β	693	D	<u>E</u>	<u>V</u>	<u>A</u>	<u>A</u>	<u>E</u>	<u>E</u>	<u>P</u>	<u>N</u>	<u>A</u>	<u>A</u>	<u>V</u>	<u>P</u>	<u>D</u>	<u>E</u>	<u>I</u>	<u>P</u>	P	L	E	G	D	E	D	716
K41007 ^a			W	V	A	*	D	T	S	Y																
K3705		D	E	V	P	A	H	V	D	G	E	L	N													
K3725B #1														E	P	D	E	V	A	M	E	V	R	M	G	
#2							V	S	T	L	I	E	L	P	D	E	I	P								
#3															D	E	I	P	E	P	A	V	P	C	G	G

mAbs, monoclonal antibodies; hHsp, human heat shock protein.

^a Reported in Nemoto et al (1997). Site II is underlined. Amino acids identical to hHsp90α (K41007) or hHsp90β (K3000 series) are indicated by bold letters.

* Deletion.

quencies of 5 amino acids emerging in the clones selected with group-3 mAbs were always higher than those selected with group-1 mAbs. In particular, it was noticeable that Ile₂₉₆ was observed only in 1 clone (14%) reactive against group-1 mAbs and in 5 clones (50%) against group-3 mAbs. Thr₂₉₈ was found in 1 clone (14%) reactive against group-1 mAbs and in 3 clones (30%) against group-3 mAbs. In contrast, the frequencies of Arg₂₉₉ and Pro₂₉₅ were comparable between the clones selected with the 2-group mAbs (Table 3). These findings strongly suggest that Ile₂₉₆ and Thr₂₉₈ are closely related to the specificity of anti-site-Ic mAbs. In fact, among the 5 amino acids of site Ic, Ile₂₉₆ is replaced by Leu in yHsc82 (see Table 8), which reasonably explains the specificity of the anti-site-Ic mAbs. So, the substitution of Asp₃₀₂ in hHsp90 to Ser in yHsc82 was not related to the binding of anti-site-Ic mAbs.

Essential amino acids constituting site II

We estimated amino acids important for the recognition of anti-site-II mAbs. As shown in Table 4, K41007 recognized Ala₇₀₅-Asp-Asp-Thr-Ser₇₀₉ of hHsp90α (Nemoto et al 1997); K3705, Asp₆₉₃-Glu-Val-Pro-Ala₆₉₇ of hHsp90β; and K3725, Pro₇₀₅-Asp-Glu-Ile-Pro₇₀₉ of hHsp90β. Reflecting amino acids of site II altered between hHsp90α and hHsp90β, the sequences recognized by the mAbs resembled either one of the hHsp90 isoforms (Table 4). Noticeably, distinct segments within site II were recognized by the mAbs. Thus, we concluded that sites Ia and Ic each formed a single epitope, whereas site II was composed of several epitopes.

Amino acids essential for K3729

We were interested in K3729, because it cross-reacted with hGrp94 (Fig 2b). Its epitope had remained ambiguous because it bound to 4 separate sites at residues 212–312 of hHsp90α, ie, residues 227–238, 258–270, 282–291, and 301–310 [see Table 5, hHsp90α (1)–(4) and Nemoto et al 1997]. Hence, we performed panning with K3729.

Although there were 4 potential epitope sites for

Table 5 Epitope mapping with K3729^a

hHsp90α	227	DKEVSDDEAE	EKEDKEE	EKEEKESE	DK	255
hHsp90β	222	<u>EKE I</u>	<u>SDDEAE</u>	<u>EKG***</u>	<u>EKEEDKDDEEK</u>	247
hGrp94	267	<u>TVEEPM</u>	<u>EEEEAAKE****</u>	<u>EKEESDDEAAV</u>		291
yHsc82	215	KEVPIPEE	EKKDEE	KK***	DEDDKKPKLE	240
K3729 #1		EVAESDDEAEV				
#2		EPGLIDDDERNT				
#3		SPLQGSTDDESA				
#4		RSDTDDERSSEG				
#5		NWGQVESDLEFE				
#6		SSLSDMHDCESV				
#7		AGVDTDWEPHGR				
hHsp90α (1)	227	DKEVSDDEAE	EKEE			238
hHsp90α (2)	258	I	EDVGSDEE	EKK		270
hHsp90α (3)	282	EKIDQE	ELN			291
hHsp90α (4)	301	PDDITN	EEYG			310

hHsp, human heat shock protein.

^a Amino acids of hHsp90β corresponding to the epitope of K3729 deduced from selected clones (#1–#7) and those of hHsp90α and hGrp94 homologous to the epitope sequence of hHsp90β are underlined. Amino acids of positive clones and hHsp90α [(1)–(4)] identical to the sequence of hHsp90β are indicated by bold letters.

* Deletion.

K3729, it was precisely defined from 1 clone (Table 5, #1), which carried the sequence identical to Glu₂₂₂-X-X-X-Ser-Asp-Asp-Glu-Ala-Glu₂₃₁ (E₂₂₂XXXSDDEAE₂₃₁) of hHsp90β (equivalent to Asp₂₂₇-Glu₂₃₆ of hHsp90α). Taking other clones into consideration, we propose that Ser₂₂₆-Asp-Asp-Glu₂₂₉ is the most important, in which Asp₂₂₇ and Glu₂₂₉ are indispensable. Ser₂₂₆ could be substituted by Thr, another amino acid harboring a hydroxyl group or charged amino acids, such as Asp and His. Noticeably, Glu₂₂₂ and Ile₂₂₅ of hHsp90β, which are the amino acids altered between the hHsp90 isoforms, were mimicked in clones #1 and #2, respectively. Considering the preferential binding of K3729 to hHsp90β, these amino acids should additionally contribute to the binding of K3729.

This assignment gave an answer as to why K3729 bound to 4 separate regions of hHsp90α in an earlier study (Nemoto et al 1997). Table 5 indicated that 1 (residues 227–238) of the 4 peptides corresponded to the true epitope of the mAb [hHsp90α(1)] but that the remaining ones [hHsp90α(2)–(4)] possessed similar sequences, ie, Ser₂₃₁-Asp-(Glu)-Glu₂₃₄, Asp₂₈₆-(Gln)-Glu₂₈₈, or Asp₃₀₁-Asp-(Ile₃₀₃). Because we used a multipin peptide library

Table 6 Epitope mapping with K41218^a

hHsp90α	68	S	K	L	D	S	*	G	K	E	L	H	I	N	L	I	81
hHsp90β	63	S	K	L	D	S	*	G	K	E	L	K	I	D	I	I	76
K41218 #1		V	L	D	*	*	G	A	E	L	K	E	R	E	N		
#2		A	G	N	Y	*	G	E	E	L	L	P	V	D			
#3		A	L	K	*	G	A	E	L	L	I	H	G	S			
#4		R	R	D	S	G	F	E	Q	I	W	G					
#5		W	D	S	*	G	F	E	L	S	G	P	G	K			

hHsp, human heat shock protein.

^a Amino acids identical to hHsp90α are indicated by bold letters.

* Deletion.

Table 7 Epitope mapping with K41002^a

hHsp90α	306	N	E	E	Y	G	E	F	Y	K	S	L	T	N	D	W	320
hHsp90β	298	Q	E	E	Y	G	E	F	Y	K	S	L	T	N	D	W	312
K41002 #1					D	R	G	E	G	P	R	I	N	A	R	C	
#2		A	I	M	P	K	E	F	D	K	S	E	W				
#3			A	G	E	F	G	D	L	R	R	A	R	A			

hHsp, human heat shock protein.

^a Amino acids identical to those of hHsp90α are indicated by bold letters.

covering amino acids 212–312 derived from hHsp90α (Nemoto et al 1997), the binding to the hHsp90α peptide was weaker than that to hHsp90β. Under the conditions used, the bindings to these 3 nonpeptide sites [hHsp90α(2)–(4)] were overestimated.

This assignment also solved the question of why K3729 was capable of interacting with hGrp94 with an affinity higher than that for hHsp90α. The hGrp94 possesses a sequence similar to the epitope of hHsp90β, and moreover, the sequence carries Glu₂₉₁, which is equivalent to Glu₂₂₂ of hHsp90β (Table 5, underlined). Note that Glu₂₂₂ of hHsp90β was partly involved in the binding of K3729 (Table 5, #1) and was altered to Asp₂₂₇ in hHsp90α.

Amino acids essential for K41218

The recognition site of K41218, which had been proved to exist within amino acids 48–196 (Nemoto et al 1997), was precisely determined to be Asp₇₁-Ser-Gly-X-Glu-Leu₇₆ in which Gly₇₃ and Glu₇₅ seemed to be essential

and Asp₇₁ and Leu₇₆ were important (Table 6). In addition, Leu₇₀ and Ile₇₈ may contribute to the binding.

Amino acids essential for K41002

The recognition site of K41002, which had been proved to exist within amino acids 290–312 (Nemoto et al 1997), was determined. As shown in Table 7, Gly₃₁₀-Glu-Phe-X-Lys-Ser₃₁₅ was recognized by K41002. Charged amino acids, such as Arg and Asp, could be located at the position of Lys₃₁₄. The region defined in this study (residues 310–315) is slightly shifted from the previous one (residues 290–312). However, this is inevitable because we had defined the region using various truncated forms, and the mAb could interact with hHsp90α 290–312 (Nemoto et al 1997).

Effect of the substitution of amino acids constituting site Ic

To evaluate the roles of the respective amino acids constituting site Ic, we expressed hHsp90α-N with mutated amino acids. When compared with the sequences corresponding to site Ic of eukaryotic Hsp90, Ile₂₉₆ is substituted to Leu in *Candida albicans* (Swoboda et al 1995), yHsc82 (Borkovich et al 1989), and heat-inducible Hsp82 (Farrelly and Finkelstein 1984) in budding yeast *S cerevisiae* (Table 8). In plant cytoplasmic Hsp90, Thr₂₉₈ is substituted to Met in bread wheat *Triticum aestivum* (NCBI accession, X98582) or to Leu in rice *O sativa* (japonica cultivar group) (AB111810). In *Tetrahymena thermophila* (AF151114) and *Toxoplasma gondii* (A344115), a double substitution to Leu₂₉₆ and Met₂₉₈ occurs. In a plastid-specific form of Hsp90, Thr₂₉₈ is replaced by Met in *Secale cereale* (Schmitz et al 1996), and the double substitution to Leu₂₉₆ and Met₂₉₈ occurs in *Arabidopsis thaliana* (Milioni and Hatzopoulos 1997). In *P tetraurelia* (Frankel et al 2001), the substitution occurs simultaneously at 3 positions (Leu₂₉₆, Met₂₉₈, and Lys₂₉₉). Considering these substitutions, we expressed hHsp90α-N with the mutations

Table 8 Comparison of site-Ic sequences of Hsp90 from various species

hHsp90α ^a /hHsp90β ^a	NKTKPIWTRNPDDI
hGrp94 ^a	NDIKPIWQRPSKEI
hTrap1 ^a	NTLQAIWMDPKDV
yHsc82 ^{a,b} /yHsp82 ^b	NKTKPLWTRNPDDI
EcHtpG ^a	NKAQALWTRNKSEI
PgHtpG ^a	INDTHPAWTKKPADL
rat Hsp90 ^b	NKTKPIWTRNPDDI
<i>Schizosaccharomyces pombe</i> Hsp90 ^b	NKTKPIWTRNPSEV
<i>Oryza sativa</i> Hsp90 ^b	NKQKPIWLKPEEI
<i>Paramecium tetraurelia</i> Hsp90	NKNKPLWMMKPPDDI

hHsp, human heat shock protein.

^a Hsp90-family members subjected to ELISA in Fig. 2.^b Hsp90-family members endogenously expressed in cells subjected to the immunoblotting in Fig. 5. In the last line, the sequence of that of *P. tetraurelia* Hsp90 is shown instead of *P. caudatum* used in this study. Amino acids identical to those of hHsp90α/hHsp90β are indicated by bold letters.

Fig. 3. Effect of mutation at Ile₂₉₆, Trp₂₉₇, Thr₂₉₈, and Arg₂₉₉ on the antibody binding - (a) Aliquots of purified hHsp90α-N wt (lane 1), Trp 297Gly (lane 2), Thr298Met (lane 3), Ile296Leu/Thr298Met (lane 4), and Arg299Gly (lane 5) were separated on SDS-PAGE and then stained with Coomassie brilliant blue (CBB) or subjected to the immunoblotting with K41110 or K41116C. Lane M, molecular markers. (b-d) ELISA was performed with hHsp90α-N wt (open circle), Trp297Gly (closed circle), Thr298Met (open square), and Ile296Leu/Thr298Met (open triangle) as antigens. Groups 1-3 were categorized based on the reactivity toward yHsc82 (see "RESULTS").

of Trp297Gly, Thr298Met, Ile296Leu/Thr298Met, and Arg299Gly. The recombinant proteins were purified by affinity chromatography (Fig 3a). hHsp90α Arg299Gly appeared to be partially degraded (lane 5).

We first analyzed the bindings of anti-site-Ic mAbs to the mutants at positions 296, 297, and 298. As a positive control, we evaluated the binding of K41102, which recognized site Ia (Table 1). The mAb equivalently bound to hHsp90α-N irrespective of the substitutions (data not shown). In contrast, the bindings of the anti-site-Ic mAbs were drastically altered by the substitutions. None of the 8 mAbs bound to Trp297Gly even at the highest antibody concentration used (Fig 3b-d). This finding faithfully reflected the appearance of Trp₂₉₇ in all clones selected by panning (Table 2). In contrast, the bindings for Thr298Met and a double mutant Ile296Leu and Thr298Met were dependent on the mAbs. The bindings of group-1 mAbs to Thr298Met were scarcely affected (Fig 3b); but the binding affinities for Thr298Met of the remaining 6 anti-site-Ic mAbs except for K41338 were decreased at least to one-third of those for the wild type. An additional substitu-

tion of Ile₂₉₆ to Leu further decreased the binding affinities for anti-site-Ic mAbs, but the effect was little for group-1 mAbs (Fig 3b). Similarly, the effect was relatively small on 2 group-2 mAbs (K41220 and K41322; Fig 3c), but 1 group-2 mAb (K41122B) and 2 group-3 mAbs (K3720 and K41331) showed a significant reduction (<1/300) in binding affinity. The remaining group-1 mAb (K41338) did not bind to the double mutant under the conditions used (Fig 3d). Therefore, we concluded that the binding affinities of group-1 mAbs for the double mutant Ile296Leu and Thr298Met were slightly decreased (1/3-1/2) but that those of the remaining 6 mAbs were significantly reduced at least to one-tenth. In particular, the substitution of Ile₂₉₆ to Leu significantly reduced the binding affinities of group-3 mAbs. With these data taken together, we conclude that Ile₂₉₆ and Thr₂₉₈ were important for group-3 mAbs (with the exception of K41338 for Thr₂₉₈) but were not critical for the recognition of group-1 mAbs.

We then evaluated the effect of the substitution of Arg₂₉₉ to Gly. The change slightly reduced the binding of

Fig. 4. Effect of mutation at Arg₂₉₉ on the antibody binding - ELISA was performed with hHsp90 α -N wt (open circle) and Arg299Gly (closed circle) as antigens by using 8 anti-site-Ic mAbs. Groups 1–3 were categorized based on the reactivity toward yHsc82.

K41116C and reduced the binding of K41110 ten-fold (Fig 4a), and caused a significant reduction ($<1/30$) in those of the remaining 6 anti-site-Ic mAbs (Fig 4 b,c). Especially, K3720 never bound to Arg299Gly even at the highest antibody concentration (Fig 4c). Thus, although Arg₂₉₉ was one of the important determinants of site Ic for most anti-site-Ic mAbs, the effect was little on the group-1 mAbs. In contrast, the mutation did not affect the binding

of mAbs that recognized the sites other than site Ic (data not shown).

We also performed immunoblotting analysis. Group-1 mAbs could blot Thr298Met, Ile296Leu and Thr298Met, and Arg299Gly, but scarcely blotted Trp297Gly (Fig 3a), consistent with the results of ELISA (Fig 3b).

Although we did not perform the mutation of Pro₂₉₅, this amino acid appeared to be important for antibody recognition because (1) the frequency (76%) of Pro₂₉₅ appearing in the selected clones was higher than the 47% of Arg₂₉₉ (Table 3) and (2) EcHtpG, in which Pro₂₉₅-Ile-Trp-Thr-Arg₂₉₉ of eukaryotic Hsp90 is converted to Ala-Leu-Trp-Thr-Arg (Table 8), was not recognized by any site-Ic mAb.

Immunogenicity of the region corresponding to site Ic of EcHtpG

Although the sequences corresponding to site Ic of hTrap1, EcHtpG, and PgHtpG were partially homologous to that of hHsp90 (Table 8), none of the anti-site-Ic mAbs bound to them (Fig 2a and data not shown). In view of the findings obtained by mutagenesis (Figs 3 and 4), either Pro₂₉₅ or Arg₂₉₉ replaced in these 3 proteins should explain this phenomenon: Pro₂₉₅ is replaced by Ala in EcHtpG and hTrap1 and Arg₂₉₉ is replaced by Lys in PgHtpG. Alteration of Ile₂₉₆ to Leu in EcHtpG and to Ala in PgHtpG, and that of Thr₂₉₈ to Met in hTrap1, may additionally contribute to the phenomenon. On the other hand, these facts raised a new question. Is the region of other Hsp90-family members equivalent to site Ic of hHsp90 immunogenic? Or is the triplet of Pro₂₉₅, Trp₂₉₇, and Arg₂₉₉ essential for this role? To address this issue, we chose EcHtpG, which carries Ala at the position of Pro₂₉₅ (Table 9). We performed panning of the peptide library with the pAb raised and purified with EcHtpG-N.

After 5 cycles of panning, all clones were positive on immunoblotting with the pAb. Then, we randomly selected 7 clones and sequenced the oligonucleotides of 36 bases. Among 7 clones, 5 clones (Table 9, #1–5) possessed 2 or 3 amino acids matching the site-Ic sequence of EcHtpG. Three clones (#2, #4, and #5) could be aligned in 2 ways. Interestingly, at the position of Arg₂₃₆ of EcHtpG, equivalent to Arg₂₉₉ of hHsp90 α , charged amino acids (Lys and Glu) were located in 3 clones (#1, #2[1], and #5[2]). On the other hand, the amino acids of both #6 and #7 were not similar to the site-Ic sequence, but they appeared to be partially homologous to a single site, Ile₃₆-Leu₄₆ (Table 9). So, we conclude that the site of EcHtpG equivalent to site Ic of hHsp90 α is also immunogenic although the sequence carries Ala at the position of Pro₂₉₅. In the 7 selected clones, we did not detect Trp at position 297. We do not know the reason for this at present.

Fig. 5. Immunoblotting analysis of the Hsp90-family proteins endogenously expressed in eukaryotes - Tissue or cell pellets of several species were denatured in SDS-sample buffer, and aliquots (5 μ l) were loaded onto SDS-PAGE gels. Separated proteins were immunoblotted with K41110, K41116C, K41220 or AC88/SPA-830 (Stressgen). Lane 1, rat liver; lane 2, budding yeast (*S. cerevisiae*); lane 3, fission yeast (*S. pombe*); lane 4, rice (*O. sativa*) and lane 5, *P. caudatum*. Lane M, kaleidoscope prestained standards.

Table 9 Epitope mapping of anti-EcHtpG-N pAbs^a

hHsp90 α	291	N	K	T	K	P	I	W	T	R	N	P	D	D	I	304
EcHtpG	228	N	K	A	Q	A	L	W	T	R	N	K	S	E	I	241
#1		A	G	G	A	V	W	N	E	V	D	L				
#2 (1)		L	K	A	L	G	L	N	M	E	P	V	P			
#2 (2)			L	K	A	L	G	L	N	M	E	P	V	P		
#3					I	G	C	I	M	R	N	A	G	T	A	N
#4 (1)		T	L	Q	A	L	K	G	C	T	R	R	Y			
#4 (2)			T	L	Q	A	L	K	G	C	T	R	R	Y		
#6 (1)					A	L	E	R	K	C	G	A	L	V	E	V
#6 (2)		A	L	E	R	K	C	G	A	L	V	E	V			
hHsp90 α	49	I	S	N	S	S	D	A	L	D	K	I	R	Y	E	62
EcHtpG	36	I	S	N	A	S	D	A	A	D	K	L	R	F	R	49
#5		I	N	V	T	L	F	V	A	D	N	L	P			
#7		T	E	T	E	V	G	L	R	D	G	L	G			

pAbs, polyclonal antibodies; hHsp, human heat shock protein.
^a Three clones (#2, #4, and #6) are aligned in 2 ways. The 5 amino acids essential for site Ic are underlined (see Table 2). Amino acids identical to those of EcHtpG are indicated by bold letters.

Immunoblotting analysis of Hsp90 endogenously expressed in various species

The findings in Figures 2b and 3c prompted us to examine the reactivity of group-1 mAbs to endogenous forms of eukaryotic Hsp90-family members (Table 8, lower 4 species). K41110 and K41116C specifically recognized 80–90 kDa species from rat liver, budding yeast (*S. cerevisiae*), fission yeast (*S. pombe*) (Aligue et al 1994), rice (*O. sativa*, AB111810), and *P. caudatum* (Fig 5). To the contrary, K41220, which belonged to group 2, did not bind to *P. caudatum* Hsp90 (lane 5) and scarcely bound to rice Hsp90 (lane 4). We also tested an anti-Hsp90 mAb AC88,

which has been reported to react with Hsp90 from various species (Riehl et al 1985). Although it was developed against eukaryotic filamentous fungi *Achlya ambisexualis*, it efficiently recognized rat Hsp90, as reported previously (Chen et al 1997). It blotted *P. caudatum* Hsp90 as a faint band and did not detect Hsp90 from other species (Fig 5).

DISCUSSION

Recently, we characterized the immunological cross-reactivity of the members of the Hsp90 family with a number of anti-hHsp90 mAbs. In combination with ELISA in this study, we concluded that 2 mAbs reacted with yHsc82 with affinity identical to that for hHsp90, and 7 mAbs reacted with yHsc82 with affinities lower than that for hHsp90. Also, 1 reacted with hGrp94 with a moderate affinity. In this study, we identified essential amino acids constituting various epitopes of hHsp90 α and hHsp90 β . In particular, 3 of the 4 highly immunogenic sites reported previously were further characterized in this study. As a result, we demonstrated that sites Ia and Ic each formed a single epitope, but showed that site II was composed of several epitopes. We could not define the amino acids essential for site Ib.

Among 11 amino acids constituting site Ia, reported previously (Nemoto et al 1997), Glu₂₅₁-Ser-X-Asp₂₅₄ of hHsp90 α was most important for the recognition by mAbs (Table 1). On the other hand, among 14 amino ac-

ids (Asn₂₉₁-Ile₃₀₄) of site Ic (Nemoto et al 1997), only 5 residues from Pro₂₉₅ to Arg₂₉₉ were required for the antibody recognition. The observations in this study indicate that the immunogenicity of Glu₂₂₂-Ala₂₃₀ in hHsp90 β is shared by Glu₂₈₁-Ala₂₈₉ of hGrp94 and that of site Ic in hHsp90 is shared by yHsc82.

The finding that hGrp94 shared immunogenicity with hHsp90 β at the epitope of K3729 (Table 5) suggests the usefulness of this mAb for the detection of hGrp94. However, usefulness of K3729 for the detection of Grp94 may be limited to mammals because the sequence is conserved in mouse (Mazzarella and Green 1987), bovine (AB025193), pig (de Crom et al 1999) Grp94 but is altered in chick (Kulomaa et al 1986), barley (Walther-Larsen et al 1993), and *A thaliana* (Ishiguro et al 2002).

Among 9 mAbs that reacted with yHsc82, 5 mAbs recognized site Ic. This may not be surprising in consideration of the sequence homology of site Ic (Table 8). However, it seems important to note that, among 8 anti-site-Ic mAbs, there were 3 anti-site-Ic mAbs that did not react with yHsc82. Among the 5 amino acids constituting the epitope of site Ic, Trp₂₉₇ was essential, and Pro₂₉₅ also appeared to be essential. The frequency of Arg₂₉₉ appearing in the panning clones followed that of Pro₂₉₅: Arg₂₉₉ was important for most site-Ic mAbs, but not essential for the immunoreactivity, because the mutant with Arg299Gly possessed potent reactivity toward 7 of the 8 anti-site-Ic mAbs (Fig 4). Moreover, K41116C showed the binding to Arg299Gly comparable with that for an intact form, and K41110 had considerable binding ability.

The frequencies of Ile₂₉₆ and Thr₂₉₈ that emerged in the panning clones were lower than the frequency of Arg₂₉₉. In fact, these 2 amino acids were less critical than Pro₂₉₅, Trp₂₉₇, and Arg₂₉₉ were for the 2 group-1 mAbs. It was not surprising that group-1 mAbs did not require Ile₂₉₆ because we classified anti-site-Ic mAbs on the basis of the reactivity to yHsc82, which carries Leu at position 296. However, it was noticeable that the 2 mAbs tolerated the substitution of Thr₂₉₈ to Met.

The 5 amino acids of site Ic are completely conserved in most eukaryotic Hsp90s, such as those of mammals (Rebbe et al 1987; Hickey et al 1989; Moore et al 1989; Huang et al 1999), chick (Binart et al 1989; Meng et al 1993), fission yeast *S pombe* (Aligue et al 1994), nematoda *Caenorhabditis elegans* (Massie et al 2003), and *Drosophila melanogaster* (Helmgren et al 1981). Moreover, Pro₂₉₅ and Trp₂₉₇ are completely conserved, and Arg₂₉₉ is highly conserved in eukaryotic Hsp90s.

The in vitro mutagenesis experiment combined with the immunoblotting analysis and ELISA demonstrated that K41110 and K41116C could interact with Pro₂₉₅-Ile(Leu)₂₉₆-Trp₂₉₇-Thr(Met)₂₉₈-Arg(Gly)₂₉₉. Rice Hsp90, which was recognized by the mAbs (Fig 5), carries Leu instead of Thr₂₉₈, which further indicates that Leu at po-

sition 298 could be blotted with the mAbs. Arg₂₉₉ was also one of the important determinants for most of the anti-site-Ic mAbs. However, 1 of the group-1 mAbs (K41116C) did not discriminate Arg299Gly from hHsp90 α -N (wild type) and another (K41110) also bound to Arg299Gly, although the binding affinity was decreased to one-tenth. Actually, they could blot *P caudatum* Hsp90 (Fig 5), which potentially possesses Arg₂₉₉, as does *P tetraurelia* Hsp90. Therefore, we conclude that anti-site-Ic mAbs that belong to group 1 recognize all eukaryotic Hsp90s.

The sequences of site Ic of hGrp94, hTrap1, EcHtpG, and PgHtpG are partially homologous to that sequence of hHsp90 (Table 8), but none of them were blotted with anti-site-Ic mAbs (Kawano et al 2004). Nevertheless, the binding affinities for the 4 recombinant proteins could be estimated by ELISA at higher antibody concentrations (Fig 2a). PgHtpG could be partially recognized by group-1 mAbs and K41122B (Fig 2b). In contrast, hGrp94 was never recognized by group-1 mAbs and was second least efficiently recognized by K41122B next to hTrap1 (Fig 2b). The site-Ic sequence of PgHtpG is Pro-Ala-Trp-Thr-Lys, of which Ile₂₉₆ and Arg₂₉₉ are substituted to Ala and Lys, respectively. Thus, the change of Arg₂₉₉ to Lys may primarily cause the loss of binding. However, as we demonstrated in Figures 3 and 4, the change of Arg₂₉₉ had little effect on group-1 mAbs, especially on K41116C. In view of this finding, we postulate that the second substitution of Ile₂₉₆ to Ala additionally abrogated the binding. It may be not surprising that the change of Ile₂₉₆ to Ala caused a more profound effect on the antibody binding than the change of Ile₂₉₆ with respect to Leu on group-1 mAbs.

Among the 7 recombinant proteins, hGrp94 was least efficiently bound by group-1 mAbs on ELISA (Fig 2a). However, the Pro₂₉₅, Trp₂₉₇, and Arg₂₉₉ triplet, and Ile₂₉₆ as well, are maintained in hGrp94 (Table 8). What does this apparent discrepancy imply? The sole amino acid substitution from Thr₂₉₈ to Gln should be responsible for this phenomenon, for Gln located at the position of Thr₂₉₈ prevented the antibody binding in a dominant-negative manner. To address this issue, we compared the amino acids located at the position of Thr₂₉₈. Thr is located at position 298 in EcHtpG and PgHtpG identical to hHsp90 α , and Met is present in hTrap1 (Table 8). In the panning experiment, Thr (4 clones), Ala (6 clones), and Ser (3 clones) were occasionally recovered; each of Met, Leu, Val, and Asp, respectively, emerged in 1 clone (Table 2), which strongly suggests that Met and Leu were permitted to be located at this position but that Gln₂₉₈ sterically interfered with the antibody binding. If this is true, inversely, the antibody raised against site Ic of hGrp94 would be expected to interact with hHsp90. In fact, we found that antibodies raised against site Ic of hGrp94 could interact with hHsp90s, although the affinity was

decreased to one-tenth (Kawano et al 2004). This discussion regarding Ala at the position of Ile₂₉₆ and Gln at the position of Thr₂₉₈ implies that Ile₂₉₆ and Thr₂₉₈ are also important for the recognition by group-1 mAbs as well as by group-3 mAbs, but that, principally, group-1 mAbs less stringently recognize Ile₂₉₆, Thr₂₉₈, and Arg₃₀₀ for their recognition than do group-3 mAbs.

We propose that K41110 and K41116C can be used as potent tools for detecting all eukaryotic Hsp90. Historically, AC88 (SPA830) has been used for the analysis of Hsp90 because it was the first-developed anti-Hsp90 mAb and interacts with Hsp90 from various species with the 2 isoforms of mammalian Hsp90 (Riehl et al 1985). However, its reactivity was limited in some groups of eukaryotes (Fig 5; Kawano et al 2004). Thus, although the contribution of AC88 in the previous studies should be admitted, K41110 and K41116C should be superior to AC88 for detection of eukaryotic Hsp90. Interestingly, Grp94 of some species, such as trypanozoma *Leishmania infantum* (Larreta et al 2000) and sea urchin *Strongylocentrotus purpuratus* (Smith et al 1996), possesses the sequence Pro-Ile-Trp-Thr-Arg identical to site Ic of hHsp90. Hence, Grp94 from these species could be recognized by all anti-site-Ic mAbs.

It was reported that antibodies against Hsp70 and Hsp90 were produced in patients with lupus erythematosus (Minota et al 1988; Twomey et al 1993). It is important to evaluate whether HtpG of *P. gingivalis* as well as that of *Actinobacillus actinomycetemcomitans*, another implicated pathogen, is truly involved in the autoimmune responses in those patients. However, we demonstrated that, on immunoblotting analysis, 19 anti-hHsp90 mAbs tested could not recognize PgHtpG or EcHtpG even when they were anti site Ic. Therefore, we suppose that, even if antibodies against hHsp90 are produced in periodontitis, the antigen is unlikely to be derived from bacteria but rather is hHsp90 of host cells.

As clearly demonstrated in this study, panning method of FliTrx peptide library is a powerful tool for determining the epitopes of mAbs. Moreover, this study further demonstrated that pAbs can be used for this technique. Serially truncated proteins and peptides produced by recombinant techniques, which are conventionally used for the identification of epitopes, could be substituted by this approach.

In addition to sites Ia and Ic, we defined essential amino acids of several anti-Hsp90 mAbs: K3705 recognized Asp₆₉₃-Ala₆₉₇ of hHsp90 β (Table 4); K3725B recognized Pro₇₀₅-Pro₇₀₉ of hHsp90 β (Table 4); K3729 primarily recognized Ser₂₂₆-Glu₂₃₁ of hHsp90 β (Table 5); K41218 recognized Asp₇₁-Leu₇₆ (Table 6); and K41002 recognized Gly₃₁₀-Ser₃₁₅ (Table 7). The information on the epitopes of these mAbs is useful for future investigation on the chaperone cycle of Hsp90.

The 3-dimensional structures of the N-terminal ATP-binding 220 or 230 amino acids of hHsp90 and yHsp82 and the middle domain (residues 273–528) of yHsp82 equivalent to residues 293–549 of hHsp90 α have been determined (Prodromou et al 1997a; Stebbins et al 1997; Meyer et al 2003), which indicates that the 5 amino acids of site Ic form a loop exposed outside the domain. Moreover, we found that anti-site-Ic mAbs (K41116C and K41220) could adsorb glucocorticoid receptor-Hsp90 complex from rat liver (J. Kishimoto and T.K. Nemoto, unpublished observation), which further indicated that site Ic is exposed to outside of an Hsp90 molecule, and moreover, that site Ic is not directly involved in client binding.

We also mapped the epitope of K41002 (amino acids Gly₃₁₀-Ser₃₁₅) on the structure. The site formed an α -helix adjacent to site Ic, and the most important amino segment, Gly₃₁₀-Glu-Phe₃₁₂, was located to the outside of the domain (Prodromou et al 1997a). Similarly, the epitope of K41218 was mapped at the surface of the N-terminal domain, forming a loop connecting an α -helix and a β -strand structure (Prodromou et al 1997a; Stebbins et al 1997). Other epitopes defined in this study may also be located on the surface of the Hsp90 molecule; however, they could not be mapped at present because they were located in either the highly charged region or the C-terminal region, the 3-dimensional structures of which remain unknown. Although the dimer-forming, C-terminal domain of HtpG of *E. coli* has been reported recently (Harris et al 2004), the C-terminal end of Hsp90 (amino acids 697–732), which carries immunogenic site II, can not be pictured yet because the corresponding region is deleted in HtpG.

We previously reported that, even if residues 289–389 and its adjacent C-terminal region, (residues 401–546) of hHsp90 were separately expressed, they formed a stable complex in vitro and in vivo (Tanaka et al 2001; Matsumoto et al 2002). With reference to the 3-dimensional structure (Meyer et al 2003), the interaction appears to be primarily maintained through the association of the 5 β -sheet strands (amino acids 324–392 in hHsp90 α) and the longest α -helix structure (amino acids 410–428). We further propose as the activation mechanism of Hsp90 molecular chaperone that heat stress induces the liberation of the oligomerization and client-binding site of residues 311–350 by disrupting the intramolecular interaction between the residues 289–389 and 401–546 (Tanaka et al 2001; Matsumoto et al 2002). In fact, there are several lines of evidence that amino acids 289–389 are involved in client binding. A binding site of hHsp90 β for Ser and Thr kinase Akt (phosphatidylinositol 3-kinase or protein kinase B) was mapped to amino acids 327–340, equivalent to 335–348 of hHsp90 α (Sato et al 2000). It has been proposed that amino acids 347–360 form a potential client-

binding site (Meyer et al 2003). We recently mapped amino acids 311–350 as an essential region for self-oligomerization/client binding (Nemoto et al 2004). Moreover, a similar β -sheet platform is involved in client binding of DnaK (Zhu et al 1996) and major histocompatibility complex–class I molecule (Fremont et al 1995). Therefore, it is interesting to investigate whether the accessibility of the antibodies to the 5 β -sheet strands as well as to the α -helix structure is modified, coupled with the heat-induced activation process of the Hsp90 molecular chaperone. Such an investigation is now under way in our laboratory.

Previous studies demonstrated that the presence of autoantibodies to Hsp90 in systemic lupus erythematosus (Minota et al 1988). Although Hsp90 is a cytosolic protein, surface expression of Hsp90 on peripheral blood mononuclear cells was found in a quarter of the patients with systemic lupus erythematosus during active state (Erkeller-Yuksel et al 1992). The level of Hsp90 has been shown to be elevated in peripheral blood mononuclear cells of a subset of systemic lupus erythematosus patients, mainly caused by transcription activation of *hsp90 β* gene (Twomey et al 1993). Kenderov et al (2002) proposed that immunopathogenesis of lupus nephritis is associated with Hsp90 as an autoantigen and that the pathology is associated with altered idiotypic regulation of the anti-Hsp90 IgG autoantibodies. On the other hand, it remains unknown whether or not there are epitopes of Hsp90 specifically involved in autoimmune responses. Thus, it seems important to determine the epitopes of Hsp90, such as those defined in this study, are involved in responses of autoimmune diseases.

ACKNOWLEDGMENTS

We greatly appreciate Ms R. Aoi (Institute of Immunology Company, Tochigi, Japan) for providing the anti-hHsp90 mAbs and Dr W.-H. Lee (University of Texas, San Antonio, TX, USA) for the gift of the hTrap1 plasmid. We also appreciate Drs I. Miyakawa (Yamaguchi University, Yamaguchi, Japan), M. Fujishima (Yamaguchi University, Yamaguchi, Japan), Y. Maeda (Tohoku University, Sendai, Japan), H. Hirano (Tokyo University, Tokyo, Japan), and H. Hayashida (Nagasaki University, Nagasaki, Japan) for providing various specimens used for the immunoblotting. This work was supported by a grant from the programs Grants-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Science, Sports, Culture, and Technology of Japan (to T.K.N.).

REFERENCES

Aligue R, Akhavan-Niak H, Russell P. 1994. A role of Hsp90 in cell cycle control: wee1 tyrosine kinase activity requires interaction with Hsp90. *EMBO J* 13: 6099–6106.

- Bardwell JC, Craig EA. 1984. Major heat shock protein gene of *Drosophila* and the *Escherichia coli* heat-inducible dnaK gene are homologous. *Proc Natl Acad Sci U S A* 81: 848–852.
- Bardwell JC, Craig EA. 1987. Eukaryotic Mr 83,000 heat shock protein has a homologue in *Escherichia coli*. *Proc Natl Acad Sci U S A* 84: 5177–5181.
- Binart N, Chambraud B, Dumas B, et al. 1989. The cDNA-derived amino acid sequence of chick heat shock protein Mr 90,000 (HSP90) reveals a “DNA like” structure: potential site of interaction with steroid receptors. *Biochem Biophys Res Commun* 159: 140–147.
- Borkovich KA, Farrelly FW, Finkelstein DB, Taulien J, Lindquist S. 1989. Hsp82 is an essential protein that is required in higher concentrations for growth of cells at higher temperatures. *Mol Cell Biol* 9: 3919–3930.
- Chen C-F, Chen Y, Dai K, Chen P-L, Riley DJ, Lee W-H. 1996. A new member of the hsp90 family molecular chaperones interacts with retinoblastoma protein during mitosis and after heat shock. *Mol Cell Biol* 16: 4691–4699.
- Chen H-S, Singh SS, Perdew GH. 1997. The Ah receptor is a sensitive target of geldanamycin-induced turnover. *Arch Biochem Biophys* 348: 190–198.
- de Crom R, van Haperen R, Janssens R, Visser P, Willemsen R, Grosveld F, van der Kamp A. 1999. Gp96/Grp94 is a putative high density lipoprotein-binding protein in liver. *Biochem Biophys Acta* 1437: 378–392.
- Erkeller-Yuksel FM, Isenberg DA, Dhillon VB, Latchman DS, Lydyard PM. 1992. Surface expression of heat shock protein 90 by blood mononuclear cells from patients with systemic lupus erythematosus. *J Autoimmun* 5: 803–814.
- Farrelly FW, Finkelstein DB. 1984. Complete sequence of the heat shock-inducible HSP90 gene of *Saccharomyces cerevisiae*. *J Biol Chem* 259: 5745–5751.
- Frankel J, Williams NE, Nelsen EM, Keeling PJ. 2001. An evaluation of Hsp90 as a mediator of cortical patterning in *Tetrahymena*. *J Eukaryot Microbiol* 48: 147–160.
- Fremont DH, Stura EA, Matsumura M, Peterson PA, Wilson IA. 1995. Crystal structure of an H-2Kb-ovalbumin peptide complex reveals the interplay of primary and secondary anchor positions in the major histocompatibility complex binding groove. *Proc Natl Acad Sci U S A* 92: 2479–2483.
- Harris SF, Shiau AK, Agard DA. 2004. The crystal structure of the carboxy-terminal dimerization domain of htpG, the *Escherichia coli* Hsp90, reveals a potential substrate binding site. *Structure* 12: 1087–1097.
- Helmgren R, Corces V, Morimoto R, Blackman R, Meselson M. 1981. Sequence homologies in the 5' regions of four *Drosophila* heat-shock genes. *Proc Natl Acad Sci U S A* 78: 3775–3778.
- Hemmingsen SM, Woolford C, van der Vies SM, Tilly K, Dennis DT, Georgopoulos CP, Hendrix RW, Ellis RJ. 1988. Homologous plant and bacterial proteins chaperone oligomeric protein assembly. *Nature* 333: 330–334.
- Hickey E, Brandon SE, Smale G, Lloyd D, Weber LA. 1989. Sequence and regulation of a gene encoding a human 89-kDa heat-shock protein. *Mol Cell Biol* 9: 2615–2626.
- Hirabayashi T, Tamura R, Mitsui I, Watanabe Y. 1983. Investigation of actin in *Tetrahymena* cells: a comparison with skeletal muscle actin by a devised two-dimensional gel electrophoresis method. *J Biochem* 93: 461–468.
- Hornbeck P, Winston SE, Fuller SA. 1994. Enzyme-linked immunosorbent assays. In: *Current Protocols in Molecular Biology*, ed. Asubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K. Current protocols (Greene Publishing Associates, Inc and John Wiley and Sons), 11.2.1–11.2.22.

- Huang H, Lee WC, Lin JH, Jian SC, Mao SJ, Yang PC, Huang TY, Liu Y. 1999. Molecular cloning and characterization of porcine cDNA encoding a 90-kDa heat shock protein and its expression following hyperthermia. *Gene* 226: 307–315.
- Hunt C, Morimoto RI. 1985. Conserved features of eukaryotic hsp70 genes revealed by comparison with the nucleotide sequence of human hsp70. *Proc Natl Acad Sci U S A* 82: 6455–6459.
- Ishiguro S, Watanabe Y, Ito N, Nonaka H, Takeda N, Sakai T, Kanaya H, Okada K. 2002. SHEPHERD is the *Arabidopsis* GRP94 responsible for the formation of functional CLAVATA proteins. *EMBO J* 21: 898–908.
- Kawano T, Kobayakawa T, Fukuma Y, et al. 2004. A Comprehensive study on the immunological reactivity of Hsp90 molecular chaperone. *J Biochem* 136: 711–722.
- Kenderov A, Minkova V, Mihailova D, et al. 2002. Lupus-specific kidney deposits of HSP90 are associated with altered IgG idiotypic interactions of anti-HSP90 autoantibodies. *Clin Exp Immunol* 129: 169–176.
- Kulomaa MS, Weigel NL, Kleinsek DA, et al. 1986. Amino acid sequence of a chicken heat shock protein derived from the complementary DNA nucleotide sequence. *Biochemistry* 25: 6244–6251.
- Larreta R, Sato M, Alonso C, Requena JM. 2000. Leishmania infantum: gene cloning of the GRP94 homologue, its expression as recombinant protein, and analysis of antigenicity. *Exp Parasitol* 96: 108–115.
- Loesche WJ, Syed SA, Schmidt E, Morrison EC. 1985. Bacterial profiles of subgingival plaque in periodontitis. *J Periodontol* 56: 447–456.
- Massie MR, Lapoczka EM, Boggs KD, Stine KE, White GE. 2003. Exposure to the metabolic inhibitor sodium azide induces stress protein expression and thermotolerance in the nematode *Caenorhabditis elegans*. *Cell Stress Chaperones* 8: 1–7.
- Matsumoto S, Tanaka S, Nemoto TK, et al. 2002. Interaction between the N-terminal and middle regions is essential for the in vivo function of HSP90 molecular chaperone. *J Biol Chem* 277: 34959–34966.
- Mazzarella RA, Green M. 1987. ERp99, an abundant, conserved glycoprotein of the endoplasmic reticulum, is homologous to the 90-kDa heat shock protein (hsp90) and the 94-kDa glucose regulated protein (GRP94). *J Biol Chem* 262: 8875–8883.
- Meng X, Jerome V, Devin J, Baulieu EE, Catelli MG. 1993. Cloning of chicken hsp90 β : the only vertebrate hsp90 insensitive to heat shock. *Biochem Biophys Res Commun* 190: 630–636.
- Meyer P, Prodromou C, Hu B, Vaughan C, Roe SM, Panaretou B, Piper PW, Pearl LH. 2003. Structural and functional analysis of the middle segment of Hsp90: implications for ATP hydrolysis and client-protein and co-chaperone interactions. *Mol Cell* 11: 647–658.
- Milioni D, Hatzopoulos P. 1997. Genomic organization of hsp90 gene family in *Arabidopsis*. *Plant Mol Biol* 35: 955–961.
- Minami M, Nakamura M, Emori Y, Minami Y. 2001. Both the N- and C-terminal chaperone sites of Hsp90 participate in protein folding. *Eur J Biochem* 268: 2520–2524.
- Minami Y, Kimura Y, Kawasaki H, Suzuki K, Yahara I. 1994. The carboxy-terminal region of mammalian HSP90 is required for its dimerization and function in vivo. *Mol Cell Biol* 14: 1459–1464.
- Minota S, Koyasu S, Yahara I, Winfield J. 1988. Autoantibodies to the heat-shock protein hsp90 in systemic lupus erythematosus. *J Clin Invest* 81: 106–109.
- Moore SK, Kozak C, Robinson EA, Ullrich SJ, Appella E. 1989. Murine 86- and 84-kDa heat shock proteins, cDNA sequences, chromosome assignments, and evolutionary origins. *J Biol Chem* 264: 5343–5351.
- Nemoto T, Matsusaka T, Ota M, Takagi T, Collinge DB, Walther-Larsen H. 1996. Dimerization characteristics of the 94-kDa glucose-regulated protein. *J Biochem* 120: 249–256.
- Nemoto T, Ohara-Nemoto Y, Ota M, Takagi T, Yokoyama K. 1995. Mechanism of dimer formation of the 90-kDa heat-shock protein. *Eur J Biochem* 233: 1–8.
- Nemoto T, Sato N, Iwanari H, Yamashita H, Takagi T. 1997. Domain structures and immunogenic regions of the 90-kDa heat-shock protein (HSP90): probing with a library of anti-HSP90 monoclonal antibodies and limited proteolysis. *J Biol Chem* 272: 26179–26187.
- Nemoto TK, Fukuma Y, Yamada S, Kobayakawa T, Ono T, Ohara-Nemoto Y. 2004. The region, adjacent to the highly immunogenic site and shielded by the middle domain, is responsible for the self-oligomerization/client binding of HSP90 molecular chaperone. *Biochemistry* 43: 7628–7636.
- Nemoto TK, Ono T, Kobayakawa T, Tanaka E, Baba TT, Tanaka K, Takagi T, Gotoh T. 2001. Domain-domain interactions of HtpG, an *Escherichia coli* homologue of eukaryotic HSP90 molecular chaperone. *Eur J Biochem* 268: 5258–5269.
- Prodromou C, Panaretou B, Chohan S, et al. 2000. The ATPase cycle of Hsp90 drives a molecular “clamp” via transient dimerization of the N-terminal domains. *EMBO J* 16: 4383–4392.
- Prodromou C, Roe SM, O'Brien R, Ladbury JE, Piper PW, Pearl LH. 1997a. Identification and structural characterization of the ATP/ADP-binding site in the Hsp90 molecular chaperone. *Cell* 90: 65–75.
- Prodromou C, Roe SM, Piper PW, Pearl LH. 1997b. A molecular clamp in the crystal structure of the N-terminal domain of the yeast Hsp90 chaperone. *Nat Struct Biol* 4: 477–482.
- Ranney RR. 1982. Bacteriology of severe periodontitis in young adult humans. *Infect Immun* 38: 1137–1148.
- Rebbe NF, Ware J, Bertina RM, Modrich P, Stafford DW. 1987. Nucleotide sequence of a cDNA for a member of the human 90-kDa heat-shock protein family. *Gene* 53: 235–245.
- Riehl RM, Sullivan WP, Vroman BT, Bauer VJ, Pearson GR, Toft DO. 1985. Immunological evidence that the nonhormone binding component of avian steroid receptors exists in a wide range of tissues and species. *Biochemistry* 24: 6586–6591.
- Roi R. 1998. Structures and expression of HSP90 family proteins [in Japanese]. *Jpn J Oral Biol* 40: 528–541.
- Sato S, Fujita N, Tsurui T. 2000. Modulation of Akt kinase activity by binding to Hsp90. *Proc Natl Acad Sci U S A* 97: 10832–10837.
- Scheibel T, Weikl T, Buchner J. 1998. Two chaperone sites in Hsp90 differing in substrate specificity and ATP dependence. *Proc Natl Acad Sci U S A* 95: 1495–1499.
- Schmitz G, Schmidt M, Feierabend J. 1996. Characterization of a plastid-specific HSP90 homologue: identification of a cDNA sequence, phylogenetic descent and analysis of its mRNA and protein expression. *Plant Mol Biol* 30: 479–492.
- Smith LC, Chang L, Britten RJ, Davidson EH. 1996. Sea urchin genes expressed in activated coelomocytes are identified by expressed sequence tags. Complement homologues and other putative immune response genes suggest immune system homology within the deuterostomes. *J Immunol* 156: 593–602.
- Song HY, Dunbar JD, Zhang YX, Guo D. 1995. Identification of a protein with homology to hsp90 that binds to type 1 tumor necrosis factor receptor. *J Biol Chem* 270: 3574–3581.
- Stebbins CE, Russo AA, Schneider C, Rosen N, Hartl FU, Pavletich NP. 1997. Crystal structure of an hsp90-geldanamycin complex: targeting of a protein chaperone by an antitumor agent. *Cell* 89: 239–250.
- Swoboda RK, Bertram G, Budge S, Gooday GW, Gow NA, Brown

- AJ. 1995. Structure and regulation of the HSP90 gene from the pathogenic fungus *Candida albicans*. *Infect Immun* 63: 4506–4514.
- Tabeta K, Yamazaki K, Hotokezaka H, Yoshie H, Hara K. 2000. Elevated humoral immune response to heat shock protein 60 (hsp60) family in periodontitis patients. *Clin Exp Immunol* 120: 285–293.
- Tanaka E, Nemoto TK, Ono T. 2001. Liberation of the intra-molecular interaction as the mechanism of heat-induced activation of HSP90 molecular chaperone. *Eur J Biochem* 268: 5270–5277.
- Twomey BM, Dhillon VB, McCallum S, Isenberg DA, Latchman DS. 1993. Elevated levels of the 90-kD heat shock proteins in patients with systemic lupus erythematosus are dependent upon enhanced transcription of the hsp90 β gene. *J Autoimmun* 6: 495–506.
- Venner TJ, Singh B, Gupta RS. 1990. Nucleotide sequences and novel structural features of human and Chinese hamster Hsp60 (chaperonin) gene families. *DNA Cell Biol* 9: 545–552.
- Walther-Larsen H, Brandt J, Collinge DB, Thordal-Christensen H. 1993. A pathogen-induced gene of barley encodes a HSP90 homologue showing striking similarity to vertebrate forms resident in the endoplasmic reticulum. *Plant Mol Biol* 21: 1097–1108.
- Young JC, Schneider C, Hartl FU. 1997. *In vitro* evidence that hsp90 contains two independent chaperone sites. *FEBS Lett* 418: 139–143.
- Zhu X, Zhao X, Burkholder WF, Gragerov A, Ogata CM, Gottesman ME, Hendrickson WA. 1996. Structural analysis of substrate binding by the molecular chaperone DnaK. *Science* 272: 1606–1614.