a similarly systematic and long lasting way as are patients with high grade lesions. Inadequate follow-up is a major factor, as suggested in the literature.⁶⁻⁹

Treatment of cervical intraepithelial neoplasia is, however, effective. In an earlier study of CIN 3 lesions,¹⁰ it was estimated that 28-39% of cases without treatment would progress to invasive cancer. Our data contained 837 cases of CIN 3 lesions in women aged 30-59, thus on the basis of the previous estimate, 234-326 would develop into invasive cervical cancers, not the three observed. The treatment effect might have been nearly 100%.

Contributors: IK contributed to the study conception and design, analysis and interpretation of the data, and drafting the manuscript. AA contributed to the study conception and design, interpretation of the data, and drafting and revising the manuscript. EP contributed to the analysis and interpretation of the data and drafting the article. PN contributed to the study conception and design, interpretation of the data, and drafting and revising the manuscript. He is guarantor.

Funding: This study was partially financed from a grant from the Finnish Cancer Organisation.

Competing interests: None declared.

Ethical approval: This study was approved by the ethical committee of obstetrics and gynecology E8 of the Helsinki-Uusimaa hospital district (decision No 150/E8/03).

- 1 International Agency for Research on Cancer. Cervical cancer screening. *IARC handbooks of cancer prevention*, vol 10. Lyon: IARC/WHO, 2005:302.
- 2 Anttila A, Läärä E. In: Sankila R, Démaret E, Hakama M, Lyng E, Schouten LJ, Parkin DM, for the European Network of Cancer Registries. *Evaluating and monitoring of screening programmes*. Brussels/Luxembourg: European Commission, Europe Against Cancer Programme, 2000.
- 3 Martin-Hirsch PL, Paraskevaidis E, Kitchener H. Surgery for cervical intraepithelial neoplasia. *Cochrane Database Syst Rev* 1999;(3):CD001318. DOI:10.1002/14651858.CD001318.
- 4 Soutter WP, de Barros Lopes A, Fletcher A, Monaghan JM, Duncan ID, Paraskevaidis E, et al. Invasive cervical cancer after conservative therapy for cervical intraepithelial neoplasia. *Lancet* 1997;349:978-80.
- 5 Teppo L, Pukkala E, Lehtonen M. Data quality and quality control of a population-based cancer registry. *Acta Oncol* 1994;33:365-9.
- 6 Andersson-Ellström A, Seidal T, Grannas M, Hagmar B. The pap-smear history of women with invasive cervical squamous carcinoma. A case-control study from Sweden. *Acta Obstet Gynecol Scand* 2000;79:221-6.
- 7 Sasieni PD, Cuzik J, Lynch-Farmery E, the National Co-ordinating Network for Cervical Screening Group. Estimating the efficacy of screening by auditing smears histories of women with and without cervical cancer. *Br J Cancer* 1996;73:1001-5.
- 8 Baldauf J-J, Dreyfuss M, Ritter J, Meyer P, Philippe E. Screening histories of incidence cases of cervical cancer and high grade SIL: a comparison. *Acta Cytol* 1997;41:1431-8.
- 9 Gornall RJ, Boyd IE, Manolitsas T, Herbert A. Interval cervical cancer following treatment for cervical intraepithelial neoplasia. *Int J Gynecol Cancer* 2000;10:198-202.
- 10 Hakama M, Räsänen-Virtanen U. Effect of a mass screening program on the risk of cervical cancer. *Am J Epidemiol* 1976;103:512-7.

(Accepted 21 October 2005)

doi 10.1136/bmj.38663.459039.7C

Tellymedicine

In the past, doctors had to see a patient before they could make a diagnosis. Doctors were also restricted in whom they could tell about this diagnosis—the patient and his or her trusted next of kin, the nurse, and perhaps the odd interested colleague. Recent advances in technology first brought us telemedicine, where doctors can assist other doctors in making a diagnosis, guide surgical procedures, or even perform surgery themselves, without being physically near the patient.

More recently, we have seen tellymedicine. This enables us to make a diagnosis in patients for whom we have no responsibility. We don't need to examine them, nor do we need a formal invitation to become involved in their medical care. In fact, we may only know of these patients from watching them on television or from reading about them in a newspaper. One of the most accomplished tellydoctors is Dr Gupta, a neurosurgeon who is also a senior medical correspondent for the US Cable News Network (CNN) (www.cnn.com/2005/HEALTH/03/30/otsc.gupta/). Dr Gupta diagnosed coronary artery disease in former US president Bill Clinton almost before his cardiologist did.

Another recent triumph of tellymedicine was recorded during the death of the late pope. The reluctance of the pontiff's doctors to release information threatened to turn things into a private Vatican matter. Fortunately, tellydoctors in several countries used their skills in remote diagnostics to provide the international media with detailed information on the pope's medical condition during his last days.

Forensic medicine is one particular field that can benefit from tellymedicine. For instance, Volkert van der Graaf, the assassin of Dutch politician Pim Fortuyn, was admitted to a clinic for forensic psychiatry so that a single expert witness could report to the court. Because Volkert refused cooperation, this

psychiatric evaluation was extremely lengthy.

Fortunately, several "Tele" psychiatrists, after studying the media, were able to conclude that Volkert suffered from Asperger's syndrome, and informed the court and public of this through the media long before his trial.

In the UK, Professor David Southall sought a child protection investigation after watching a television programme. He claimed to see what others had not but for his pains was found guilty of serious professional misconduct by the General Medical Council.

Tellymedicine implies that an insightful diagnosis can be made through careful observation alone, and that modern physicians have lost this gift and have come to rely too much on their tradition of diagnostic tests and procedures, requiring the physical presence of the patient. Tellymedicine obviates all these costly and time consuming tests. Patients are spared pointless visits to our clinics. As an additional benefit, they can read their diagnosis in the newspapers or watch it on television in the comfort of their home.

Jurgen A H R Claassen *consultant geriatrician, department of geriatric medicine, Radboud University Medical Centre, Nijmegen, Netherlands*
(j.claassen@ger.umcn.nl)

We welcome articles up to 600 words on topics such as *A memorable patient, A paper that changed my practice, My most unfortunate mistake*, or any other piece conveying instruction, pathos, or humour. Please submit the article on <http://submit.bmj.com>. Permission is needed from the patient or a relative if an identifiable patient is referred to. We also welcome contributions for "Endpieces," consisting of quotations of up to 80 words (but most are considerably shorter) from any source, ancient or modern, which have appealed to the reader.