Purplish, pruritic papules on the limbs

JENNIFER CLAY CATHER, MD, AND M. ALAN MENTER, MD

A 46-year-old white man presented with an acute 3- to 4-week history of an intensely pruritic eruption that mainly involved the limbs. Individual lesions appeared to predominate around the wrist flexures. Examination revealed multiple, flat, purplish lesions, 5 to 15 mm in diameter, with coalescence in areas of scratching (*Figure 1*). Whitish plaques were

seen on the oral mucous membrane (*Figure 2*). The patient was otherwise well, although on further questioning he stated that he felt less energetic than usual. He reported no recent illness or medication ingestion.

What is the diagnosis, and what further laboratory tests are indicated?

Figure 1. Purplish, flat-topped papules on the wrist; a few are in a linear arrangement.

Figure 2. A white reticulated pattern can be seen inside the right cheek; lesions cannot be removed with a tongue blade.

From the Division of Dermatology, Department of Internal Medicine, Baylor University Medical Center, Dallas, Texas.

Corresponding author: Jennifer Clay Cather, MD, 5310 Harvest Hill Road, Suite 260, Dallas, Texas 75230.

DIAGNOSIS: Lichen planus.

DISCUSSION

Lichen planus is a T-lymphocyte-mediated disease that affects both the skin and mucous membranes (1). Typically, skin lesions are pruritic, purple, polygonal, flat-topped papules that preferentially involve the flexor surfaces (especially the wrist) and trunk and only rarely involve the face (2). Close examination of a papule with a magnifying lens reveals a network of gray streaks (Wickham’s striae). Rare variants of lichen planus include lesions that are hypertrophic (on the shins), bullous (on the oral mucosa), and ulcerative or erosive (on the oral mucosa, palms, and soles). The isomorphic response, or Koebner’s phenomenon, is characterized by the production of lesions at areas of trauma (i.e., scratches, bites, surgical sites, and healed lesions of herpes zoster) and occurs commonly in lichen planus. Koebner’s phenomenon is a helpful clinical clue because it is found in only a few dermatological diseases (Table 1). Resolution of lesions may result in hyper- or hypopigmentation, frequently persisting for months to years.

Mucous membranes are involved in two thirds of patients with skin lesions and may be the sole area of involvement. The inside cheeks (buccal mucosa) characteristically have silver-white, pin-sized papules in a lacelike distribution. Lesions may also occur on the lips, palate, and tongue and less commonly on the vaginal mucous membranes. Rarely, oral lichen planus may ulcerate, producing chronic, painful erosions of the gums and mucosa. Long-standing mucosal lichen planus, especially oral ulcerative lichen planus, has been associated with carcinoma (3, 4). On the glans penis, minute, flat, polygonal papules may be arranged in rings.

Nail changes are seen in 5% to 10% of patients (2). Pterygium formation is distinctive in lichen planus (Figure 3). Other nail changes include longitudinal grooving, ridging, and splitting and a peculiar midline fissure. Proximal onycholysis (loosening) may occur with lichen planus.

Scalp lesions are seen infrequently. When present they may produce a scarring alopecia (2). Follicular lichen planus (lichen planopilaris) is characterized by spiny erythematous follicular papules in the scalp that result in a scarring alopecia. Scalp le-

Figure 3. Absent right thumbnail plate with tenting of skin and scar.

Table 2. Disorders and other conditions associated with lichen planus

Autoimmune disorders	Chronic liver disease
Alopecia areata	Hepatitis B and C
Bullous pemphigoid	Primary biliary cirrhosis
Dermatitis herpetiformis	
Dermatomyositis	Other
Diabetes mellitus	HLA-DR1 antigen
Hashimoto's thyroiditis	Thymoma
Morphea	Urolithiasis
Myasthenia gravis	
Pemphigus (foliaceus/vulgaris)	
Pernicious anemia	
Sjögren's syndrome	
Systemic sclerosis	
Ulcerative colitis	
Vitiligo	

Table 1. Dermatological diseases in which Koebner's phenomenon occurs

Contact dermatitis	Molluscum contagiosum
Darier's disease	Pellagra
Dermatographism	Perforating disorders
Eruptive xanthoma	Pityriasis rubra pilaris
Erythema multiforme	Porokeratosis
Henoch-Schönlein purpura	Psoriasis*
Kyrle's disease	Reactive perforating collagenosis
Lichen nitidus	Rhus dermatitis*
Lichen planus*	Verrucae*
Lichen sclerosus et atrophicus	Vitiligo

*Most frequently seen

sions may occur alone or with other lesions on skin or mucosal surfaces.

Very rarely, lichen planus has been described in the esophagus and the rest of the gastrointestinal tract.

Skin biopsy specimens reveal a “saw-tooth” epidermal hyperplasia and a lichenoid interface infiltrate with vacuolar alteration along the basement membrane. Necrotic keratinocytes are present along the basal layer of the epidermis and in the dermis (Civatte bodies). Biopsy specimens of drug-induced lichen planus may have a considerable number of eosinophils. Direct immunofluorescence reveals subepidermal clumps of immunoglobulin (Ig) M (less often IgG, IgA, or C3), which correspond to the Civatte bodies (2).

Numerous diseases have been associated with lichen planus (Table 2). Most recently the association with hepatitis, particularly hepatitis C, has been stressed in the literature (5, 6). Thus, all patients with a clinical or histological diagnosis of lichen planus should be evaluated with this in mind. In addition, medications have been associated with lichen planus-type drug eruptions (Table 3) (7).

Table 3. Medications frequently associated with lichen planus*

Antimalarials	Penicillamine
Arsenic	Phenothiazine derivatives
Chlorpropamide	Propranolol
Gold	Quinidine
Methyldopa	Spironolactone
Naproxen	Thiazide derivatives

*Compiled from reference 7.

Treatment is primarily required for symptomatic patients (i.e., those with pruritus or pain from ulcers) or patients with generalized lichen planus. Corticosteroids (topical, intralesional, and systemic) are the mainstay of therapy. Retinoids (topical and systemic) (8–11), dapsone (12), cyclophosphamide (13), cyclosporine (14–16), metronidazole, hydroxychloroquine (17), tacrolimus (18), thalidomide (19), and phototherapy have also been helpful. The recent introduction of topical tacrolimus has been valuable for patients with intractable mouth involvement (18).

- Schiller PI, Flaig MJ, Puchta U, Kind P, Sander CA. Detection of clonal T cells in lichen planus. *Arch Dermatol Res* 2000;292:568–569.
- Boyd AS, Nelder KH. Lichen planus. *J Am Acad Dermatol* 1991;25:593–619.
- Mignogna MD, Lo Muzio LL, Lo Russo LL, Fedele S, Ruoppo E, Bucci E. Clinical guidelines in early detection of oral squamous cell carcinoma arising in oral lichen planus: a 5-year experience. *Oral Oncol* 2001;37:262–267.
- Sigurgeirsson B, Lindelof B. Lichen planus and malignancy. An epidemiologic study of 2071 patients and a review of the literature. *Arch Dermatol* 1991;127:1684–1688.

- Beaird LM, Kahloon N, Franco J, Fairley JA. Incidence of hepatitis C in lichen planus. *J Am Acad Dermatol* 2001;44:311–312.
- Bonkovsky HL, Mehta S. Hepatitis C: a review and update. *J Am Acad Dermatol* 2001;44:159–182.
- Ellgehausen P, Elsner P, Burg G. Drug-induced lichen planus. *Clin Dermatol* 1998;16:325–332.
- Giustina TA, Stewart JC, Ellis CN, Regezi JA, Annesley T, Woo TY, Voorhees JJ. Topical application of isotretinoin gel improves oral lichen planus. A double-blind study. *Arch Dermatol* 1986;122:534–536.
- Woo TY. Systemic isotretinoin treatment of oral and cutaneous lichen planus. *Cutis* 1985;35:385–386, 390–391, 393.
- Sloberg K, Hersle K, Mobacken H, Thilander H. Severe oral lichen planus: remission and maintenance with vitamin A analogues. *J Oral Pathol* 1983;12:473–477.
- Ott F, Bollag W, Geiger JM. Efficacy of oral low-dose tretinoin (all-trans-retinoic acid) in lichen planus. *Dermatology* 1996;192:334–336.
- Falk DK, Latour DL, King LE Jr. Dapsone in the treatment of erosive lichen planus. *J Am Acad Dermatol* 1985;12:567–570.
- Paslin DA. Sustained remission of generalized lichen planus induced by cyclophosphamide. *Arch Dermatol* 1985;121:236–239.
- Eisen D, Ellis CN, Duell EA, Griffiths CE, Voorhees JJ. Effect of topical cyclosporine rinse on oral lichen planus. A double-blind analysis. *N Engl J Med* 1990;323:290–294.
- Harpenau LA, Plemons JM, Rees TD. Effectiveness of a low dose of cyclosporine in the management of patients with erosive oral lichen planus. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995;80:161–167.
- Pigatto PD, Chiappino G, Bigardi A, Mozzanica N, Finzi AF. Cyclosporin A for treatment of severe lichen planus. *Br J Dermatol* 1990;122:121–123.
- Eisen D. Hydroxychloroquine sulfate (Plaquenil) improves oral lichen planus: an open trial. *J Am Acad Dermatol* 1993;28:609–612.
- Lener EV, Brieve J, Schachter M, West LE, West DP, el-Azhary RA. Successful treatment of erosive lichen planus with topical tacrolimus. *Arch Dermatol* 2001;137:419–422.
- Dereure O, Basset-Seguin N, Guihou JJ. Erosive lichen planus: dramatic response to thalidomide. *Arch Dermatol* 1996;132:1392–1393.