1. Johnson C, Pinschmidt S. *Cardiac Rehab Statistics*. Washington, DC: Cardiology Preeminence Roundtable, the Advisory Board Company, 1999.
2. Schank RC, Cleary C. *Engines for Education*. Hillsdale, NJ: Lawrence Erlbaum Associates, 1995.
3. National Institutes of Health. *Second Report of the Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II)*. Bethesda, Md: National Institutes of Health, 1993.
4. Smith SC Jr, Blair SN, Criqui MH, Fletcher GF, Fuster V, Gersh BJ, Gotto AM, Gould KL, Greenland P, Grundy SM, et al. AHA consensus panel statement. Preventing heart attack and death in patients with coronary disease. The Secondary Prevention Panel. *J Am Coll Cardiol* 1995;26:292-294.
5. Kottke TE, Battista RN, DeFries GH, Brekke ML. Attributes of successful smoking cessation interventions in medical practice. A meta-analysis of 39 controlled trials. *JAMA* 1988;259:2883-2889.
6. Lando HA, Hellerstedt WL, Pirie PL, McGovern PG. Brief supportive telephone outreach as a recruitment and intervention strategy for smoking cessation. *Am J Public Health* 1992;82:41-46.
7. Hamalainen H, Luurila OJ, Kallio V, Knuts LR, Arstila M, Hakkila J. Long-term reduction in sudden deaths after a multifactorial intervention programme in patients with myocardial infarction: 10-year results of a controlled investigation. *Eur Heart J* 1989;10:55-62.
8. Kallio V, Hamalainen H, Hakkila J, Luurila OJ. Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979;2:1091-1094.
9. Prochaska JO, DiClemente RC. Toward a comprehensive model of change. In Miller WE, Heather N, eds. *Treating Addictive Behaviors: Process of Change*. New York: Plenum, 1986:3-26.
10. Ware JE, Kosinski M, Keller SD. *SF-12: How to Score the SF-12, Physical and Mental Health Summary Scales*, 3rd ed. Lincoln: QualityMetric Inc, 1998.
11. Campbell MK, DeVellis BM, Strecher VJ, Ammerman AS, DeVellis RF, Sandler RS. Improving dietary behavior: the effectiveness of tailored messages in primary care settings. *Am J Public Health* 1994;84:783-787.
12. Bennett SJ, Hays LM, Embree JL, Arnould M. Heart messages: a tailored message intervention for improving heart failure outcomes. *J Cardiovasc Nurs* 2000;14:94-105.

Invited commentary

In the past decade, a wealth of data from both clinical trial and interventional studies has proven that clinicians who treat patients with coronary heart disease (CHD) can reduce the future risk of recurrent CHD events, reduce mortality, and improve quality of life. These treatments include the acute management of myocardial infarction and unstable angina with thrombolytics, primary angioplasty, and antithrombotic agents, as well as long-term treatment with beta-blockers, aspirin, angiotensin-converting enzyme inhibitors, and statins. Major changes in lifestyle behavior have also shown benefit, such as smoking cessation, low-fat diets, weight reduction, and regular exercise.

While most clinicians readily acknowledge the effectiveness of these treatments in the post-acute coronary syndrome (ACS) hospital setting, they are often frustrated by their patients' lack of adherence to these lifesaving therapies. The reasons for this reduced adherence and persistence to recommended lifestyle changes and drug treatments are many and complex. Most of the frequently cited reasons involve shortcomings of the health care delivery system and are potentially correctable. Some of these are lack of time to fully educate patients in the hospital setting prior to discharge, inadequate numbers of or proximity to cardiac rehabilitation programs for aftercare participation, inadequate insurance coverage for these programs, limited time and low reimbursement for private physician office education, and limited insurance coverage for prescription medications. These barriers may be overwhelming and can only be overcome by a coordinated effort from managed care organizations or large integrated health care providers. The article in this issue of *Baylor University Medical Center Proceedings* demonstrates how such a coordinated multidisciplinary approach can provide a vehicle to help patients adhere to post-ACS lifestyle and medication programs.

The Leap for Life program of the Baylor Health Care System was born out of frustration with the lack of educational opportunities for CHD patients who had no access to traditional phase II cardiac rehabilitation programs. The program's subsequent success allowed it to expand its mission to include all patients discharged from the system's hospitals following a myocardial infarction or cardiac catheterization. The authors describe their experience with 152 participants over a 12-month period, focusing on patient perceptions of goal achievement, satisfaction, and hospital readmission. Three quarters of the participants felt that the program helped them meet all or some of their goals, and 100% of them thought the workshops met their educational needs.

The participants' success rate for achieving their own preset goals is satisfying, particularly for smoking cessation and control of blood pressure and diabetes. On the other hand, the program shows us how difficult some lifestyle changes can be, specifically reducing weight and stress. Although not described, it would be important to measure medication adherence and persistence and quality of life. Ultimately, determining the impact that goal achievement has on recurrent CHD events and hospital admissions would allow an analysis of the cost-effectiveness of the Leap for Life program. The Baylor Health Care System has initiated an important educational program for the post-ACS patient. Its initial success should be applauded and should encourage other managed or integrated health care groups to follow suit. Expanding the outcomes measures of the Leap for Life program will most likely prove that these educational initiatives are not only cost saving, but life saving as well.

—PETER H. JONES, MD

Section of Atherosclerosis and Lipoproteins,
Baylor College of Medicine