The nursing profession's coming of age

The end of medicine as a profession

EDITOR—Young describes the coming of age of the nursing profession.¹ All professions and occupations are human creations and therefore are subject to change. With the widening of roles of many professions, it becomes harder to define what a doctor can do that can't be done by an appropriately trained member of another profession. Perhaps major surgery and certifying cause of death are all that remains. With the increasing difficulty of getting junior doctors to do anything that may be regarded as work (rather than education) we doctors may be engineering the abolition of our profession, and we may move from distinct professions to practitioners from varied backgrounds being trained for a particular function.

David L Bisset *consultant histopathologist*
Royal Bolton Hospital, Bolton BL4 0JR
david.bisset@rbh.nhs.uk

Competing interests: I am a registered medical practitioner.

¹ Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

The death of nursing, the dumbing down of medicine

EDITOR—Young claims that the distinction between medicine (diagnosis and cure) and nursing (care) has become increasingly blurred.¹ No it has not—patients still have the same illnesses and still require the same care as they did before. It is the abandonment of nursing by modern nurses that is new, to the detriment of health care in this country.

"The only aim to consider is that we respond safely, sensitively, and effectively to our patients' needs."¹ Exactly. So can we please get back to basics and have trained nurses to nurse and trained doctors to doctor. Instead we have care assistants playing at being nurses, nurses playing at being doctors, doctors playing at being managers, and managers just playing.

Here is a radical idea. Why not have doctors who are trained to doctor, do the doctoring, and nurses who are trained to nurse, do the nursing. That way the patients might actually get "the best care possible" that the Department of Health propaganda unit keeps telling us about.

Robbie S Coull *general practitioner*
Inverness IV1 1LD
locum@coull.net

Competing interests: None declared.

¹ Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Contribution of "maxi" nurses is exaggerated

EDITOR—Like many "maxi" nurses, Young seems to have an exaggerated notion not only of her worth but also of the contribution she makes.¹ She trots out the cliché that nurses are better than doctors at communication and are more "holistic." If I saw the numbers of patients the "maxi" nurses I see at work then I too would be a fountain of compassion and empathy with all the time in the world, because I would only be seeing a third of the patients I now assess.

Susan Duncan *consultant neurologist*
Hope Hospital, Salford M6 8HD
susan@susanduncan.co.uk

Competing interests: SD is a registered medical practitioner.

¹ Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Plus ça change

EDITOR—Some of us who worked in hospitals may remember the time when general practitioners boasted of their superiority to specialists because of the fact that their care and treatment centred on the whole patient. It is amusing now to find Young claim that a consultation with a nurse is preferred to one with a general practitioner for exactly the same reason.¹

Memo G S Spathis *retired physician*
St Helier Hospital, Carshalton, Surrey SM5 1AA
spathis@doctors.org.uk

Competing interests: None declared.

¹ Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

My experience of nurse practitioners

EDITOR—Having worked at some primary care centres managed by primary care trusts promoting the use of nurse practitioners,¹ I am worried about the standard of care these practitioners deliver for some patients with

undiagnosed illnesses. For example, they seem routinely not to clinically examine patients with headaches or abdominal pain. These nurse practitioners are not students; they are getting extended prescribing rights and are training other nurses to become nurse practitioners. It truly is the blind leading the blind.

Anjan Chakraborty *general practitioner locum*
London
anjan@blueyonder.co.uk

Competing interests: AC is a general practitioner.

¹ Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Not as successful as it looks

EDITOR—Young says that some practices are even nurse led—it is the nurses who employ the general practitioners, and patients are formally registered with the practice, not the doctor.¹ What she neglects to point out is that one of these flagship nurse led practices has had to be taken over by the primary care trust, the nurses' contract being terminated. This was apparently because of the numerous patients' complaints about the service with which they were being provided.

The fundamental difference between doctors and nurses is that doctors have a holistic view of all patho-physiology and at least a decade of training in diagnosis and diagnostic uncertainty. Nurse practitioners do not have the depth and breadth of training to manage all aspects of patients' care on their own. Pretending that nurses can be doctors is not helpful. Instead of spending 12 years learning to be a specialist in one area, perhaps Young would have better spent the time training to be a doctor and then a general practitioner, so as to be able to be a generalist with unlimited independent practice.

Trefor J Roscoe *general practitioner*
Sothall Medical Centre, Sheffield S20 5JX
Trefor@nhs.net

Competing interests: TJR is a registered medical practitioner.

¹ Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Sign of the times

EDITOR—Support for extended powers to nurses is limited and comes from short-sighted consultants who seem to have forgotten the worth of repetitive tasks of the nature presently assigned to nurse practitioners in shaping the consultant of tomorrow.¹ Extending the role of nurses to include

surgical care makes a mockery of a surgical training scheme that has believed in apprenticeship after a period of intense study in medical school.

Vinay P Rao *research fellow*
Mayo Clinic, Rochester, MN 55905, USA
vinayprao@yahoo.com

Competing interests: VPR is a medical practitioner and has been a junior doctor in surgical training in the NHS.

1 Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

A difference in cultures

EDITOR—There may also be a natural self-selection process in this debate in terms of skills, etc—most people who choose nursing are fairly good communicators and most people who choose surgery like being dextrous.¹

Nurses are also arguably better than doctors at protocol driven decision making. Doctors, however, manage risk as part of their core training—making the diagnosis and managing it thereafter—which nurses historically have not.

I think it is great that nurses learn more skills and maximise their potential with MSc degrees, diplomas, procedures, etc, but I also think we should all do what we do best. Perhaps medical educators need to work out a way that nurses can be fast-tracked into becoming doctors. This would be far better than what we have now: ambiguity and even the blurring of roles.

Ayan S Panja *general practitioner*
The Greens Medical Practice, 96 Umfreville Road, London N4 1TL
ayanpanja@hotmail.com

Competing interests: ASP is a general practitioner.

1 Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Separate but complementary

EDITOR—In my several years of postgraduate training I have had the privilege to work with, and learn from, numerous nurses, male and female.¹ They never pretended to be anything else than their title implied, and they expected me to authorise their suggestions about patient management, even though their experience (particularly in specialist units) was longer than my own. Through this collaboration I have come to appreciate the complementary skills that diverse professions can bring into the care (and occasionally the cure) of the whole person. Blurring the roles can never accomplish as much as mutual respect and cooperation between varied skills.

Anthony Papagiannis *respiratory physician*
St Luke's Hospital, Thessaloniki, Greece
antpap@the.forthnet.gr

Competing interests: None declared.

1 Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Doctors can use their skills to the maximum

EDITOR—Nurse practitioners do not seek to be doctors, but surely it is more cost effective to have an experienced nurse deal with the minor self limiting illnesses that make up much of general practice and allow doctors to use their skills to the maximum.¹ Having worked in this practice for 16 years as a practice nurse before training to be a nurse practitioner, I see it as a natural progression to be able to deal with anything presented to me. I think also the confusion between someone calling themselves a nurse practitioner and working to protocols and someone who has trained properly can finally be resolved, with the Nursing and Midwifery Council acknowledging the role.

Sandra Magennis *nurse practitioner*
Idle Medical Centre, Bradford BD10 8RU
sandra.magennis@bradford.nhs.uk

Competing interests: Ghislaine Young was SM's mentor when she trained to be a nurse practitioner.

1 Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

A team effort required?

EDITOR—My work as a nurse practitioner is primarily several duties previously carried out by a senior house officer: routine clerking of preoperative patients, post-operative reviews, and assessment of ophthalmic emergencies and minor surgical procedures.¹ Training has been in the form of academic postgraduate training and one to one teaching with several (extremely patient) consultant colleagues. I have not tried to usurp the role of the medical practitioner, and I hope that I draw on knowledge from the domains of both medicine and nursing during my clinical interactions. Indeed, much of the impetus for the role came not from any desire by ophthalmic nurses to be backdoor doctors, but on the suggestion of the deanery that training grade doctors should not be trapped in routine work at the expense of more esoteric clinical experience.

Mark Davies *nurse practitioner in ophthalmology*
North Glamorgan NHS Trust, CF47 9DT
mark.davies@nglam-tr.wales.nhs.uk

Competing interests: MD is a nurse practitioner.

1 Young G. The nursing profession's coming of age. *BMJ* 2005;331:1415. (10 December.)

Rise and demise of the hospital

Managing hospital inpatient care

EDITOR—Black calls for nurses to transform our hospitals.¹ This will not happen without a major change in the clinical culture that pervades hospital practice.

Observational studies show inadequate communication between doctors and other clinical staff.² Attempts to foster collaboration are undermined by differential power and status, lack of interprofessional socialisation, and inadequate attention to team building. The problem is intensified by the pressure on senior nurses to enhance their managerial roles. Time spent on management too often leaves them with inadequate time in which to provide the leadership necessary to maintain standards of clinical care.³ This saps morale and may play an important part in the high turnover of nurses in most of our hospitals. Meanwhile senior nurses seek career progression through management or by becoming nurse specialists rather than continuing as ward sisters, a role that was once regarded as the apex of the nursing profession.

No one would wish to return to the days when "senior nurses made their hospital ward their home," but there seems to be an overwhelming need for training clinical nurse consultants who know their patients, who define the standards of nursing on their wards and who play a key role in the training of junior doctors. And they should be paid accordingly—the top of the specialist nurse scale.

Since the turn of the century the training of medical students has changed dramatically with the emphasis on professional development in accord with the directives of the General Medical Council.⁴ Is it not time for the nursing profession to rethink its aims and objectives, and to improve the quality of clinical training?

Graham Neale *visiting professor*
g.neale@imperial.ac.uk

Sisse Olsen *research fellow, Clinical Safety Research Unit*
Department of Biosurgery and Surgical Technology, Imperial College, St Mary's Hospital, London W2 1NY

Competing interests: None declared.

1 Black N. Rise and demise of the hospital: a reappraisal of nursing. *BMJ* 2005;331:1394-6. (10 December.)

2 Reeves S, Lewin S. Interprofessional collaboration in the hospital: strategies and meanings. *J Health Serv Res Policy* 2004;9:218-25.

3 Johns C. Clinical supervision as a model for clinical leadership. *J Nurs Manage* 2003;11:25-34.

4 General Medical Council. *Tomorrow's doctors*. London: GMC, 2003.

Backwards is the new forward: time for the heritage hospital

EDITOR—May I suggest as a start to Black's transformations¹:

- Reinstatement of a strictly pyramidal shape to the ward management structure, in which the sister knows every patient under her care
- Replace teams with a flexible "whole ward" approach to allocating nursing tasks
- Restore a more formal and, by inference, professional relationship between staff when in the clinical area—reconsider, for example, whether it is appropriate to use first names either for staff or for patients
- Treat nursing time as our most valuable resource—not to be squandered on writing off-duty rotas or over-elaborate nursing

records but to be spent in the company of their patients

- Reinstate the consultant ward round as the focus for the business of the week.

My plans for green tiling, shiny floors, and Sunday afternoon matron's ward round are under wraps until I have had a chance to pilot them—after all, we have to have a good evidence base.

Richard Neil Harrison *consultant physician*
University Hospital of North Tees, Stockton
TS19 8PE
richard.harrison@nth.nhs.uk

Competing interests: None declared.

- 1 Black N. Rise and demise of the hospital: a reappraisal of nursing. *BMJ* 2005;331:1394-6. (10 December.)

Is methadone too dangerous for opiate addiction?

Issue is one of toxicity v acceptability

EDITOR—We disagree with Luty et al's suggestion that buprenorphine should replace methadone.¹ Doctors and patients need different treatment options in different phases of treatment, and buprenorphine does not suit everyone. Since January 2004, 54% of the patients who started maintenance treatment with us chose buprenorphine initially. However, 64% of them asked to switch to methadone within two months because buprenorphine did not suit them.

Ensuring truly supervised consumption for slowly dissolving sublingual tablets is difficult. We do not discount the risks of recreational illicit methadone use, but illicit buprenorphine is not hazard-free. The risk of respiratory depression might be less, but the likelihood of injecting is greater and there is still a risk of pulmonary oedema.

Of the 167 methadone related deaths in 2003, 75% were caused by multidrugs. Overall, opiate overdose deaths fell by 21% to 1040 compared with 2002.² The proportion of polydrug deaths could have been even higher because contributory drugs are often under-reported.³ Postmortem blood concentrations of methadone are hard to interpret because it is lipophilic, and concentrations can increase up to fourfold by redistribution after death whereas morphine concentrations show relatively little change.⁴ Pirnay et al showed the difficulty in determining methadone's role in polydrug deaths.¹ In 30% of methadone related deaths other substances were also present at toxic concentrations, whereas some of the highest methadone concentrations were found in cases of carbon monoxide poisoning and a firearm homicide.¹

The highest drug related death rate in 2003 was in Brighton,² where methadone provision was considered to be poor. Only a minority of addicts currently seek treatment. Rather than reducing the treatment menu, treatments should be made as accessible and attractive as possible. Methadone still has the

best outcomes on meta-analysis.⁵ Buprenorphine certainly has a place but so have dihydrocodeine and morphine.

Adam Bakker *general practitioner*
Vanessa Sibanda *clinical nurse specialist (substance misuse)*
Lisson Grove Health Centre, London NW8 8EG
bakker@nhs.net

Competing interests: None declared.

- 1 Luty J, O'Garra C, Sessay M. Is methadone too dangerous for opiate addiction? *BMJ* 2005;331:1352-3. (10 December.)
- 2 Ghodse H, Schifano F, Oyefeso A, Bannister D, Cobain K, Dryden R, et al. *Drug-related deaths as reported by participating procurator fiscal and coroners*. London: St George's Hospital Medical School, 2004. (International Centre for Drug Policy Reports 11 and 13.)
- 3 Zador D, Kidd B, Hutchinson S, Taylor A, Hickman M, Fahey T, et al. *National Investigation into drug related deaths in Scotland, 2003*. Edinburgh: Scottish Executive Social Research, Substance Misuse Research Programme, 2005.
- 4 Drummer OH. Post mortem toxicology of drugs of abuse. *Forens Sci Int* 2004;142:101-13.
- 5 Mattick RP, Kimber J, Breen C, Davoli M. Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane Database Syst Rev* 2003;(2):CD002207.

Methadone is still needed in addiction treatments

EDITOR—Luty et al's enthusiasm for buprenorphine over methadone risks losing baby and bath water.¹ Some patients fare well on one drug yet poorly on the other, so a choice is needed, as in depression, arthritis, or hypertension. The references cited do not support the assertion that buprenorphine is at least as effective as methadone in maintenance treatment. Most randomised control trials have showed adequate methadone doses (60-100 mg/day) to be associated with modestly higher retention rates and lower illicit opiate use than buprenorphine, with low dose methadone (20-30 mg/day) clearly giving poorer results than adequate doses.²

What randomised trials can tell us about dose-response relations in methadone maintenance is limited, and the use of randomised trials of non-equivalent medications has ethical limitations. Furthermore, a finding of "no significant difference" is often loosely interpreted to mean clinical equivalence, which it is not.³ There is other clinical evidence of a dose-response relation in methadone maintenance treatment, extending to doses above 100 mg/day.⁴

Luty et al point out the risks of unsupervised dispensing of methadone. However, they assume that reducing availability of methadone will reduce overdoses, despite strong contrary evidence that increased methadone availability actually reduces overdose rates.⁵

This is not an either/or situation. Better outcomes in addiction treatments are likely to be achieved by increasing the range of treatments available. The answer is not for methadone to be replaced by a more expensive and sometimes less effective alternative but for appropriate supervision of dispensing of opioid replacement treatment, with

take-home doses for people who have demonstrated stability in treatment.

Andrew Byrne *addiction physician*
Richard Hallinan *addiction physician*
75 Redfern Street, Redfern, NSW 2016, Australia
ajbyrne@ozemail.com.au

Competing interests: AB charges a fee for dispensing maintenance opioids in his surgery.

- 1 Luty J, O'Garra C, Sessay M. Is methadone too dangerous for opiate addiction? *BMJ* 2005;331:1352-3. (10 December.)
- 2 Mattick RP, Kimber J, Breen C, Davoli M. Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane Database Syst Rev* 2004;(3):CD002207.
- 3 Ling W, Wesson DR. Clinical efficacy of buprenorphine: comparisons to methadone and placebo. *Drug Alcohol Depend* 2003;70:S49-57.
- 4 Hallinan R, Ray J, Byrne A, Agho K, Attia J. Therapeutic thresholds in methadone maintenance treatment: A receiver operating characteristic analysis. *Drug Alcohol Depend* 2005 Jul 15 (Epub ahead of print).
- 5 Grönbladh L, Öhlund LS, Gunne LM. Mortality in heroin addiction: impact of methadone treatment. *Acta Psychiatr Scand* 1990;82:223-7.

Are GPs who specialise in dermatology effective?

Be wary of generalising research conclusions

EDITOR—The conclusions of the study reported by Salisbury et al cannot be generalised.¹ Some primary care trusts seem to see the appointment of general practitioners with a special interest at best as a "quick fix" for their waiting times problems and at worst as an alternative to the proper provision of secondary care dermatology services.

Appointing such specialists is not the only way in which dermatology services may be improved. The Department of Health's dermatology workforce group has concluded that the traditional model, with the general practitioner as the "gatekeeper" to intermediate or secondary care services is inimical to optimal care of patients with long term skin diseases. It sees the key to sustainable service improvement as being the separation of diagnosis from treatment.

All patients should have rapid access to diagnosis of their skin disease and that diagnosis should be undertaken at whatever level in the healthcare system is necessary. General practitioners should be able to diagnose the common skin disorders in their characteristic forms, but it may be necessary for patients to be referred promptly to intermediate or secondary care and, thereafter, even to tertiary care for the diagnosis of skin diseases that are beyond the general practitioner's experience.

Once properly diagnosed, and while some complex or unresponsive conditions will still require active treatment and management in primary or secondary care, most common inflammatory skin diseases can be managed by patients themselves, provided they are properly supported by a (probably nurse led) inflammatory skin disease clinic in primary care. A model of

this sort offers the double benefit of improved patient satisfaction and substantial reduction of the burden of skin disease in primary care. Most of the 15 pilot studies funded by the Department of Health's Action on Dermatology programme concerned such clinics, which were shown to be effective and popular with patients.

Peter M Lapsley *chief executive*
Skin Care Campaign, London N19 5NA
plapsley@eczema.org

Competing interests: PL is deputy chair of the Department of Health's dermatology workforce group.

1 Salisbury C, Noble A, Horrocks S, Crosby Z, Harrison V, Coast C, et al. Evaluation of a general practitioner with special interest service for dermatology: randomised controlled trial. *BMJ* 2005;331:1441-6. (17 December.)

Result for teamwork and planning is positive

EDITOR—Lapsley (previous letter) is right that it is inappropriate to extrapolate the positive findings from the study reported by Salisbury et al to all schemes¹: they have not all been set up in the same way or as well. The question "What is a general practitioner with a special interest?" needs to be asked and answered. Training and accreditation for these doctors are not as yet nationally defined or regulated.

In Southampton we approached the problem of "inappropriate" cases being referred to dermatologists (six years' dermatology training) by general practitioners (six days' dermatology training) by developing intermediate level doctors with six months' (equivalent) dermatology training. General practitioners with a special interest would not be necessary if all general practitioners had much more training in dermatology and there were more dermatologists, but they are seen as a realistic compromise for now.

The issue of calculating cost effectiveness is contentious. The manager of my primary care trust told me that my cost per case is £38, as opposed to £250 per case for the hospital. I tend to question how this was calculated, and I am surprised at the high cost per case in the paper.

Stephen F Hayes *general practitioner (with a special interest in dermatology)*
Bitterne Walk In Centre, Southampton
stephen.hayes007@onetel.net

Competing interests: SH was involved in developing a general practitioner with a special interest scheme in Southampton, works as a general practitioner with a special interest in dermatology, is paid for teaching dermatology to general practitioners and nurses, and is a member of the Primary Care Dermatology Society committee.

1 Salisbury C, Noble A, Horrocks S, Crosby Z, Harrison V, Coast C, et al. Evaluation of a general practitioner with special interest service for dermatology: randomised controlled trial. *BMJ* 2005;331:1441-6. (17 December.)

High cost of general practitioners with special interests is a fallacy

EDITOR—The high cost of the service in the articles by Salisbury et al and Coast et al

resulted from its failure to attract enough patients to make it economic.^{1,2} A proper, community based skin service relies on accuracy, speed, and a rapid reply to the referring general practitioner, as well as being embedded in the community. I set up such a service in Ealing in 2000, after passing the diploma in practical dermatology, being a clinical assistant, and doing a skin surgery course. It was fully accredited by the British Association of Dermatology and had the support of the local dermatologists.

Local general practitioners referred patients directly to me by fax. I did three clinics, at three hours each, per fortnight. In 14 months I saw 658 new patients and 498 follow-ups, a total of 1156 consultations. I typed the replies and faxed the letter to the general practitioner that night. I saw patients within 2.5 weeks, had a rate of non-attendance of 5%, and referred 3% to hospital. I also did over 100 minor operations. I had no administrative or nursing support at any time, but I had a typist for six months in the four years that the service ran.

My costs were about £10 per patient (£155 a clinic), £80 per operation, £30 a week to type all the clinic letters (up to 25), and minimal costs to rent rooms. The total cost was about £27 000 for 1156 patients, £23.35 a patient, about a 10th of the costs in the Salisbury model (£207.91) and about a fifth of the consultant costs.

The papers by Salisbury et al and Coast et al paint a very different picture to the work most general practitioners with special interests do, as well as the costs pertaining to such a service. The conclusions of the articles are suspect; the reality is different.

Cornelius J Crowley *general practitioner principal*
Queens Walk Surgery, London W5 1TP
neil.crowley@gp-e85057.nhs.uk

Competing interests: CJC was a general practitioner with a special interest in Ealing until he resigned in March 2005. He is a member of the Primary Care Dermatology Society.

- 1 Salisbury C, Noble A, Horrocks S, Crosby Z, Harrison V, Coast C, et al. Evaluation of a general practitioner with special interest service for dermatology: randomised controlled trial. *BMJ* 2005;331:1441-6. (17 December.)
- 2 Coast J, Noble S, Noble A, Horrocks S, Asim O, Peters TJ, et al. Economic evaluation of a general practitioner with special interest led dermatology service in primary care [with commentary by M Roland]. *BMJ* 2005;331:1444-9. (17 December.)

Cost calculations were impenetrable

EDITOR—Salisbury et al have provided evidence that community dermatology, delivered by a general practitioner with a special interest, is accessible, well liked by patients, and effective.¹ This comes as no surprise to those of us working in this service. However, I was unable to understand how Coast et al found it to be nearly

twice as expensive for the NHS as hospital outpatient care.² I cannot refute the findings of this particular study, as the paper is impenetrable to all but health economists; however, I think it important to point out that dermatology posts for general practitioners with a special interest are not all the same.

I work for Norwich and South Norfolk primary care trusts in two separate clinics. Not only do the doctors have a list, but the specialist dermatology nurse has her own clinic running concurrently. She is able to treat a number of dermatological conditions within a limited list and with the support of

her medical colleagues. Norwich primary care trust is confident that significant savings are being made through the appropriate use of a general practitioner service.

My remit, apart from the clinical dermatology service, is to teach undergraduates at University of East Anglia medical school in the university and in the community clinics and to provide feedback sessions to my general practitioner colleagues. Recently negotiations started with a nascent, practice based commissioning group, to provide teaching sessions directly to their general practitioners.

A value added part of a general practitioner with special interest service for dermatology delivered in the community is the relation achieved with local general practitioners and medical students. Whether this relationship will have a beneficial effect on inappropriate referrals is too early to say, but it is unreasonable to use "economic evaluation" of a single model as a stick to beat us all with.

Simon F Cooper *general practitioner principal*
Theatre Royal Surgery, Dereham NR19 2QG
simon.cooper@nhs.net

Competing interests: SFC is a general practitioner with a special interest in dermatology for South Norfolk and Norwich primary care trusts.

- 1 Salisbury C, Noble A, Horrocks S, Crosby Z, Harrison V, Coast C, et al. Evaluation of a general practitioner with special interest service for dermatology: randomised controlled trial. *BMJ* 2005;331:1441-6. (17 December.)
- 2 Coast J, Noble S, Noble A, Horrocks S, Asim O, Peters TJ, et al. Economic evaluation of a general practitioner with special interest led dermatology service in primary care [with commentary by M Roland]. *BMJ* 2005;331:1444-9. (17 December.)

bmj.com

We select the letters for these pages from the rapid responses posted on bmj.com within five days of publication of the article to which they refer.

Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.