JAMA's policy on industry sponsored studies

EDITOR—The editorial by Rothman and Evans, that accused the JAMA editorial policy governing industry sponsored studies of being “unfair and absurd,” contains numerous errors and misconceptions.^{1,2}

The title of the editorial incorrectly implies that the JAMA policy applies only to clinical trials. An independent statistical analysis is required for any type of industry sponsored study in which the data analysis is conducted only by statisticians employed by the research sponsor. Rothman and Evans also incorrectly say that the policy requires authors to hire an academic statistician before their submission will be considered by JAMA. The policy clearly states that studies in which the data analysis has been conducted only by statisticians employed by the company sponsoring the research will not be accepted for publication in JAMA. Failure to have the independent analysis completed at the time the paper is submitted does not preclude initial evaluation and review of the paper. The authors also contend the policy may require arbitrary distinctions, such as defining academic statistician. We clearly indicated that the independent analysis must be conducted by statisticians at an academic institution, such as a medical school, academic medical centre, or government research institute. This analysis, which is conducted using the entire raw dataset, serves to corroborate independently the analysis performed by statisticians employed by the sponsor, and by virtue of involvement of an academic centre, provides an additional layer of institutional oversight.

Ironically, this blatant and unbalanced criticism of the JAMA policy was published just two days after the *New England Journal of Medicine* issued its expression of concern about the VIGOR study, which directly resulted from concerns about data integrity, analysis, and reporting in a major industry sponsored study.³ This example, as well as numerous other serious scientific and ethical lapses involving industry and industry sponsored studies, such as incomplete reporting of data on celecoxib,⁴ suppression of studies of paroxetine,⁵ and delayed reporting of design defects in implantable cardioverter defibrillators,⁶ have resulted in doctors and the public now having an unprecedented lack of trust and confidence in manufacturers of drugs and devices.

We believe our policy is necessary and sound. What we find unfair and absurd is that some companies and research sponsors

will resort to tactics that manipulate clinical research data and misrepresent scientific information; that some commentators, such as Rothman and Evans, apparently do not fully acknowledge the magnitude and implications of these problems; and, surprisingly, that a journal like the *BMJ* would criticise another journal's policy on such a pressing issue without properly vetting the material for accuracy.

To help ensure confidence in industry sponsored studies, JAMA will continue to require an author who is independent of the sponsor to take responsibility for the completeness and integrity of all the data in the study and will continue to require an independent statistical analysis for studies in which the data analysis was conducted only by statisticians employed by the sponsor.

Phil B Fontanarosa executive deputy editor

Phil.Fontanarosa@jama-archives.org

Catherine D DeAngelis editor-in-chief

JAMA, 515 N State Street, Chicago, IL 60610, USA

Competing interests: None declared.

1 Rothman KJ, Evans S. Extra scrutiny for industry funded trials. *BMJ* 2005;331:1350-1. (10 December.)

2 Fontanarosa PB, Flanagan A, DeAngelis CD. Reporting conflicts of interest, financial aspects of research, and role of sponsors in funded titles. *JAMA* 2005;294:110-1.

3 Curfman GD, Morrissey S, Drazen JM. Expression of concern: Bombardier et al. “Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis,” *N Engl J Med* 2000;343:1520-8. *N Engl J Med* 2005;2813-4.

4 Hrachovec JB, Wright JM. Reporting of 6-month vs 12-month data in a clinical trial of celecoxib. *JAMA* 2001;286:2398.

5 Gibson L. GlaxoSmithKline to publish clinical trials after US lawsuit. *BMJ* 2004;328:1513.

6 Maisel WH. Safety issues involving medical devices: implications of recent implantable cardioverter-defibrillator malfunctions. *JAMA* 2005;294:955-8.

US flu mortality estimates are based on solid science

EDITOR—Doshi implies that the use of statistical models to estimate flu related mortality is inappropriate.¹ Not so. Epidemiologists rely on statistical models because the International Classification of Diseases (ICD) code for influenza (ICD9 487) severely undercounts the true number of flu related deaths. Doshi also implies that the Centers for Disease Control and Prevention (CDC) deliberately exaggerates flu mortality for the benefit of the pharmaceutical industry, while other scientists stand by, meek and mute. The *BMJ*'s decision to publish Doshi's commentary with no counterpoint from an

expert in the field served only to sow unnecessary confusion.

We need statistical models partly because flu is rarely confirmed in the laboratory and partly because flu brings about many more deaths from secondary bacterial pneumonia or exacerbations of underlying chronic diseases than from primary flu pneumonia. The classic Serfling modelling approach measures the total seasonal number of flu related deaths as the excess mortality above an expected winter baseline of all winter deaths.^{2,3} Regression models guided by virus surveillance data have been developed more recently.^{4,5} These modelling approaches yield similar estimates when applied to the same mortality data for the same range of years (table). The deaths identified by the models only occur when flu causes epidemics² and are most numerous for flu seasons dominated by severe A(H3N2) viruses.³ The important point is that the deaths attributed to flu by these models are flu related in that they would not have occurred without flu.

Why, Doshi asks, are the newer CDC estimates much higher than the older estimates of seasonal flu related deaths? The disparity between older and newer estimates is simply a consequence of various authors studying different time periods over decades when the elderly population simultaneously grew and aged (table); specifically, the proportion of elderly people over 85 had quadrupled by the end of the 1990s, which led to increasing mortality estimates because the risk of dying from flu increases exponentially in the last decades of life.³

And why, Doshi asks, is the estimate of 34 000 deaths for the 1968 pandemic lower than the current CDC estimate for an average season? One cannot compare crude mortality estimates across decades because

Average seasonal estimates of flu related deaths for specific time periods, based on applying various models to all US deaths

	1972-92	1976-99	1990-9
Crude estimates of all cause excess mortality			
Former CDC Serfling model ²	20 000	—	—
Current CDC virus-guided model ^{4*}	—	34 470	51 203
NIH Serfling model ³	24 400	33 400	48 700
Annual regression model ⁵	—	—	51 900
Age adjusted estimates of all cause excess mortality†			
NIH Serfling model ³	19 800	23 600	30 800

*Crude all cause excess deaths from tables 2 and 3 in Thompson et al.⁴

†Carefully adjusted for population ageing but not for increased circulation of virulent A(H3N2) viruses in 1990s—a factor explaining much of residual increase compared with previous time periods.³

of the ageing of the population; what makes the mild 1968 pandemic stand out is the shift in mortality to younger people—nearly half the 1968 flu related deaths occurred in people under 65, compared to around 10% of deaths in recent seasons.

Lone Simonsen senior epidemiologist
L.Simonsen@niaid.nih.gov

Robert Taylor contract analyst

Cecile Viboud visiting fellow

Jonathan Dushoff research scientist

Mark Miller director, Division of Epidemiology and

International Population Studies

National Institutes of Health, Bethesda, MD 20892, USA

Competing interests: None declared.

- 1 Doshi P. Are US flu death figures more PR than science? *BMJ* 2005;331:412. (10 December.)
- 2 Simonsen L, Clarke MJ, Williamson GD, Stroup DF, Arden NH, Schonberger LB. The impact of influenza epidemics on mortality: introducing a severity index. *Am J Public Health* 1997;87:1944-50.
- 3 Simonsen L, Reichert TA, Viboud C, Blackwelder WC, Taylor RJ, Miller MA. Impact of influenza vaccination on seasonal mortality in the US elderly population. *Arch Intern Med* 2005;165:265-72.
- 4 Thompson WW, Shay DK, Weintraub E, Brammer L, Cox W, Anderson LJ, et al. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* 2003;289:179-86.
- 5 Dushoff J, Plotkin JB, Viboud C, Earn DJD, Simonsen L. Mortality due to influenza in the United States—an annual regression approach using multiple-cause mortality data. *Am J Epidemiol* 2006;163:181-7.

Stockpiling of raw material for oseltamivir is not justified

EDITOR—Mudur reports that Roche has granted a sublicense for the production of the antiviral drug oseltamivir (Tamiflu) to an Indian company and points out that the supply of raw material, shikimic acid (extracted from the Chinese spice star anise *Illicium verum*), is limited.¹

We did not find any effect of shikimic acid (1 µmol/1-10 pmol/1) on the release of two predictors of antiviral activity—cytokines interleukin-8 (IL8) and interleukin-6 (IL6)²—in human peripheral blood mononuclear cells or when the mononuclear cells were stimulated by lipopolysaccharide.³ We also tested quercetin, another natural polyphenol found with shikimic acid in plants such as *Hypericum laricifolium* (romerillo) and *Cyclopia subternata* (honeybush tea)—traditional herbal remedies used in Ecuador and South Africa. Only the concomitant presence of the two compounds (concentration 10 pmol/1) produced a nearly twofold increase in IL8 (9440 (SD 778) pg/ml *v* 5657 (247) pg/ml) and an 18-fold increase in IL6 (3959 (2479) *v* 216 (52) pg/ml) compared with controls.

Other less well known, but more common sources of shikimic acid are grape juice and white wine. The analysis of six Italian and four Austrian white wines yielded concentrations of shikimic acid of 15-27 mg/l and quercetin up to 10.6 mg/l.

Alberto A E Bertelli research physician
Department of Human Morphology, University of Milan, via Mangiagalli 31, 20133 Milan, Italy
studiojh@usa.net

Competing interests: None declared.

- 1 Mudur G. Indian firm to produce oseltamivir for developing countries. *BMJ* 2006;332:5. (7 January.)
- 2 Barak V, Halperin T, Kalickman I. The effect of sambucol, a black elderberry-based, natural product, on the production of human cytokines. I. Inflammatory cytokines. *Eur Cytokine Netw* 2001;12:290-6.
- 3 Bertelli AA, Migliori M, Panichi V, Longoni B, Origlia N, Ferretti A, et al. Oxidative stress and inflammatory reaction modulation by white wine. *Ann N Y Acad Sci* 2002;957:295-301.

Dummies and SIDS

Causality has not been established

EDITOR—The relative risk for associated factors should not be confused with the attributable risk for causative factors. Li et al express the risk associated with sudden infant death syndrome (SIDS) and use of a dummy (pacifier) during sleep as a 90% reduction in risk, which could be potentially misunderstood and brings unwarranted media attention to what otherwise is a very interesting paper.¹

Any calculation of attributable risk would be limited to the prevalence of exposure, which for dummy users in the Californian infant population would be 23% (based on the control data). However, projecting risk reduction or quantifying attributable risk in this study is inappropriate. As the authors point out in the discussion, a causal effect between dummy use and SIDS has not been established. The discrepancy in dummy use has already been reported in previous larger studies,²⁻⁴ with better ascertainment levels providing a more cautious interpretation of the data.

Researchers in SIDS still need to determine what the protective mechanism might be, if it exists, or whether dummy use is a marker for something else. The discrepancy in dummy use between SIDS cases and age matched control infants for the last sleep does not seem to be as marked among routine users; this may indicate that infants are at greater risk of SIDS if they routinely use a dummy but have not been given their dummy on a particular night. Any discussion of potentially modifiable factors and impact on infant care practices is better served by a review on the subject taking into account a more detailed study of infant sleep behaviour and any detrimental effect of the exposure outside this research area.⁵

Peter S Blair research fellow
p.s.blair@bris.ac.uk

Peter J Fleming professor of infant health and developmental physiology
Institute of Child Life and Health, University of Bristol, Bristol BS2 8AE

Competing interests: None declared.

- 1 Li D-K, Willinger M, Petitti DB, Odouli R, Liu L, Hoffman H. Use of a dummy (pacifier) during sleep and risk of sudden infant death syndrome. *BMJ* 2006;332:18-21. (7 January.)
- 2 Altman DG, Bland JM. Statistics notes: interaction revisited: the difference between two estimates. *BMJ* 2003;326:219.

- 2 Mitchell EA, Taylor BJ, Ford RP, Stewart AW, Becroft DM, Thompson JM, et al. Dummies and the sudden infant death syndrome. *Arch Dis Child* 1993;68:501-4.
- 3 Fleming PJ, Blair PS, Pollard K, Ward Platt M, Leach C, Smith I, et al. Pacifier use and sudden infant death syndrome: results from the CESDI/SUDI case control study. *Arch Dis Child* 1999;81:112-116.
- 4 McGarvey C, McDonnell M, Chong A, O'Regan M, Matthews T. Factors relating to the infant's last sleep environment in sudden infant death syndrome in the Republic of Ireland. *Arch Dis Child* 2003;88:1058-64.
- 5 Mitchell EA, Blair PS, L'Hoir MP. Should pacifiers be recommended to prevent SIDS? *Pediatrics* (in press).

Low response rates generate considerable uncertainty

EDITOR—Li et al found that in infants who died from sudden infant death syndrome (SIDS) the reported use of a dummy (pacifier) in the index sleep was 4.1%, in the 50% of eligible mothers who agreed to take part.¹ The use of a dummy in the other 50% of eligible infants with SIDS is unknown. In the control infants the reported use of a dummy in the last sleep before interview was 23.6%, in the 40% of eligible mothers who agreed to take part. The use of a dummy in the other 60% of control infants is also unknown.

Thus, potentially the overall dummy use in all the cases was between 2% and 54% and in all the controls between 9% and 69%. None of the confidence intervals reported in the paper reflects uncertainty generated by those who declined to take part. Because of the low response rates, unadjusted odds ratios of any where between 0.01 and 7.7

are compatible with the study findings, when the total eligible population is considered.

The authors conclude in their abstract that use of a dummy seems to reduce the risk of SIDS, but it could just be that reported dummy use in infants who died from SIDS was low because more mothers who gave their infants dummies declined to take part in the study.

The abstract also concludes that use of a dummy possibly reduces the influence of known risk factors in the sleep environment, but this is based on a spurious use of confidence intervals for the subgroups involved.² The more correct test of interaction reported in the results section showed that the difference in odds ratios between infants who did or did not use a dummy was not significant.

Christopher J Cates general practitioner
Manor View Practice, Bushey Health Centre, Bushey, Hertfordshire WD23 2NN
chrisates@nhs.net

Competing interests: None declared.

- 1 Li D-K, Willinger M, Petitti DB, Odouli R, Liu L, Hoffman H. Use of a dummy (pacifier) during sleep and risk of sudden infant death syndrome. *BMJ* 2006;332:18-21. (7 January.)
- 2 Altman DG, Bland JM. Statistics notes: interaction revisited: the difference between two estimates. *BMJ* 2003;326:219.

Funduscopy: to dilate or not?

Precipitation of angle closure may not be a disservice

EDITOR—Liew et al encourage clinicians to dilate pupils for thorough funduscopy.¹ The rare occurrence of precipitated angle closure with tropicamide 0.5% may not necessarily be a disservice to the patient, for two reasons.

Firstly, anyone whose angle may be provoked into closure by a mild mydriatic is at risk of spontaneous angle closure glaucoma. The fact that it has been precipitated in a healthcare setting, rather than occurring in the community, is of some benefit. The patient is likely to be within easy access of specialist care. We see patients who are at risk of angle closure but who, for example, have a propensity to travel (either during the day or for extended durations, as with the armed forces) or who cannot travel (because of being snow bound without ambulance access for several days).

Secondly, such angle closure would occur shortly after dilating drops are wearing off. Thus the symptoms are likely to be attributed to the preceding use of mydriatic drops and the correct diagnosis should be made quickly and treatment given promptly. Conversely, in cases of spontaneous angle closure patients often do not take notice of, or report, early symptoms. Clinicians too, may misinterpret symptoms or signs, leading to delay in diagnosis and a potentially poorer outcome. The fear of precipitating angle closure glaucoma should therefore not affect the decision to dilate to perform accurate funduscopy.

Kanchan J Bhan *specialist registrar, ophthalmology*
starfish1944@yahoo.com

Andrew Bastawrous *senior house officer, ophthalmology*
Keith G Davey *consultant ophthalmologist*
Department of Ophthalmology, Huddersfield
Royal Infirmary, Huddersfield HD3 3EA

Competing interests: None declared.

1 Liew G, Mitchell P, Wang JJ, Wong TY. Funduscopy: to dilate or not to dilate? *BMJ* 2006;332:3. (7 January.)

Other drugs can cause partial pupil dilatation

EDITOR—Liew et al highlight the very small risks associated with precipitating an attack of angle closure glaucoma when dilating a pupil for the purposes of funduscopy.¹

The initiating event in acute angle closure glaucoma is incomplete mydriasis (pupil dilatation) resulting in pupil block and the prevention of aqueous drainage through the trabecular meshwork. This partial mydriasis is normally caused by poor ambient lighting conditions, but drugs with adrenergic and anticholinergic properties have also been identified as causative agents. These drugs include ipratropium and tricyclic antidepressants.²⁻⁵ Serotonin selective reuptake inhibitors (SSRIs) such as paroxetine and fluoxetine have also been implicated through their anticholinergic properties.¹

The *British National Formulary* advises caution with using these drugs in patients

with a history of angle closure glaucoma.⁵ This is, however, misleading, as patients who have had an attack of angle closure glaucoma will hopefully have been treated with either iridotomies or trabeculectomies and will therefore no longer be at risk. We therefore recommend that all patients taking ipratropium, tricyclics, and SSRIs be warned of the risk of ophthalmic symptoms (painful red eye, blurring of vision, headache, nausea and vomiting), especially those at greater risk (elderly, female, hypermetropic refractive error). This should reduce ocular morbidity in patients who develop this complication.

Peter D Cackett *ophthalmology specialist registrar*
Princess Alexandra Eye Pavilion, Edinburgh
EH3 9HA
pete@pdackett.demon.co.uk

Competing interests: None declared.

- 1 Liew G, Mitchell P, Wang JJ, Wong TY. Funduscopy: to dilate or not to dilate? *BMJ* 2006;332:3. (7 January.)
- 2 Singh J, O'Brien C, Wright M. Nebulized bronchodilator therapy causes acute angle closure glaucoma in predisposed individuals. *Respir Med* 1993;87:559-61.
- 3 Ritch R, Krupin T, Henry C, Kurata F. Oral imipramine and acute angle closure glaucoma. *Arch Ophthalmol* 1994;112:67-8.
- 4 Browning AC, Reck AC, Chisholm IH, Nischal KK. Acute angle closure glaucoma presenting in a young patient after administration of paroxetine. *Eye* 2000;14:406-8.
- 5 Joint Formulary Committee. *British national formulary*. 50th ed. London: British Medical Association and Royal Pharmaceutical Society of Great Britain; September 2005.

Mobile medical teams rethink

EDITOR—The new emergency planning guidance from the Department of Health specifically addresses several of the issues raised by Walsh.¹⁻² The new guidance places the responsibility for medical support to a scene with the ambulance services and the strategic health authorities.

Medical incident commanders should come from a recognised pool of staff with an appropriate background and evidence of training, preparation, and experience for the role. They should form part of an on-call system that allows for a lead medical incident commander and numerous supporting clinicians to take the key medical leadership roles, such as in the casualty clearing station, described as needed by Walsh in the event of an incident. Mobile medical teams are replaced by MERITs (medical emergency response incident teams), and the strategic health authority has a responsibility to ensure that sufficient staffed and trained MERITs are available to provide for their area. Each hospital is no longer required to provide a medical team.

Pre-hospital medical training has been available through BASICS education and accredited by the Royal College of Surgeons of Edinburgh for over 10 years. This provides the specific skill set for responding to scene for the ambulance services and outlines clinical priorities in the event of a major incident. Specific training in major incident management has been provided for over 10 years by the internationally recognised MIMMS (major incident medical management and support) course administered by the Advanced Life Support Group.³

Darren Walter *consultant in emergency medicine*
South Manchester University Hospitals NHS Trust,
Wythenshawe Hospital, Manchester M23 9LT
darren.walter@smtr.nhs.uk

Competing interests: DW is an active BASICS doctor, a member of BASICS Education Committee, a member of the Board of the Faculty of Pre-hospital Care at the Royal College of Surgeons of Edinburgh, and a contributor to two subcommittees on emergency planning at the Department of Health.

- 1 Walsh EM. Is it time to rethink mobile medical teams? *BMJ* 2006;332:59. (7 January.)
- 2 Department of Health. *NHS emergency planning guidance 2005*. [www.dh.gov.uk/PublicationsAndStatistics/PublicationsPolicyAndGuidance/PublicationsPolicyAndGuidanceArticle/fs/en?CONTENT_ID=4121072&chk=yCJURQ](http://www.dh.gov.uk/PublicationsAndStatistics/Publications/PublicationsPolicyAndGuidance/PublicationsPolicyAndGuidanceArticle/fs/en?CONTENT_ID=4121072&chk=yCJURQ) (accessed 8 Jan 2006).
- 3 Advanced Life Support Group. Major incident medical management and support. www.walsg.org.uk/index.php?id=44 (accessed 8 Jan 2006).

Resistance to HIV drugs in UK may be lower in some areas

EDITOR—The UK Group on Transmitted HIV Drug Resistance reports an estimated prevalence of transmitted HIV drug resistance of 14.2% between 1996 and 2003 which is increasing over time.¹ Using the same definition of drug resistance as the UK group, a study of primary HIV infection at St Mary's Hospital, London, showed that the prevalence and annual incidence of transmitted drug resistance have remained low and stable, at 6% (9/140) between 2000 and 2005. Resistance to nucleoside reverse transcriptase inhibitors was detected in 1/9, to non-nucleoside reverse transcriptase inhibitors in 5/9, to protease inhibitors in 1/9, and to more than one class of drug in 2/9.

The disparity in estimated prevalence of transmitted drug resistance is interesting and may reflect the differences between the two cohorts. In our cohort, all individuals acquired HIV within six months of baseline genotyping and were predominantly white with homosexually acquired B clade viruses (125/140, 89%).

The UK group described a large cohort of 2357 individuals with chronic HIV infection presumed to be naive to antiretroviral therapy. Of these, 172 (7%) were infected within the previous 18 months, 22% of whom had drug resistance. The country of acquisition of HIV infection may contribute to the disparity since the use of antiretroviral therapy globally differs considerably, as in the use of nevirapine to prevent mother to child transmission.²⁻³ Most of our patients (98%) were infected in the UK. Although this was not assessed by the authors, the diversity of viral clades reported implies greater HIV acquisition outside the UK.⁴

The advantage of prospectively recruiting individuals with primary HIV infection and confirmed naive to antiretroviral treatment avoids the difficulties acknowledged by the group in relying on treatment histories reported by patients and doctors. Prevalence estimates of transmitted drug resistance in newly infected patients may not therefore be generalised to patients with established

infection who are reportedly treatment naive.

Julie M Fox *clinical lecturer*
julie.fox@imperial.ac.uk

Sarah Fidler *clinical senior lecturer*

Jonathan Weber *Jefferriss professor*
Section of Infectious Diseases, Faculty of Medicine,
Imperial College, London W2 1PG

Competing interests: None declared.

- 1 UK Group on Transmitted HIV Drug Resistance. Time trends in primary resistance to HIV drugs in the United Kingdom: multicentre observational study. *BMJ* 2005;331:1368. (10 December.)
- 2 McIntyre J. Prevention of mother-to-child transmission of HIV: treatment options. *Expert Rev Anti Infect Ther* 2005;3:971-80.
- 3 Jourdain G, Ngo-Giang-Huong N, Le Coeur S, Bowonwatanuwong C, Kantipong P, Leechanachai P, et al. Intrapartum exposure to nevirapine and subsequent maternal responses to nevirapine-based antiretroviral therapy. *N Engl J Med* 2004;351:229-40.
- 4 Tatt ID, Barlow KL, Clewley JP, Gill ON, Parry JV. Surveillance of HIV-1 subtypes among heterosexuals in England and Wales, 1997-2000. *J Acquir Immune Defic Syndr* 2004;36:1092-9.

Patients get four choices for NHS treatments

Choose and book has not left the station

EDITOR—UK general practitioners have no contractual obligation to offer patients a choice of provider.¹ My practice calculated that with some 50 elective referrals to hospital each week we would need to spend nearly £4500 this year for additional staff time to implement the service (patient population 6400). If we do receive the £0.95 per patient as payment for implementing the whole "Connecting 4 Health" package of which choose and book is but a part, my practice may achieve a net profit before tax of nearly £1500.

This is meagre compensation for the three partners, who must devote additional time and effort to assist staff to implement choose and book for our patients, although evidence of the clinical benefit for my patients is lacking. The additional time per appointment that may ensue will reduce our overall availability to see patients, thus jeopardising our facility to see everyone within 48 hours. Is this really the way to maintain morale in general practice and support an already overstretched service?

David M Lewis *general practitioner*
Tudor Surgery, Watford WD24 7PH
david.lewis@nhs.net

Competing interests: DML is a GMS general practitioner.

- 1 Coombes R. Patients get four choices for NHS treatments. *BMJ* 2006;332:8. (7 January.)

Choose and book will hinder development of good outpatient services

EDITOR—Is it not time that as a profession we state unequivocally that we are opposed to having choose and book imposed on us?¹ There are also cogent reasons for resistance by the hospital sector, principally that choose and book will hinder the development of good outpatient services.

Specialist "one stop clinics" have been one of the outstanding improvements of

hospital outpatient services in the past decade. They have been developed in many specialties (for example, prostrate, endoscopy, breast), bringing obvious benefits to patients' care, as well as improving clinic efficiency. They could be expanded to other disciplines, but this will be much more difficult with choose and book. Having instant access to diagnostic facilities such as fine needle aspiration cytology is difficult to organise and requires a substantial and potentially expensive time commitment from service sector departments and their staff, particularly consultants.

To have such facilities available every time a clinic in a specialty is held and in every location is not practical. Patients who are most likely to need and benefit from one stop assessment need to be concentrated in the clinics where it is available, so referrals inevitably need to be vetted and streamed. Choose and book is the antithesis of this process and will therefore reduce the quality and efficiency of clinic services. The scheme can therefore be opposed for reasons that are more to do with quality and standards of care than practicality and expense.

Jeremy Wood *consultant breast surgeon*
London N10 2BS
jeremy.wood@blueyonder.co.uk

Competing interests: None declared.

- 1 Coombes R. Patients get four choices for NHS treatments. *BMJ* 2006;332:8. (7 January.)

Wishful thinking about primary care in 2015

EDITOR—With regard to the five visions of primary care,¹⁻⁵ my heart hopes that the aspirations of Lakhani and Baker prevail, but my head suggests that the alternative scenarios outlined by Meldrum reflect the future.

I have had two contrasting careers—17 years as a general practitioner principal and then 13 years in medical management as a medical adviser to a health authority and a full time medical director to three primary care trusts. I am now returning to my clinician roots via the general practitioner returner scheme.

Interpersonal care, continuity, the advocacy role, and the longitudinal relationship between a practice team and a family have driven my career. However, in all the NHS management networks that I have worked in these values are never discussed at the management table. The Royal College of General Practitioners and the NHS seem to be in parallel universes. In NHS management the core values are the "3Cs": choice, contestability (whatever that is), and control.

Unless there is a massive revolt against what seems to be emerging policy, general practice will suffer the same fate as the local branches of banks, the local solicitor, or the local shop—and the public will only miss us when we are no longer there.

The core function of the consultation is still the most fascinating of intellectual challenges. But explaining some of the current

policies to patients—such as choose and book and changes in referral pathways and out of hours coverage—is met with bewilderment and then resignation, particularly as implementing choose and book here requires patients to ring a call centre. On what basis is this a patient led NHS?

Robert D Walker *general practitioner returner*
8 Moor Road, Workington, Cumbria CA14 1TS
robert@robertwalker98.wanadoo.co.uk

Competing interests: None declared.

- 1 Lakhani M, Baker M. Good general practitioners will continue to be essential. *BMJ* 2006;332:41-3. (7 January.)
- 2 Lapsley P. The magic roundabout. *BMJ* 2006;332:43. (7 January.)
- 3 Jeffries D. Return to a system based on social values. *BMJ* 2006;332:44-5. (7 January.)
- 4 Meldrum H. General practice under pressure. *BMJ* 2006;332:46-7. (7 January.)
- 5 Black C. Still striving for utopia. *BMJ* 2006;332:47-8. (7 January.)

Don't discount magnet therapy

EDITOR—Not all magnetic devices are equally non-efficacious.¹ Magnetic fields from bracelets and such may have little effect on human tissue, but it would be wrong to discount all effect on human tissue, even theoretically.

There is a strong research literature using transcranial magnetic stimulation (TMS) to induce neuronal firing in the brain and spinal cord.² Small pilot studies have been done using this technique in, for example, stroke,³ and Parkinson's disease,⁴ as well as many other neurological and psychiatric conditions. A neglect-like syndrome can be transiently induced in healthy subjects,⁵ shedding light on physiological and cognitive processes. While no large clinical trials have been done yet, it is crucial that not all magnetic therapy should be discounted.

James T H Teo *clinical research fellow*
Sobell Department of Motor Neuroscience,
Institute of Neurology, London WC1N 3BG
jte@ion.ucl.ac.uk

Competing interests: JTH is a clinical researcher working in neurophysiology.

- 1 Finegold L, Flamm BL. Magnet therapy. *BMJ* 2006;332:4. (7 January.)
- 2 Liepert J. Transcranial magnetic stimulation in neurorehabilitation. *Acta Neurochir* 2005;93(suppl):71-4.
- 3 Mansur CG, Fregni F, Boggio PS, Riberto M, Gallucci-Neto J, Santos CM, et al. A sham stimulation-controlled trial of rTMS of the unaffected hemisphere in stroke patients. *Neurology* 2005;64:1802-4.
- 4 Khedr EM, Farweez HM, Islam H. Therapeutic effect of repetitive transcranial magnetic stimulation on motor function in Parkinson's disease patients. *Eur J Neurol* 2003;10:567-72.
- 5 Fierro B, Brighina F, Oliveri M, Piazza A, La Bua V, Buffa D, Bisiach E. Contralateral neglect induced by right posterior parietal rTMS in healthy subjects. *Neuroreport* 2000;11:1519-21.

bmj.com

We select the letters for these pages from the rapid responses posted on bmj.com within five days of publication of the article to which they refer.

Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.