on glibenclamide, chlorpropamide, metformin, and insulin. The subjects were less severely diabetic than most given sulphonylureas, in that after diet treatment they had no symptoms but still had a raised fasting plasma glucose concentration (mean 8 mmol/l (144 mg/100 ml)). Therapy was aimed at reducing the fasting plasma glucose value to near normal (6 mmol/l (109 mg/100 ml)).

In a published report on 286 subjects the proportions whose doctors reported a single episode of any hypoglycaemic symptoms in one year were 31% for glibenclamide, 7% for chlorpropamide, 0% for metformin, and 8% for once daily insulin.¹ Most of these hypoglycaemic symptoms were mild and transient, occurring some two to four hours after glibenclamide was taken. A report from Sweden, however, described 57 cases of serious hypoglycaemia, including 10 deaths, in subjects taking glibenclamide.² In many cases there were contributing factors such as liver, kidney, or cardiac disease, and the median age of the patients was 70 years, but one man of 32 was left with permanent brain dysfunction. The median dose in the series was 10 mg daily but over one third of patients were receiving only 2.5 mg or 5 mg daily.

Campbell calculated that glibenclamide induced hypoglycaemia and metformin induced lactic acidosis carry a similar risk of mortality in Sweden, but he noted that the 10 deaths from glibenclamide in 300 645 patient years of use in Sweden contrasted with the absence of reports of a case of metformin induced lactic acidosis in the UK in 330 000 patient years of use from 1972 to 1982.³ It is usually said that chlorpropamide should be used with caution in elderly patients in view of its long plasma half life and the possibility of high plasma concentrations in patients with impaired renal function. These recent reports suggest that there is also a risk of hypoglycaemia with glibenclamide. Such episodes are generally transient and mild but may be more severe and even life threatening, especially in elderly subjects or in those with impairment of hepatic, renal, or cardiac function. Glibenclamide should be used carefully in such subjects, particularly in those with milder degrees of hyperglycaemia, and, as with other sulphonylureas, it is important to monitor the response to therapy by blood glucose measurements.

JOHN S YUDKIN

Academic Unit of Diabetes and Endocrinology,
Whittington Hospital,
London N19 5NF

- 1 Multicentre study. UK prospective study of therapies of maturity-onset diabetes. 1. Effect of diet, sulphonylurea, insulin or biguanide therapy on fasting plasma glucose and body weight over one year. *Diabetologia* 1983;24:404-11.
- 2 Asplund K, Wiholm BE, Lithner F. Glibenclamide-associated hypoglycaemia: a report on 57 cases. *Diabetologia* 1983;24:412-7.
- 3 Campbell IW. Metformin and glibenclamide: comparative risks. *Br Med J* 1984;289:289.

Management consultancy studies on cancer treatment services in the Thames regions

SIR,—The management consultants Coopers and Lybrand Associates have recently reviewed cancer treatment services in the North East Thames region¹ and are currently undertaking similar reviews for the North West and South West regions—each for fees reportedly of the order of £60 000 to £100 000. However, for the North East Thames region several fallacies have been included in the Coopers and Lybrand model for cancer prevalence. These could lead to significant errors in the estimates of the needs of future patients with cancer and therefore nullify their recommendations for reorganising cancer treatment services in

Fallacies in the Coopers and Lybrand model of cancer prevalence in the North East Thames region

Fallacy	Suggested modification
(1) Cancer site groups contain subsites with very different survival rates—eg, the head and neck group includes tongue, pharynx, larynx, and brain; the upper digestive tract group includes oesophagus, stomach, and pancreas. It is not known if the cancer site "mix" will change substantially with time.	(1) The site groups should be more precise. The following 10 should be included individually since they account for some two thirds of all cancer in England and Wales: oesophagus, stomach, colon, rectum, lung, skin, breast, prostate, bladder, leukaemia.
(2) The definition of "cure" from cancer is incorrect. The survival distributions which were inspected were for all causes of death and a levelling off in survival for a period of time is an artefact, possibly due to small sample sizes. It is not a demonstration of "cure."	(2 & 3) Data should be assembled by site and age, not only the overall survival but also for recurrence free periods. As well as regional and hospital cancer registry records, those of hospices and centres for continuing care can provide data on the cancer services requirements for the terminally ill.
(3) It is implied that those who eventually die from cancer provide a problem for cancer services from initial treatment to death. This is untrue since many experience long periods with a good quality of life. During this time only regular follow up services are required.	
(4) Statements on new courses of radiotherapy treatment for "new" and for "recurrent" cancers are inconsistent: Volume I, table 7—13 200 courses needed for new and recurrent cancer ¹ ; Volume II, para 1.26—4650 courses needed for new cancer. ¹ This implies that only 35% of courses are for new (previously untreated) cancers. Data from other sources indicates that this figure is 60% (Mould RF, Hanham IWF, statistics prepared for Copper and Lybrand review, 1986); 35% must be an underestimate.	(4) A review of cancer treatment statistics should include consistent data on the requirement for radiotherapy treatment of new and recurrent cancer. It should also estimate, by site group, those patients who will remain untreated except for receiving hospice or continuing care—eg, in North West Thames in 1980 20% of lung cancer registrations were untreated. ²

North East Thames. Their prevalence model should be corrected: see table.

R F MOULD

Westminster Hospital Cancer Registry,
Department of Medical Physics,
London SW1P 2AP

- 1 Coopers and Lybrand Associates, North East Thames Regional Health Authority. *Review of cancer treatment services*. Vol I. A review of cancer services, technical annexes. Vol II. London: NETHERA, 1985.
- 2 Mould RF, Williams RJ. Survival of histologically proven carcinoma of the lung registered in the North West Thames Region, 1975-9. *British Journal of Cancer* 1982;46:999-1003.

So what will fibrinolytic therapy offer your patients with myocardial infarction?

SIR,—It is hard to agree with Professor J R A Mitchell (12 April, p 973) that we must look forward to a decade of controversy over the intracoronary versus intravenous method of applying one of the lytic agents now available. He himself has pointed out that the former as a routine is really quite impracticable. Once the intracoronary method has shown that prompt fibrinolytic treatment does restore patency to a substantial proportion of coronary occlusions it is surely no longer ethical to subject ill patients to academic experiments. One wonders too at the ethics of further experiments on fibrinolysis versus routine coronary care if they may mean postponing the optimal time for a coronary bypass operation on some of the participants so that they may

complete their follow up protocol. There does, however, seem to be a need for investigating the most suitable of the lytic agents to use, but the other really important point is how can coronary thrombolysis be brought most quickly and efficiently to the patient. There is one aspect of this which seems to have been overlooked.

When in 1977 Smith, Hacking, and I treated eight patients with intracoronary streptokinase we always gave intravenous plasminogen at the earliest opportunity while coronary arteriography was being prepared.¹ It would have been much better if it had been given before the patient entered hospital, during the phase of propagating thrombosis which succeeds the initial occlusion. The rationale behind this procedure is well illustrated by in vitro experiment. If one makes a fibrin plate by clotting a solution of fibrinogen in a Petri dish with thrombin and then applies to the surface a drop of some fibrinolytic activator, it commonly requires 18 hours of incubation at 37°C before lysis is evident. But if one has first enriched the fibrinogen with extra plasminogen then the period of incubation can be reduced to two hours.²

I suggest that experiments be conducted to see if this early enrichment of fibrin(ogen) is really worth while, especially now that non-invasive methods are being developed to detect the moment when blood flow has been restored (Richardson SG *et al.* British Cardiac Society Meeting, 1986). At present plasminogen is expensive and must be kept in a refrigerator but then this used to be true of penicillin. It does, however, have a reasonably long half life in the body so that an intravenous bolus rather than a drip should suffice. If pretreatment with it does prove of value then the family practitioner and perhaps specially trained ambulance crews may have a new part to play in the early management of this still dangerous disease.

H A DEWAR

Wylam,
Northumberland NE41 8AS

- 1 Dewar HA, Smith RH, Hacking PM. Acute lysis of coronary thrombi. *Br Heart J* 1978;40:436.
- 2 Marsh NA, Gaffney PJ. The rapid fibrin plate—a method for plasminogen activator assay. *Thromb Haemost* 1977;38:545-51.

Toxicity of vitamins

SIR,—The article by Drs C D H Evans and J Hubert Lacey (22 February, p 509) includes an appendix which perpetuates the myth that withdrawal of pyridoxine treatment can result in symptoms of dependency.

Their appendix II lists the toxic effects of vitamin B₆ (pyridoxine) as dependency, peripheral sensory neuropathy and ataxia, and decrease in therapeutic effect of levodopa.¹ Of these the latter two effects are well documented. Peripheral neuropathy and associated ataxia have been reported in patients taking high doses of pyridoxine (200 mg-6 g/day) over periods of two months to three years.^{2,3} This neuropathy gradually improves after cessation of pyridoxine. The peripheral decarboxylation of levodopa is enhanced by pyridoxine, which results in reduced effectiveness of levodopa treatment of Parkinson's disease. Pyridoxine and multivitamin preparations containing vitamin B₆ should therefore be avoided in patients receiving levodopa. This interaction does not occur when levodopa is given with a decarboxylase inhibitor such as benserazide (Madopar).⁴

However, dependency after pyridoxine treatment is not well substantiated. On checking the two references quoted by Davidson as the sources for the pyridoxine entry in his paper¹ there is no mention in either of them of dependency due to

pyridoxine.^{5,6} There is one published paper which showed that three out of eight subjects had an abnormal electroencephalogram eight days after supplementation of pyridoxine 200 mg daily was stopped.⁷ This study, performed in 1964, has been interpreted as showing that symptoms of dependency occur after withdrawal of pyridoxine treatment. These findings have not been reproduced, nor have practitioners with experience of treating large series of patients with pyridoxine in doses of 20-150 mg encountered symptoms of dependency after cessation of treatment (M Brush, personal communication).

It is unfortunate that tables are modified for publication without the primary sources of reference being verified. This can result in unsubstantiated effects being incorporated into published reports as fact, solely by virtue of repetition.

A J SHIPMAN
S J PARKER (Miss)

Roche Products Ltd,
Welwyn Garden City,
Hertfordshire AL7 3AY

- Davidson RA. Complications of megavitamin therapy *South Med J* 1984;77:200.
- Schaumburg H, Kaplan J, Windebank A, *et al.* Sensory neuropathy with low dose pyridoxine abuse. A new megavitamin syndrome. *N Engl J Med* 1983;309:445.
- Parry G, Bredesen DE. Sensory neuropathy with low-dose pyridoxine. *Neurology* 1985;35:1466.
- Marcus R, Coulston AM. Water-soluble vitamins. The vitamin B complex and ascorbic acid. In: Gilman AG, Goodman LS, Rall TW, Murad F, eds. *Goodman and Gilman's the pharmacological basis of therapeutics*. 7th ed. New York: Macmillan, 1985.
- Schaumburg H, Kaplan J, Rasmus S, *et al.* Pyridoxine megavitaminosis produces sensory neuropathy (neuropathy) in humans. *Ann Neurol* 1982;12:107.
- Taft G, Fieldhouse P. Vitamin C and the common cold. *Public Health* 1978;92:19.
- Canham JE, Nunes WT, Eberlin ET, *et al.* Electroencephalographic and central nervous system manifestations of B₆ deficiency and induced B₆ dependency in normal human adults. *Proceedings of the sixth international congress of nutrition*. Edinburgh: Livingstone, 1964:537.

Ampoules, infusions and filters

SIR,—The statement in the leading article by Dr D G Walker and Professor C F George (15 March, p 714) that "in line filtration will reduce the risks of contamination with micro-organisms" needs some qualification, as it could be incorrectly interpreted to mean that these filters reduce the rate of all forms of cannula associated infection. The major source of bacteria that colonise these lines appears to be the patients' skin flora. These bacteria usually gain access to indwelling cannulas by the cannula-skin junction.¹ Less often infection results from intrinsically contaminated intravenous fluid, contamination in the giving set, or haematogenous spread. Only infection resulting from intrinsically contaminated fluid or contamination in the giving set occurring above the filter will be prevented by the use of these filters. Conversely, the addition of another device with consequent increased manipulation of the giving set may actually increase the risk of infection.¹

Care should also be exercised in selecting an appropriate filter pore size. Dr Waller and Professor George suggest pore sizes of between 0.2 µm and 0.5 µm. A porosity of 0.5 µm, however, will not effectively retain all small Gram negative bacilli such as pseudomonas.² A 0.22 µm filter would be needed for an adequate barrier, but, unfortunately, viscous solutions cannot pass through such a filter without a pump. Thus the use of infusion filters will not affect most line associated infections, and suitable bacterial filters may lead to practical problems. Although the use of infusion filters may reduce the incidence of infusion associated phlebitis due to particulate matter, they should be used with

circumspection and not as a substitute for careful aseptic procedures in the insertion and maintenance of intravenous lines.

S J PEDLER
T S J ELLIOTT

Department of Microbiology,
Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP

- Friedland G. Infusion-related phlebitis—is the in-line filter the solution? *N Engl J Med* 1985;312:113-5.
- Brismar B, Malmberg A-S, Nystrom B, Strandberg A. Bacterial contamination of intravenous cannula injection ports and stopcocks. *Hum Nutr Clin Nutr* 1984;iii:23-6.

Tumour markers in germ cell tumours

SIR,—The title of the leading article by Dr Gordon J S Rustin (15 March, p 713) led readers to expect an update on these markers, but the title misfits the text, and an interesting discussion of a supra-regional assay scheme followed. There was a brief aside to include markers other than the established α fetoprotein and β subunit of human gonadotrophin; raised activities of serum alkaline phosphatase occur in half the patients with seminoma; some patients also have raised lactate dehydrogenase activity. No mention is made, however, of a recent report on B5, a new marker for testicular tumour.¹ This report includes 28 patients who were tested at diagnosis, 27 of whom yielded positive results on B5 testing (including all of the 10 patients with seminoma). When compared with α fetoprotein and β subunit of human gonadotrophin, B5 proved to be a sensitive indicator for residual disease or early recurrence. The B5 marker is on erythrocytes and is detected by a simple haemagglutination test; it is non-specific for tumour type but highly sensitive for testicular tumour. Monitoring showed that patients who were free of disease after treatment became negative on B5 testing, the decay time relating to the erythrocyte life span.

We have now extended the programme to monitor more than 90 patients at Cookridge Hospital, Leeds (in collaboration with the Yorkshire Urological Cancer Group, including Mr R Ewing, Mr J Hetherington, and Dr H Close), and the overall results show a correlation between increased B5 state and active disease in more than 80% of cases. Thus we feel that the simple B5 test provides a valuable addition to α fetoprotein and β subunit of human gonadotrophin for monitoring disease state, especially when α fetoprotein and β subunit of human gonadotrophin activities remain normal.

Department of Surgery,
Addenbrooke's Hospital,
Cambridge

SU METCALFE

W G JONES

Regional Radiotherapy Centre,
Cookridge Hospital,
Leeds LS16 6QB

- Metcalfe S, Sikora K. A new marker for testicular cancer. *Br J Cancer* 1985;52:127-9.

Reaction to food additives

SIR,—“I knew that food additives provoked strong feelings among lay people,” says Caroline Richmond (15 March, p 754). Indeed so. This is partly because the views of expert government advisory committees over the years are now being brought to light.

Take azo dyes, now reliably identified as toxic to an unknown proportion of children.^{1,2} In 1924 a Ministry of Health departmental committee reported: “Colour is frequently used to cover up objectionable or inferior materials, or to give a factitious appearance, so that the objects so

coloured masquerade for something which they are not.”³

The Ministry of Food (as was) appointed a Food Standards Committee on Colouring Matters, which reported in 1954. Its view? “We cannot accept the contention that, because ‘coal-tar’ colours have been used in foods for many years without giving rise to complaint of illness, they are, therefore, harmless substances. Such negative evidence in our view merely illustrates that in the amounts customarily used in food the colours are not acutely toxic but give no certain indication of any possible chronic effects. Any chronic effects would be insidious.”⁴

By 1979 toxicity of azo dyes was better understood, as indicated by the current report, of the Food Additives and Contaminants Committee, on colouring matter. This classified 17 colours (including 12 azo dyes) as “provisionally acceptable” pending studies (usually for carcinogenicity, mutagenicity, and teratogenicity) to be carried out within three to five years, adding: “We would strongly emphasise that if the work required by the Committee on Toxicity is not carried out on any colouring matter by the time suggested, we will not be prepared to recommend its continued use in food.”⁵

Since this there has been silence. This is the context in which the Food Additives Campaign Team, representing more than 20 relevant organisations and supported by legislators of all parties, is among other things calling for the withdrawal of azo dyes from foods claimed or described to be suitable for children under 5 years of age.⁶ This is not, I think, to use Ms Richmond's phrase, a “food additives scare.” She thinks she has made “the first attempt in the popular press to introduce a little science into the discussion” on food additives, citing her “hypothesis” that hyperactivity may be caused by pushchairs and television. Is there a reference forthcoming to support this novel idea?

GEOFFREY CANNON

Food Additives Campaign Team,
London N5 1XL

- The Royal College of Physicians and the British Nutrition Foundation. Food intolerance and food aversion. *J R Coll Physicians Lond* 1984;18:3-42.
- Egger J, Graham P, Carter C, Gumley D, Sothill J. Controlled trial of oligoantigenic treatment in the hyperkinetic syndrome. *Lancet* 1985;ii:540-5.
- Ministry of Health. *Report of the departmental committee on the use of preservatives and colouring matters in food*. London: HMSO, 1924.
- Ministry of Food. *Report on colouring matters*. Food Standards Committee. London: HMSO, 1954.
- Ministry of Agriculture, Fisheries, and Food. *Interim report on the Review of the Colouring Matter in Food Regulations 1973*. Food Additives and Contaminants Committee. London: HMSO, 1979.
- Food Additives Campaign Team. *Food additives—a new national policy*. London: FACT.

Trial of strategy for reducing the use of laboratory tests

SIR,—The study by Dr F G R Fowkes and others (29 March, p 883) showing a reduction in laboratory tests after the introduction of guidelines and a weekly review of medical records in a hospital medical unit is most encouraging. The study parallels work on guidelines in other areas, including antibiotics.¹

The methods used by Dr Fowkes and others included awareness by staff who were given the guidelines that their requests for laboratory tests were being monitored during the intervention period. Staff in the control group were not so advised. Might this awareness have led to a “Hawthorne effect”—that is, knowledge of monitoring could have been a major influence leading to the reduction in laboratory test use? If this were the case it would be difficult to determine whether the