CORRESPONDENCE

Informed consent R R Hall, FRCS, and P H Smith, FRCS; T B Brewin, FRCP; P J Wyld, MRCPATH; C W Burke, FRCP 389	Dilated cardiomyopathy, myocarditis, and the biotope P J Lowry, MRCP, and others; E G J Olsen, FRCPATH 392	A better way of selecting people for jobs? M P Ward Platt, MRCP 394
Neurological complications of coronary artery bypass graft surgery A V Pollock, FRCS 390	Transcutaneous oxygen tension during exercise in patients with claudication D H Roberts, MRCP, and others; E L Gilliland, FRCS, and others 392	Whatever happened to the Black report? R R Gordon 394
Investigation of plasma lipids—which tests and when? H J Dodd, MRCP, and I Sarkany, FRCP 391	Respite care on a children's ward Josephine E Hammond, FRCPED; Leila B Cooke, MRCPsych, and Ruth M Walters, MRCPsych 393	Do keyboards cause loss of masculinity? Caroline Richmond, MSC 394
Opiate withdrawal: inpatient versus outpatient programmes C Brewer, MRCPsych 391	Diffuse peritonitis and chronic ascites due to infection with Chlamydia trachomatis G M Cawdell, MRCOG 393	The Wendy Savage case B Raymond 394
Bleeding from peptic ulcers and use of non-steroidal anti-inflammatory drugs J O'Brien, MRCP, and W R Burnham, MRCP ... 391	Hiding behind a box number R R Livings, MRCP 393	Points Drowning and near drowning (G A R Morgan and R J Winter); 1,25 Dihydroxy- vitamin D₃ in psoriasis (R M Francis and D M Beaumont); Paying for old age (J R Caldwell); Opiate withdrawal: inpatient versus out- patient programmes (D H Marjot); Measles, mumps, and rubella (C M Robertson and others); Reye's syndrome and aspirin (J Anderson); Cigarette smoking and risk of premature stroke in men and women (J S McCormick) 395
Drugs in developing countries Philippa Saunders 391	Selection of medical students by medical schools S Lessels, MB 394	

Because we receive many more letters than we have room to publish we may shorten those that we do publish to allow readers as wide a selection as possible. In particular, when we receive several letters on the same topic we reserve the right to abridge individual letters. Our usual policy is to reserve our correspondence columns for letters commenting on issues discussed recently (within six weeks) in the BMJ.

Letters critical of a paper may be sent to the authors of the paper so that their reply may appear in the same issue. We may also forward letters that we decide not to publish to the authors of the paper on which they comment.

Letters should not exceed 400 words and should be typed double spaced and signed by all authors, who should include their main degree.

Informed consent

SIR,—Jonathan Glover's leading article (19 July, p 157) has drawn attention to the Medical Research Council's trial of immediate and deferred orchidectomy in prostate cancer, which has been criticised by certain national newspapers and the medical press in Britain and North America.¹

These uninformed reports, which have caused considerable distress to patients with prostate cancer, some of whom have not been in the trial, have been initiated by articles in the March and April issues of the *Bulletin* of the Institute of Medical Ethics in which the editor, Dr R Nicholson, has stated, among many other things, that: "Old men... are being castrated without their informed consent, for the benefit of a trial... that is of little, and possibly no, scientific value." "This trial is too late... both the scientific understanding of prostatic carcinoma and its treatment have advanced to the point at which it is no longer certain that the patients in this trial are being offered the best treatment." "The MRC protocol does not require subjects to be told of the alternatives to castration... and it appears that none of them are given this information." "The MRC... is content that research carried out under its auspices should contravene all current national and international guidelines for the conduct of research on humans."

We wish to inform you that these claims misrepresent the facts concerning this trial.

Many, perhaps most, surgeons still treat all patients whose carcinoma has spread beyond the prostatic capsule. Orchidectomy is usually the treatment of choice in the United Kingdom. However, data described elsewhere (Handley RC *et al*, Second International Symposium on Prostatic Cancer, Paris 1986)² show that a significant number of British men with prostate cancer do not need treatment for their malignancy and for these patients a policy of deferred treatment is the most

appropriate. Most patients do need treatment but it is not known if this is best given immediately or when disease progression occurs because the necessary comparative trial has never been conducted. We agree that this trial should have been done 20 years ago but it was not and, as a result, the question remains unanswered. It is thus even more important that the trial is conducted now.

The reason Dr Nicholson claims the trial to be too late is because of his belief that other treatments have superseded orchidectomy, in particular the introduction of luteinising hormone releasing hormone analogues and total androgen ablation as advocated by Labrie.³ In this respect Dr Nicholson has allowed himself to be persuaded by the results of an uncontrolled study that are not supported by the randomised study of the European Organisation for Research and Treatment of Cancer or by the early results of the Canadian randomised trial that is testing the "Labrie hypothesis" (Beland F *et al*, British Association of Urological Surgeons and Canadian Urological Association meeting, London 1986). His suggestion that patients treated in the MRC trial are being denied the best treatment is unsubstantiated. Since it may lead some patients to believe that they are being treated inadequately his statement is probably unethical.

"Informed consent" is Dr Nicholson's main concern and was discussed in part in your leading article. It is true that the trial protocol allowed that some patients could be treated without detailed informed consent if in the opinion of the surgeon this was thought to be in the patient's best interest. This is in agreement with previous MRC guidelines that recognise "for example, to awaken patients with a possibly fatal illness to the existence of doubts about effective treatment may not always be in their best interest."

That this should be accepted by the MRC and

the many ethical committees that have approved the trial does not contravene all current national and international guidelines for the conduct of research on humans. The Declaration of Helsinki states, "If the doctor considers it essential not to obtain informed consent, the specific reasons should be stated for transmission to the independent committee." The guidelines on the practice of ethics committees in medical research published by the Royal College of Physicians of London in 1984 state: "In general patients should be told that a trial is in progress and that they are being given the best available treatment or one which may be better or worse. Only under exceptional circumstances when it would cause more distress to reveal the nature of the experiment is there an argument for not telling the patients; however, this should be a deliberate decision taken as part of the ethical review."

Contrary to Dr Nicholson's belief, British urologists do talk to their patients about their prostate cancer and its treatment. The average British patient is 10 years older than his American counterpart and the awareness of cancer and discussion about treatment is different from that which is common in the USA. We accept that every patient has an absolute right to be informed but we also have to work in a world in which some patients do not wish to exercise this right and their families do not wish them to be told that they have prostate cancer. Even more do not wish to choose their own treatment from the several available, and the phrase "You decide what's best" is still to be heard from patients in prostate clinics, despite a diligent attempt by the urologist to explain treatment options and the desirability of inclusion in a randomised trial. It may be argued that such patients should be excluded from clinical trials but is it better that their treatment should be chosen by a surgeon's personal preference, with no more

explanation, than by randomisation? The continuing debate on "informed consent" is an indication of the different opinions that still exist among medical and lay workers in the caring professions. It is for this reason that district ethical committees have been made the final arbiters of ethical practice in the health service and have been established in preference to a single, national ethical committee.

Dr Nicholson claims that some ethics committees have "rubber stamped" the MRC trial protocol without proper discussion. If so, this is the fault of the ethical committees concerned and not of the MRC, whose protocols always require local ethical corroboration.

We met Dr Nicholson before his initial publication on this matter. We would be pleased to meet him again together with his urological advisers and senior members of the Institute of Medical Ethics should they still have reservations about the scientific or ethical basis of this MRC trial.

R R HALL

Chairman,

MRC working party on urological cancer

Freeman Hospital,
Newcastle upon Tyne

P H SMITH

Chairman,

MRC subgroup on prostatic cancer

St James Hospital,
Leeds

- 1 Anonymous. Castration without consent? *Medical World News for Obstetricians, Gynecologists, Urologists*. 1986;29 May.
- 2 Newling DW, Hall RR, Richards B, Robinson MRG, Hetherington JW. The natural history of prostate cancer—the argument for a no treatment policy. In: Pavone-Macaluso, Smith, Bagshaw, eds. *Testicular cancer and other tumours of the genitourinary tract*. New York: Plenum Press, 1985:443-8.
- 3 Labrie F, Belanger A, Dupont A, Edmond J, Lacoursiere Y, Monfette G. Combined treatment with LHRH agonist and pure antiandrogen in advanced carcinoma of prostate. *Lancet* 1984;ii:1090.

SIR,—In referring to the problem of informed consent when treatment for cancer is randomised Jonathan Glover says, "There are two possible arguments for specific exceptions to the requirement." With respect, not only are there others that he does not mention, there are better ones.¹⁻³

In several passages—for example, when he talks of "unacceptable risks to patients" or when he asks, "Can it be right to impose such a trial on uninformed and unconsenting patients?"—Dr Glover seems to be perpetuating a common misconception: the idea that when a patient's treatment is decided by randomisation he automatically risks being treated in a way that would not normally be considered appropriate. This is not so.

There are many ways of forming an opinion as to the wisdom or correctness of a patient's treatment. Checking to see if the treatment was randomised is not one of them. A doctor randomising treatment is just as responsible for each alternative as one giving non-randomised treatment. Those who show excessive distrust and suspicion when treatment is randomised also show excessive confidence and approval when it is not. In both cases everyone hopes that the patient will get the best treatment, but "best" is full of complexities. Both early and late advantages and disadvantages are relevant. If it were not for the problem of selection bias valid comparisons could be made without the need for randomisation, but selection bias leads easily to false conclusions.

A really enlightened public would be calling for far more randomised trials to be done to protect them from methods that have never been properly compared with other—perhaps less drastic—alternatives.^{4,5}

T B BREWIN

Institute of Radiotherapeutics and Oncology,
Glasgow G11 6NT

- 1 Hill AB. Medical ethics and controlled trials. *Br Med J* 1963;ii:1043-9.

- 2 Atkins H. Conduct of a controlled clinical trial. *Br Med J* 1966;iii:377-9.
- 3 Brewin TB. Consent to randomised treatment. *Lancet* 1982;ii:919-21.
- 4 Silverman WA. *Human experimentation*. Oxford: Oxford University Press, 1985.
- 5 Chalmers I. Scientific enquiry and authoritarianism in perinatal care and education. *Birth* 1983;10:151-64.

SIR,—I agree with Jonathan Glover that fuller debate of exceptions to the general rule of informed consent would be valuable. Otherwise I believe some clinicians participating in MRC trials may find themselves faced with more problems—not least from local ethical committees, who could make participation in the trials increasingly difficult.

There are, for example, real problems with gaining truly informed consent from adults with some malignant diseases. I do not think one can simply ask a patient newly diagnosed with acute leukaemia to give consent for a randomised trial after an initial discussion. In many cases informed consent is part of the developing relationship of mutual trust that forms between patient and doctor. This occurs during days and weeks after initial diagnosis, and, in my own practice, informed consent is arrived at over such a period. In the same way information about the patient's disease is gradually imparted. We must give patients a minimum of information at diagnosis, and I suspect that this may, with some difficulty, need to be more clearly defined. There are wide variations in individual clinicians' practices.

We still depend on the common sense of our peers who organise clinical trials. It is important that all those who participate keep closely concerned with the trial and have an opportunity to discuss all aspects with the organisers. In haematology we are fortunate that things are arranged in this way.

We must not lose sight of the fact that we must do what is best for the patient. The amount of information he can absorb at the time of diagnosis is variably limited. We certainly have a duty to inform, but just how much and when and to whom is the clinician's judgment. Despite my earlier caveat this must still broadly remain the doctor's decision.

PETER J WYLD

Department of Haematology,
Royal Hospital,
Chesterfield S44 5BL

SIR,—Jonathan Glover is right to express concern about informed consent. When I was chairman of the central Oxford ethics committee for a number of years 90% of the queries to applicants concerned informed consent. It is clearly vital, but his discussion of exceptions to the rule is superficial.

Research for its own sake is justifiable in the test tube but not in humans. To be ethically justifiable in humans a research programme must be capable of providing an answer. For this reason ethics committees' chairmen find themselves turning down rather non-invasive but trivial research—for example, questionnaires may be as invasive as coronary catheters in certain circumstances while the subject matter may be trivial and hence the research unethical—yet they may give ethical approval to quite major and invasive research (especially if in the presence of informed consent the question being answered is of fundamental human importance).

It follows from this that any research is justifiable only if it is capable of producing an answer, and a research protocol submitted where an answer to the question being posed (which may or may not be worth while) is incompatible with informed consent is unethical. One such was a trial of carotid surgery versus conservative management in carotid

stenosis. You cannot randomise people to surgery or not surgery without doing sham operations, acceptable in the rat but not in a hospital. In that particular trial therefore the randomisation, it was eventually agreed, had to be without informed consent, though obviously consent was given by those randomised to surgery. This is, of course, ethical because in certain centres the patient might well have been advised to have surgery anyway, and the purpose of the trial was to find out whether that was in fact ethically or scientifically acceptable.

It is noteworthy how strongly many of the self styled "institutes" of medicine, medicine and the law, law and human ethics, and so on, lack any constructive medical input or on the ground experience. While the points they make should be scrutinised and listened to see if they are appropriate or not, there is no guarantee that they are any more or any less appropriate than those of an enthusiastic research investigator. Enthusiasm takes different forms.

C W BURKE

Department of Endocrinology,
Radcliffe Infirmary,
Oxford OX2 6HE

Neurological complications of coronary artery bypass graft surgery

SIR,—In the 19 July issue of the journal are two important papers—an editorial by Jonathan Glover on informed consent (p 157) and a follow up study by Dr Pamela J Shaw and her colleagues of 312 patients who had coronary artery surgery (p 165). Four of the 312 patients were severely incapacitated by the after effects of major peri-operative strokes. When these four are added to the one who died postoperatively with neurological complications it appears that the risk of fatal or permanent neurological damage after coronary artery bypass graft surgery in their series was 1.6%. The question now arises: should patients about to undergo such operations be specifically warned about the possibility of death or permanent disability from brain damage during operation? Lord Scarman, giving judgment in the Sidaway case, accepted the four propositions expounded in the United States Court of Appeals, District of Columbia Circuit, *Canterbury v Spence* (1972) (464 F 2d 772). These were:

(1) The root premise is the concept that every human being of adult years and of sound mind has a right to determine what shall be done with his own body. (2) The consent is the informed exercise of a choice, and that entails an opportunity to evaluate knowledgeably the options available and the risks attendant on each. (3) The doctors must, therefore, disclose all "material risks"; what risks are "material" is determined by the "prudent patient" test, which was formulated by the court (464 F 2d 772 at 787): "(a) risk is . . . material when a reasonable person, in what the physician knows or should know to be the patient's position, would be likely to attach significance to the risk or cluster of risks in deciding whether or not to forego the proposed therapy." (4) The doctor, however, has what the court called a "therapeutic privilege." This exception enables a doctor to withhold from his patient information as to risk if it can be shown that a reasonable medical assessment of the patient would have indicated to the doctor that disclosure would have posed a serious threat of psychological detriment to the patient.

The relationship between surgeons and patients is changing, and the traditional consent form is no protection against an accusation of negligence by trespass if we fail to instruct our patients in detail not only about the benefits but also about the costs of any operation.

A V POLLOCK

Scarborough Hospital,
North Yorkshire YO12 6QL