Published in final edited form as:

Nature. 2005 November 17; 438(7066): 318–324.

Structure of the *E. coli* protein-conducting channel bound to a translating ribosome

Kakoli Mitra^{1,*}, Christiane Schaffitzel^{2,*}, Tanvir Shaikh³, Florence Tama⁴, Simon Jenni², Charles L. Brooks III⁴, Nenad Ban², and Joachim Frank^{1,3,5}

¹ Howard Hughes Medical Institute, Health Research, Inc. at the Wadsworth Center, Empire State Plaza, Albany, NY 12201-0509, USA

² Institute for Molecular Biology and Biophysics, Eidgenössische Technische Hochschule Hönggerberg (ETH Zürich), CH-8093 Zürich, Switzerland.

³ Wadsworth Center, Empire State Plaza, Albany, NY 12201-0509, USA

⁴ Department of Molecular Biology (TPC6), The Scripps Research Institute, 10550 Torrey Pines Road, La Jolla, CA 92037, USA

⁵ Department of Biomedical Sciences, State University of New York at Albany, Empire State Plaza, Albany, NY 12201-0509, USA

Abstract

Secreted and membrane proteins are translocated across/into cell membranes via a protein-conducting channel (PCC). We present a cryo-EM reconstruction of the *E. coli* PCC, SecYEG, complexed with the ribosome and a signal anchor containing nascent chain, showing mRNA, three tRNAs, the nascent chain, and detailed features of both a translocating PCC and a second, non-translocating PCC bound to mRNA hairpins. The translocating PCC forms connections with ribosomal RNA hairpins on two sides and ribosomal proteins at the back, leaving a frontal opening. Normal mode-based flexible fitting of the archaeal SecYEG structure into the PCC EM densities favors a front-to-front arrangement of two SecYEG complexes in the PCC, and supports channel formation by the opening of two linked SecY halves during polypeptide translocation. Based on our observation in the translocating PCC of two segregated pores with different degrees of access to bulk lipid, we propose a model for co-translational protein translocation.

INTRODUCTION

The proper functioning of every living cell requires that secreted, soluble proteins be translocated across, and membrane proteins be inserted into, a cell membrane. Protein translocation at the membrane occurs through a proteinaceous channel ¹, the translocon ², at the core of which lies the protein-conducting channel (PCC). The PCC is a heterotrimeric integral membrane protein complex ^{3, 4} composed of the α -subunit (SecY in eubacteria/archaea, Sec61 α in mammals), the β -subunit (SecE in eubacteria/archaea, Sec61 β in mammals) and the γ -subunit (SecE in eubacteria/archaea, Sec61 γ in mammals). Most membrane proteins and some soluble proteins are translocated co-translationally, a process in which the ribosome binds to the PCC, and nascent chain translocation is concomitant with polypeptide elongation on the ribosome. The PCC is a dynamic complex, which must be able to open and close an aqueous channel ^{5, 6} relatively isolated from hydrophobic lipids and running perpendicularly to the membrane plane to allow hydrophilic regions of a polypeptide across. The PCC must also be able to open and close laterally ⁷ to regulate the lipid-mediated

*These authors contributed equally to this work

partitioning of hydrophobic transmembrane helices (TMHs) into the plane of the lipid bilayer⁸.

Low-resolution cryo-electron microscopy (cryo-EM) studies of eukaryotic co-translational translocation complexes^{9–12} have demonstrated that the functional PCC is an oligomer of the Sec61 $\alpha\beta\gamma$ complex. The recent x-ray structure of a non-translocating, monomeric, archaeal SecYE β heterotrimer¹³ – uncomplexed with a ribosome or substrate polypeptide – shows that the α -subunit, SecY, is divided into two independent N-terminal (TMHs 1-5) and C-terminal (TMHs 6-10) halves, forming a ‘clam shell’¹⁴, which on one side is hinged by the loop between TMHs 5/6 and clamped together by the γ -subunit, SecE, while the other side – forming the lateral gate – is unconstrained. Each SecY half contributes to the formation of a transmembrane funnel-like cavity in the center of the complex, which is blocked by a plug (TMH 2a). It is hypothesized that the signal sequence (SS)/TMH signal anchor (SA) of a translocating nascent chain displaces the plug and wedges itself into the lateral gate, thus opening the two SecY halves in the plane of the membrane, enabling either translocation across of a hydrophilic region of nascent chain or the lateral partitioning of a hydrophobic TMH into the lipid bilayer^{13, 14}. Additionally, the x-ray structure shows that the long cytoplasmic loops between TMHs 6/7 and 8/9 extend ~20 Å above the membrane plane for interaction with cytosolic factors, such as the large ribosomal subunit^{15–17}.

We have used a combination of single-particle cryo-EM and computational methods to determine and interpret the structure of a functional, co-translational translocation complex from *E. coli*, comprising SecYEG and a ribosome-nascent chain (RNC) complex. Our cryo-EM reconstruction is of sufficient resolution to determine the precise number of SecYEG monomers in the PCC and to differentiate, using computational methods, between two principal models of SecYEG monomer arrangements, helping to clarify the structural and mechanistic details of co-translational translocation.

RESULTS

Overall structure of the *E. coli* RNC-SecYEG complex

E. coli ribosomes programmed in a cell-free translation system with mRNA encoding for a chimeric nascent chain (Suppl. Fig. 1), which includes the TMH SA of FtsQ and the SecM stalling sequence¹⁸, were complexed with detergent-solubilized SecYEG. These reconstituted RNC-SecYEG complexes were used for structure determination by cryo-EM using single-particle reconstruction. EM data (Fourier shell correlation characteristics ~11 Å at 3 σ , ~15 Å at 0.5) extended out to ~1/10 Å⁻¹ in Fourier space, i.e. structural features down to 10 Å in size could be discerned (see Suppl. Fig. 4). In this reconstruction, tRNAs are visible in all three canonical sites, A, P, and E (Fig. 1a). The path of over 100 nucleotides of the mRNA from its entrance to past its exit site in the 30S subunit can be traced, as can most of the path of the nascent chain in the polypeptide exit tunnel. Greatly improving on the globular appearance of the PCC in previous single-particle cryo-EM studies^{9–12}, detailed rod- and lamella-like features, corresponding to groupings of transmembrane helices in the PCC, are discernible in the current reconstruction (Fig. 1a,b,c). As observed previously there is a large frontal opening between the ribosome and the PCC at the polypeptide exit site, through which the translocating polypeptide is accessible (Fig. 1b, and see Suppl. Discussion 2.1 for a discussion on the discrepancy between existing fluorescence and structural data).

Quite surprising is the observation of a second PCC bound at the exit of the mRNA channel (occupancy ~70%). When viewed perpendicularly to the membrane plane (Fig. 1c,e), the shapes of the two PCCs are different: the PCC at the mRNA channel exit is an elliptic cylinder (110 Å $\times$ 70 Å, height ~45 Å) with pseudo-222 symmetry (Suppl. Fig. 2); whereas the PCC at the polypeptide exit site comprises a circular cylinder (~95 Å diameter, height ~45 Å). Since

the latter PCC contains the TMH SA of FtsQ (see Methods), it represents a translocating state. The PCC at the mRNA channel exit is non-physiologically bound and so must represent a non-translocating state. Volume calculations on the isolated densities of both PCCs indicate that each PCC contains two copies of SecYEG, as also suggested biochemically^{19, 20} (see Suppl. Results 1.1, Suppl. Discussion 2.2 and Suppl. Fig. 5 for a discussion on the effect of low resolution on PCC EM density).

Normal mode-based flexible fitting of PCC atomic coordinates

Two copies of a homology-modeled atomic model of *E. coli* SecYEG (SecYEG_{Ec}) (see Methods) were docked rigidly into the isolated densities of both the non-translocating and translocating PCC. While docking into the translocating PCC density was unsuccessful, a good fit (cross-correlation coefficient (CC) 0.78) to the non-translocating PCC density was achieved when two SecYEG_{Ec} heterotrimers were placed in a front-to-front arrangement. Thus, the structure of the PCC at the mRNA channel exit may represent a physiologically relevant non-translocating state, providing a starting structure for determining the proper conformation of the translocating PCC at the polypeptide exit site. Since relative motions of the N- and C-terminal halves of SecY are postulated to underlie SecYEG function in a translocating PCC^{13, 14}, we opted to utilize a computational method, normal mode-based flexible fitting (NMFF)²¹. Normal modes provide information on the natural vibrations of a molecule and the preferential directions of collective, many-atom displacements, while maintaining the steric and geometric constraints of the overall structure. Normal mode analysis (NMA)²² applied to plug-less SecYEM_j (see Methods) confirmed the expected opening motion of linked SecY halves¹³ (Suppl. Fig. 3). We thus proceeded to use a progression of docking and NMFF steps (see Suppl. Results 1.2 and Suppl. Fig. 4) to obtain a model for the translocating PCC, both in a front-to-front and the recently proposed back-to-back¹³ arrangement.

The front-to-front model fit well (CC_{final} 0.79) into the non-translocating PCC density, while the back-to-back model fit poorly (CC_{final} 0.64). The bulk of each dimeric PCC model could be docked into the translocating PCC density (CC_{initial} 0.65 (front-to-front), 0.61 (back-to-back)) only when the ribosome-binding cytoplasmic loops between TMHs 6/7 and 8/9 (combined cross-section of Cα backbone ~20 Å) of both SecYEG heterotrimers were docked into the connection densities to the ribosome located on the sides (C1,C2), which have a roughly circular cross-section of ~20 Å (back connection (C3): cross-sectional dimensions ~45×20 Å) (Fig. 2b,c,f,g). Upon application of NMFF to these models, the two SecY halves in each heterotrimer in the front-to-front model (CC_{final} 0.79) display significant opening motions within the membrane plane (Fig. 2a,d,i,j), (distance between the centers of mass of linked SecY halves increases by an average of ~5 Å), while no opening motion is seen in the back-to-back model (CC_{final} 0.65). Thus, the formation of a pore wide enough to enable polypeptide translocation, a process hypothesized to be necessary for translocation^{13, 14}, is not observed as in the front-to-front case (Fig. 2a,d,e,h, Suppl. Fig. 4c,d). SecYEG has been shown to be inactivated when SecE is disulfide-crosslinked to a neighboring SecE or SecY molecule, indicating that SecE dynamics, which are restricted when SecE clamps are juxtaposed tightly as in the back-to-back arrangement, are essential for protein translocation^{23, 24}.

Our experimental and computational data thus provide two major arguments – one structural, the other functional – for why the back-to-back model is less likely. Furthermore, key biochemical observations cannot be explained on the basis of a back-to-back arrangement SecYEG heterotrimers, in which the lateral gates in each heterotrimer face away from each other, towards the bulk lipid in the membrane. During post-translational translocation – a process in which a fully translated preprotein is translocated through the PCC in the absence of a ribosome – disulfide-bonded, partially folded regions of preprotein can cross through the PCC channel²⁵. Forming an aqueous pore large enough for this process^{25, 26} would require

the lateral gate in a heterotrimer to open up beyond the point where the wedged SS could sterically block lipid diffusion into the pore, making the translocation of a hydrophilic polypeptide energetically very unfavorable. Similarly, at a certain point during co-translational translocation the SS/SA is no longer associated with SecYEG/Sec61 $\alpha\beta\gamma$ ²⁷, leaving the lateral gate permeable to lipids and making the continuing translocation of a hydrophilic nascent chain energetically very unfavorable. The front-to-front model, however, is consistent with these biochemical observations and certain observed cross-links in SecY²⁰, since the lateral gates of each heterotrimer face each other, and thus a consolidated channel – segregated from lipid – could potentially form.

Details of ribosome-PCC interactions

In order to determine the details of interaction between the PCC and ribosomal elements, an *E. coli* ribosome atomic model was refined into our cryo-EM reconstruction (see Methods). As previously observed^{11, 12} both ribosomal proteins and RNA surround the polypeptide exit site at which the PCC is located. Ribosomal proteins L17, L32 and L22 are located in the front above – and not interacting with – the PCC (Fig. 3a); whereas proteins L24, L29, and L23 interact with the PCC in the back (Fig. 3b). The PCC makes three connections with the ribosome (Fig. 2b,c and Fig. 3c). The connections at the sides, C1 and C2, are interactions between ribosomal RNA (rRNA) hairpins (helix 24 in C1 and helix 59 in C2) and the cytosolic factor-associating domains (CFAD) of SecY, the cytoplasmic regions between TMHs 6/7 and 8/9 (Fig. 3a,c), as has been suggested biochemically^{15–17}. The wall of connection at the back (C3) appears to be mediated mostly through interactions of L29 and L23 with the cytoplasmic region of the SecG subunit and possibly of the N-terminal part of SecE (Fig. 3b,c). Neither the SecG/ β /61 β subunit nor the N-terminal portion of SecE are essential for viability^{28, 29}. However, SecG/ β /61 β has been observed to stabilize the ribosome-PCC complex and facilitate nascent chain translocation^{28, 30}. It is thus likely that connection C3 lends stability and directionality (to be discussed later) to the PCC-ribosome association, but that connections C1 and C2 are required for PCC attachment to the ribosome¹⁶.

Modeling of 121 nucleotides of RNA into the mRNA density isolated from our reconstruction using sequence-specific secondary structure information indicated the presence of two major RNA hairpins at the channel exit, immediately above the CFAD of both heterotrimers of the non-translocating PCC (Fig. 3d). Biochemical competition experiments have demonstrated that mRNA can compete with rRNA for PCC binding¹⁵. This inferred interaction of the CFADs with immobilized, structured mRNA hairpins is analogous to the interaction observed at the polypeptide exit site with rRNA hairpins and may explain the occurrence of the non-translocating PCC at the mRNA channel exit. It is possible that the structure of this PCC may reflect that of a non-translocating PCC when it is bound to the ribosome at the polypeptide exit site.

The path of the translocating nascent chain

Upon NMFF of the dimeric plug-less SecYEG complex into the translocating PCC density, prominent regions of density remain unaccounted for (green and yellow asterisks, Fig. 4a), most notably a long rod-like region (green asterisk) at the front interface of the two heterotrimers. This density most probably corresponds to the nascent chain TMH SA, close to SecY TMH 2 and 7, as suggested by cross-linking experiments³¹. Furthermore, the observation that the interface formed by the two heterotrimers at the front changes significantly from the non-translocating to the translocating state may reflect SA binding at this site (Fig. 2i,j). The other empty regions of density (yellow asterisks) in the middle of both heterotrimers could represent either the plug (TMH 2a) or the remaining, hydrophilic region of the nascent chain. An atomic model was fit into the isolated thread of density corresponding to the nascent chain inside the ribosome, which facilitated the identification of putative interactions with

ribosomal RNA (Mitra *et al.*, manuscript in preparation) and protein. The real-space refined positions of ribosomal protein domains suggest an interaction of the nascent chain with finger-like projections of proteins L4 and L22 in the polypeptide exit tunnel, as has been demonstrated biochemically^{18, 32}. Towards the end of the polypeptide exit tunnel the projection of L23 is in proximity to the nascent chain, indicating another potential interaction site (Fig. 4b,d).

Co-translational translocation through the PCC

The cryo-EM reconstruction of the *E. coli* RNC-SecYEG presented in this work offers glimpses into the architecture of the ribosome-bound PCC in its translocating and possibly also non-translocating state. Analysis of atomic models obtained by NMFF suggests that the PCC is most likely formed by a front-to-front association of two SecYEG/Sec61αβγ heterotrimers in both states. Our data suggest that during the translocation of a hydrophilic region of the nascent chain, neither of the two heterotrimers opens to form one central, consolidated channel, but both retain two segregated pores. The translocating PCC density displays an asymmetry in the distribution of features, with separated rod-like features dominating at the front and contiguous lamella-like features at the back, possibly reflecting greater lipid accessibility at the front (Fig. 1c). Analysis of the atomic model of the translocating PCC supports this hypothesis. The 'hook'-shaped C-terminal SecY half in Sec₁YEG forces a path to the front, to the bulk lipid, whereas in Sec₂YEG the 'hook' dictates a path to the back, which is blocked to lipid by a contiguous wall formed by the TMHs of SecG and SecE (Fig. 2i,j and Fig. 3a,c). An SS/TMH exiting laterally from the Sec₁YEG/Sec61αβγ pore would interact with SecY/61α TMH 2 and 7³¹ at the front interface between both heterotrimers, from where a TMH might then interact with YidC/TRAM^{33, 34} and/or diffuse into the lipid bilayer (Fig. 4a,c,d).

Our NMA results support the initial role of the plug to retain tightly juxtaposed SecY halves in the non-translocating state until the plug is displaced by the nascent chain, at which point the SecY halves can open laterally¹³ (see methods and Suppl. Results 1.2). We suggest that the partially hydrophobic nature of the plug drives its equilibrium state towards burial between linked SecY halves when not sterically obstructed by the presence of a nascent chain, ensuring the maintenance of the permeability barrier even if linked SecY halves are not tightly juxtaposed. Thus, while the nascent chain is being looped through the pore in one heterotrimer, the pore in the other heterotrimer may remain 'plugged'. The frontal alignment of the ribosome/PCC opening with the partition site of nascent chain TMHs at the PCC/membrane may ensure reproducible, efficient folding and release of nascent membrane proteins by providing a dedicated site for the association with the nascent chain and the ribosome-PCC complex of cytosolic proteins, such as chaperones and SecA^{35–37}, through the frontal opening (~20×40 Å, and see Suppl. Discussion 2.1) and membrane proteins at the front PCC/membrane interface. The observation of possible interactions of the projections of ribosomal proteins L4, L22, and L23 with the nascent chain is intriguing and may suggest some form of nascent chain-dependent communication between the ribosome and the PCC.

METHODS

Generation of the *E. coli* RNC-SecYEG complex

E. coli ribosomes were programmed in a cell-free translation system with mRNA encoding for a chimeric nascent chain (Suppl. Fig. 1), consisting of an N-terminal Strep-tag followed by the first 74 residues of *E. coli* FtsQ, which include the TMH signal anchor, and residues 132–170 of *E. coli* SecM, which include the 17 amino acid SecM stalling sequence¹⁸. A nascent chain consisting of FtsQ residues 1–108 has previously been cross-linked to SecY during cotranslational translocation³⁵ and the SecM stalling sequence has been shown to interact tightly with the ribosomal polypeptide exit tunnel while efficiently arresting ribosome elongation¹⁸. Thus, stable RNCs were generated with nascent chains of a uniform length and

sequence without using truncated mRNA. After in vitro translation, the RNCs were affinity-purified via the N-terminal Strep-tag and concentrated by ribosomal pelleting (Suppl. Fig. 1b,c,d). SecYEG was overexpressed in *E. coli*, detergent-solubilized, and purified to homogeneity by standard techniques. The RNC-SecYEG complex was reconstituted and analyzed (Suppl. Fig. 1e). The binding affinity for detergent-solubilized SecYEG was estimated to be ten-fold higher for a ribosome containing a nascent chain compared to an empty ribosome. The presence of the nascent chain thus proved to be essential, since in contrast to eukaryotes, empty *E. coli* ribosomes do not have sufficient affinity (apparent $K_D \sim 2 \mu M$) for the detergent-solubilized PCC to form ribosome-SecYEG complexes under the conditions used for cryo-EM (32 nM). The sample used for cryo-EM contained a ten-fold molar excess of SecYEG over RNCs.

Electron microscopy, image processing, and fitting of atomic ribosome models

Samples were applied to carbon-coated holey grids as published³⁸. Micrographs were recorded under low-dose conditions on a Tecnai F30 field-emission gun electron microscope at 300 kV using a defocus range of 1.5 to 4.3 μm , and scanned on a Zeiss/Imaging Scanner (Z/I Imaging Corporation, Huntsville, AL), corresponding to a pixel size of 3.59 Å on the object scale. The data were analyzed using the SPIDER software package. After automated particle picking followed by visual inspection, supervised classification³⁹ was performed on the starting set of 104,257 picked particles. 53,325 particles corresponding to a translocating PCC bound to the RNC (PCC occupancy 100%) were used for the final, CTF-corrected reconstruction. The falloff of Fourier amplitudes toward higher spatial frequencies was corrected using the x-ray solution scattering intensity distribution of 70S ribosomes from *E. coli*⁴⁰ during each round of refinement. Refinement of an *E. coli* ribosome atomic model into the cryo-EM reconstruction was performed using the program RSRef 41, a real-space refinement module for the TNT program⁴². An almost complete atomic *E. coli* ribosome model obtained by RSRef was adapted from Gao *et al.*⁴³ and complemented with the addition of RNA and protein models from the x-ray structure of the *D. radiodurans* large ribosomal subunit⁴⁴.

Normal mode-based flexible fitting of SecYEG models

Normal mode analysis (NMA) with an elastic network⁴⁵ was performed on the x-ray structure of the SecY β heterotrimer from *Methanococcus jannaschii*¹³. Residues corresponding to the plug (SecY:44-68, *M.j.* (40-75, *E.c.*)) and the Sec β subunit were removed for observation of low-frequency, high-amplitude normal modes corresponding to the opening of SecY halves and the motions of loops (Suppl. Fig. 2). Sec β , which lies at the periphery of the SecY β β_{Mj} structure and makes few interchain contacts, contributes to high-frequency normal modes and masks the low-frequency normal modes corresponding to inter-domain movement. This chain therefore needed to be removed. For normal mode-based flexible fitting (NMFF) a model of *E. coli* SecYEG (SecYEG_{Ec}) was generated by homology-modeling (similarity/identity between SecY of *M. jannaschii* and *E. coli* (40%/20%) and mouse (34%/55%)) based on SecY β β_{Mj} . The additional helices of *E. coli* were modeled according to the placement of helices by van den Berg *et al.*¹³ into the 2-D electron crystal structure of *E. coli* SecYEG determined by Breyton *et al.*⁴⁶. Only the ten lowest-frequency normal modes were utilized to direct flexible fitting of atomic models into the PCC EM density. In order to observe normal modes corresponding to the opening of SecY halves, models of plug-less SecYEG_{Ec} were used in which the two linked SecY halves in each complex were separated by 1 Å on the hingeless side. Only C α coordinates were used for NMFF and NMA, since density for sidechains is not well resolved in the current moderate-resolution EM map. The models obtained upon NMFF were energy-minimized using CNS version 1.1⁴⁷. The reliability of the PCC models generated by NMFF extends to the orientation of domains and the general arrangement of helices within

these domains, but not to the exact positioning of individual helices/residues or the conformation of loops.

STATEMENT OF AUTHOR CONTRIBUTIONS

Grid preparation, cryo-electron microscopy, data processing, atomic model generation, fitting and refinement and interpretation were done by K.M. (laboratory of J.F.). *E. coli* SecYEG-RNC complex was prepared and PCC EM density symmetry analysis performed by C.S. and S.J. (laboratory of N.B.). T.S. (J.F.) assisted in cryo-EM data processing. F.T. (laboratory of C.L.B.III) performed NMA and NMFF.

ACCESSION CODES

Coordinates for the translocating and non-translocating PCC have been deposited in the RCSB, with accession codes 2AKI and 2AKH, respectively. The cryo-EM map of the *E. coli* RNC-SecYEG complex has been deposited in the EBI Macromolecular Structure Database, accession code EMD-1143.

Acknowledgements

We thank R.A. Grassucci for training on the Tecnai F30 electron microscope; G.S. Allen for assistance with supervised classification, O. and discussion of the manuscript; and M. Watters for assistance with the illustrations. This work was supported by HHMI, NSF DBI 9871347 and NIH grants R37 GM29169 and R01 GM 55440 (to J.F.), Multiscale Modeling Tools for Structural Biology (MMTSB) funded by NIH RR12255 (to C.L.B. III), the NCCR Structural Biology program of the Swiss National Science Foundation (SNSF; to N.B.) and a Young Investigator grant from the Human Frontier Science Program (to N.B.). C.S. was supported by post-doctoral fellowships of the Roche Research Foundation and the Ernst Schering Research Foundation.

References

1. Simon SM, Blobel G. A protein-conducting channel in the endoplasmic reticulum. *Cell* 1991;65:371–380. [PubMed: 1902142]
2. Wickner W, Driessen AJM, Hartl FU. The enzymology of protein translocation across the *Escherichia coli* plasma membrane. *Annu Rev Biochem* 1991;60:101–124. [PubMed: 1831965]
3. Brundage L, et al. The purified *E. coli* integral membrane protein SecY/E is sufficient for reconstitution of SecA-dependent precursor protein translocation. *Cell* 1990;62:649–657. [PubMed: 2167176]
4. Gorlich D, Rapoport TA. Protein translocation into proteoliposomes reconstituted from purified components of the endoplasmic reticulum membrane. *Cell* 1993;75:615–30. [PubMed: 8242738]
5. Gilmore R, Blobel G. Translocation of secretory proteins across the microsomal membrane occurs through an environment accessible to aqueous perturbants. *Cell* 1985;42:497–505. [PubMed: 2992801]
6. Simon SM, Blobel G, Zimmerberg J. Large aqueous channels in membrane vesicles derived from the rough endoplasmic reticulum of canine pancreas or the plasma membrane of *Escherichia coli*. *Proc Natl Acad Sci U S A* 1989;86:6176–6180. [PubMed: 2474828]
7. Mothes W, et al. Molecular mechanism of membrane protein integration into the endoplasmic reticulum. *Cell* 1997;89:523–33. [PubMed: 9160744]
8. Hessa T, et al. Recognition of transmembrane helices by the endoplasmic reticulum translocon. *Nature* 2005;433:377–381. [PubMed: 15674282]
9. Beckmann R, et al. Alignment of conduits for the nascent polypeptide chain in the ribosome- Sec61 complex. *Science* 1997;278:2123–6. [PubMed: 9405348]
10. Menetret JF, et al. The structure of ribosome-channel complexes engaged in protein translocation. *Mol Cell* 2000;6:1219–1232. [PubMed: 11106759]
11. Beckmann R, et al. Architecture of the protein-conducting channel associated with the translating 80S ribosome. *Cell* 2001;107:361–372. [PubMed: 11701126]
12. Morgan DG, et al. Structure of the mammalian ribosome-channel complex at 17 Å resolution. *J Mol Biol* 2002;324:871–886. [PubMed: 12460584]

13. van den Berg B, et al. X-ray structure of a protein-conducting channel. *Nature* 2004;427:36–44. [PubMed: 14661030]
14. Rapoport TA, Goder V, Heinrich SU, Matlack KE. Membrane-protein integration and the role of the translocation channel. *Trends Cell Biol* 2004;14:568–575. [PubMed: 15450979]
15. Prinz A, et al. Evolutionarily conserved binding of ribosomes to the translocation channel via the large ribosomal tRNA. *EMBO J* 2000;19:1900–1906. [PubMed: 10775273]
16. Raden D, Song W, Gilmore R. Role of the cytoplasmic segments of Sec61 α in the ribosome-binding and translocation-promoting activities of the Sec61 complex. *J Cell Biol* 2000;150:53–64. [PubMed: 10893256]
17. Cheng Z, Jiang Y, Mandon EC, Gilmore R. Identification of cytoplasmic residues of Sec61p involved in ribosome binding and cotranslational translocation. *J Cell Biol* 2005;168:67–77. [PubMed: 15631991]
18. Nakatogawa H, Ito K. The ribosomal exit tunnel functions as a discriminating gate. *Cell* 2002;108:629–636. [PubMed: 11893334]
19. Bessonneau P, Besson V, Collinson I, Duong F. The SecYEG preprotein translocation channel is a conformationally dynamic and dimeric structure. *EMBO J* 2002;21:995–1003. [PubMed: 11867527]
20. van der Sluis EO, Nouwen N, Driessen AJM. SecY-SecY and SecY-SecE contacts revealed by site-specific crosslinking. *FEBS Lett* 2002;527:159–165. [PubMed: 12220653]
21. Tama F, Miyashita O, Brooks CL III. NMFF: Flexible high-resolution annotation of low-resolution experimental data from cryo-EM maps using normal mode analysis. *J Struct Biol* 2004;147:315–326. [PubMed: 15450300]
22. Go N, Noguti T, Nishikawa T. Dynamics of a small globular protein in terms of low-frequency vibrational modes. *Proc Natl Acad Sci U S A* 1983;80:3696–700. [PubMed: 6574507]
23. Kaufmann A, et al. Cysteine-directed cross-linking demonstrates that helix 3 of SecE is close to helix 2 of SecY and helix 3 of a neighboring SecE. *Biochemistry* 1999;38:9115–9125. [PubMed: 10413486]
24. Veenendaal A, van der Does C, Driessen A. Mapping the sites of interaction between SecY and SecE by cysteine scanning mutagenesis. *J Biol Chem* 2001;276:32559–32566. [PubMed: 11445571]
25. Tani K, Tokuda H, Mizushima S. Translocation of ProOmpA possessing an intramolecular disulfide bridge into membrane vesicles of *Escherichia coli*. Effect of membrane energization. *J Biol Chem* 1990;265:17341–17347. [PubMed: 2211627]
26. Wirth A, et al. The Sec61p complex is a dynamic precursor activated channel. *Mol Cell* 2003;12:261–268. [PubMed: 12887911]
27. Martoglio B, Hofmann MW, Brunner J, Dobberstein B. The protein-conducting channel in the membrane of the endoplasmic reticulum is open laterally toward the lipid bilayer. *Cell* 1995;81:207–214. [PubMed: 7736572]
28. Nishiyama, Ki; Mizushima, S.; Tokuda, H. A novel membrane protein involved in protein translocation across the cytoplasmic membrane of *Escherichia coli*. *EMBO J* 1993;12:3409–3415. [PubMed: 8253068]
29. Schatz PJ, et al. One of three transmembrane stretches is sufficient for the functioning of the SecE protein, a membrane component of the *E. coli* secretion machinery. *EMBO J* 1991;10:1749–1757. [PubMed: 2050112]
30. Kalies KU, Rapoport TA, Hartmann E. The β subunit of the Sec61 complex facilitates cotranslational protein transport and interacts with the signal peptidase during translocation. *J Cell Biol* 1998;141:887–894. [PubMed: 9585408]
31. Plath K, et al. Signal sequence recognition in posttranslational protein transport across the yeast ER membrane. *Cell* 1998;94:795–807. [PubMed: 9753326]
32. Laird V, High S. Discrete cross-linking products identified during membrane protein biosynthesis. *J Biol Chem* 1997;272:1983–9. [PubMed: 8999890]
33. Scotti PA, et al. YidC, the *E. coli* homologue of mitochondrial Oxa1p, is a component of the Sec translocase. *EMBO J* 2000;19:542–549. [PubMed: 10675323]
34. High S, et al. Site-specific photocross-linking reveals that Sec61p and TRAM contact different regions of a membrane-inserted signal sequence. *J Biol Chem* 1993;268:26745–51. [PubMed: 8253810]

35. Valent QA, et al. The *Escherichia coli* SRP and SecB targeting pathways converge at the translocon. EMBO J 1998;17:2504–12. [PubMed: 9564033]
36. Neumann-Haefelin C, Schafer U, Muller M, Koch HG. SRP-dependent cotranslational targeting and SecA-dependent translocation analyzed as individual steps in the export of a bacterial protein. EMBO J 2000;19:6419–6426. [PubMed: 11101515]
37. Zito CR, Oliver D. Two-stage binding of SecA to the bacterial translocon regulates ribosome-translocon interaction. J Biol Chem 2003;278:40640–40646. [PubMed: 12907673]
38. Wagenknecht T, Grassucci R, Frank J. Electron microscopy and computer image averaging of ice-embedded large ribosomal subunits from *Escherichia coli*. J Mol Biol 1988;199:137–147. [PubMed: 2451023]
39. Valle M, al e. Cryo-EM reveals an active role for aminoacyl-tRNA in the accommodation process. EMBO J 2002;21:3557–3567. [PubMed: 12093756]
40. Gabashvili IS, et al. Solution structure of the *E. coli* 70S ribosome at 11.5 Å resolution. Cell 2000;100:537–549. [PubMed: 10721991]
41. Chapman MS. Restrained real-space macromolecular atomic refinement using a new resolution-dependent electron density function. Acta Crystallogr 1995;A51:69–80.
42. Tronrud DE, Ten Eyck LF, Matthews BW. An efficient general-purpose least-squares refinement program for macromolecular structures. Acta Crystallogr 1987;A43:489–501.
43. Gao H, et al. Study of the structural dynamics of the *E. coli* 70S ribosome using real-space refinement. Cell 2003;113:789–801. [PubMed: 12809609]
44. Harms JM, et al. Alterations at the peptidyl transferase centre of the ribosome induced by the synergistic action of the streptogramins dalbavancin and quinupristin. BMC Biol 2004;2:4. [PubMed: 15059283]
45. Tirion MM. Large amplitude elastic motions in proteins from a single-parameter, atomic analysis. Phys Review Lett 1996;77:1905–1908.
46. Breyton C, et al. Three-dimensional structure of the bacterial protein-translocation complex SecYEG. Nature 2002;418:662–665. [PubMed: 12167867]
47. Brunger AT, et al. Crystallography & NMR system: a new software suite for macromolecular structure determination. Acta Crystallogr 1998;D54:905–921.

Figure 1.

General features of the cryo-EM reconstruction of *E. coli* RNC-SecYEG complex. Cryo-EM densities are shown for the small (30S, yellow) and large (50S, sky-blue) subunits, and the A-, P- and E-site tRNAs (magenta, green, and orange). Isolated densities for the mRNA (cyan), the nascent chain in the ribosomal polypeptide exit tunnel (gold), the protein-conducting channel (PCC) at the polypeptide exit site (translocating PCC, blue) and the PCC at the 5' mRNA channel exit (non-translocating PCC, red) are also shown. **a**, General overview of the RNC-SecYEG complex. **b,c**, Closeup of the translocating PCC, shown in the front view (**b**) with the nascent chain visible through the large frontal opening between the ribosome and the PCC (gold arrow). The line of view is perpendicular to the plane of the membrane (**c**). **d, e**, Closeups of the non-translocating PCC.

Figure 2.

Normal mode-based flexible fitting (NMFF) of the SecYEG complex into cryo-EM density. **a,d**, Model after NMFF of a front-to-front arrangement of two SecYEG_{Ec} heterotrimers into the non-translocating (**a**) and translocating (**d**) PCC density. **b,c**, Placement of the non-translocating front-to-front model into the translocating PCC density, such that the ribosome-binding SecY cytoplasmic loops are placed within connection densities. The three connection regions (C1–C3) are shown in cross-section (line of view perpendicular to the membrane plane) (**b**) and in the front view (**c**), with the frontal opening visible between C1 and C2. EM density is shown in cyan as a semi-transparent surface. **e–h**, Parallel analysis of the back-to-back SecYEG_{Ec} model. **i,j**, van der Waals surface representations of the front-to-front models

obtained by fitting to the non-translocating (**i**) and translocating PCC density (**j**) (C-terminal SecY halves transparent). The green and yellow arrows indicate the change in the heterotrimer interface at the front, and linked SecY half opening, respectively. In the atomic models one heterotrimer is colored in shades of blue-green, the other in shades of red. The view is within the plane of the membrane, with the ribosomal side behind the plane. In **a,d,e,h**, atomic models, with TMHs depicted as rods, are shown docked into the experimental EM density (cyan mesh).

Figure 3.

Stereo views of RNA and protein elements in the ribosome-PCC junction. Real-space refined models of *E. coli* ribosomal proteins are rendered as ribbons and ribosomal RNA regions interacting with the PCC as thick, light-grey backbone rattlers. The PCC is colored and rendered as in Fig. 2d with the cryo-EM density in cyan mesh. **ac**, The ribosome-PCC junction at the polypeptide exit site (nascent chain density semi-transparent yellow surface) is shown in the front (**a**) and back (**b**) views. **c**, Ribosomal elements near the polypeptide exit site and parts of the PCC in the three connection regions are illustrated. The line of view is perpendicular to the membrane plane. Connection regions between ribosome and PCC are circled in orange

and labeled. **d**, The non-translocating PCC interacts via its CFADs with hairpins in the mRNA (semi-transparent purple surface with non-interacting mRNA rattler regions in yellow).

Figure 4.

The path of the nascent chain through the ribosome and PCC. **a**, Fitting of a front-to-front SecYEG_{Ec} model into the translocating PCC EM density leaves prominent regions of density unaccounted for (green and yellow asterisks). The PCC is viewed within the plane of the membrane, with the ribosome behind the plane, and colored/rendered as before with TMHs numbered. **b**, The nascent chain (yellow rattler) was fitted into the isolated polypeptide density inside the ribosome. The front view is shown. Schematized versions of the PCC at the polypeptide exit site of the ribosome in views corresponding to **a** (**c**) and **b** (**d**). The PCC and ribosomal elements are colored as before, with the nascent chain in green (TMH signal anchor) and yellow. See text for discussion.