Ruptured ectopic pregnancy after a decline in chorionic gonadotropin

Laurie Montgomery Irvine

J R Soc Med 2006;99:90

The advent of automated assays for beta human chorionic gonadotropin (β -hCG) has made expectant management a realistic option in selected cases of suspected ectopic pregnancy.

CASE HISTORY

A woman of 37, a multigravida, was referred because of continuous and increasingly heavy bleeding since her last menstrual period two weeks earlier and worsening left-sided pelvic pain for the past four days. She had been using a progesterone-only pill for contraception. A home urinary pregnancy test had been positive. On examination she was haemodynamically stable; she had lower abdominal tenderness on the left side without peritonism and a left adnexal mass was felt *per vaginam*. The diagnosis was in doubt but she was judged to have an intrauterine pregnancy. A transvaginal scan showed a thickened endometrium and the possibility of a right simple cyst, there was also some free fluid in the pouch of Douglas. Serum β -hCG was 123 iu/mL. She was followed up as an outpatient, and two further β -hCGs two days apart were 62 and 21 iu/mL; she was due to have a repeat scan at one week. Six days after her initial presentation severe lower abdominal pain developed and she was admitted to hospital. At diagnostic laparoscopy she was found to have a leaking left fimbrial ectopic pregnancy; she also had a right simple ovarian cyst. Partial salpingectomy was performed laparoscopically, and the ectopic pregnancy was confirmed histologically.

COMMENT

Ectopic pregnancy remains an important cause of maternal deaths in the UK; there were 11 fatal cases in the last triennial report.¹ This is despite the advent of a rapid qualitative urine β -hCG assay which if negative virtually excludes ectopic pregnancy. Other investigations include a

quantitative serum β -hCG assay which can be used with transvaginal scanning as part of an algorithm in suspected ectopic pregnancy. In the present case the low and falling β -hCG suggested the pregnancy was unlikely to be ectopic.² Even if a pregnancy is indeed ectopic, a falling β -hCG points to a self-limiting form of pregnancy that requires no intervention.³

Review of the published work indicates a consensus that, whatever the site of pregnancy, a falling β -hCG concentration justifies expectant management.⁴ Moreover, women whose ectopic pregnancy resolved spontaneously have been found to have lower initial β -hCGs (mean 246 iu/L) than those who required surgery (mean 628 iu/L).⁵ Against this must be set rare cases in which a decline in β -hCG was followed by tubal rupture.⁶ Possible mechanisms in these cases are blood-vessel erosion by a surviving clump of trophoblastic tissue (Padwick ML, personal communication) or pressure necrosis from an intraluminal clot.⁶

The special feature of the present case is the exceptionally low level at which symptoms of rupture developed (21 iu/L); the lowest previously reported was 97 iu/L. For clinicians the message is that, when expectant management is decided upon, the patient requires regular hospital review and the β -hCG must be followed down to 'non-pregnant' levels.⁷

REFERENCES

- 1 Confidential Enquiry into Maternal and Child Death. *Why Mothers Die*. London: RCOG Press, 2004
- 2 Dart RG, Mitterando J, Dart LM. Rate of change of serial beta human chorionic gonadotrophin values as a predictor of ectopic pregnancy in patients with indeterminate transvaginal scan findings. *Ann Emerg Med* 1999;34:707–10
- 3 Ankum WM. Diagnosing suspected ectopic pregnancy. *BMJ* 2000;321:1235–6
- 4 Ankum WM, Van der Veen F, Hamerlynck JVT, Lammes FB. Suspected ectopic pregnancy—what to do when human chorionic gonadotrophin levels are below the discriminatory zone. *J Reprod Med* 1995;40:525–8
- 5 Elson J, Tailor A, Banerjee S, Salim R, Hillaby K, Jurkovic D. Expectant management of tubal ectopic pregnancy: prediction of successful outcome using decision tree analysis. *Ultrasound Obstet Gynecol* 2004;23:552–6
- 6 Gretz E, Quagliarello J. Declining serum concentrations of the β -subunit of human chorionic gonadotrophin and ruptured ectopic pregnancy. *Am J Obstet Gynecol* 1987;156:940–1
- 7 Irvine LM, Padwick M. Serial serum HCG measurements in a patient with an ectopic pregnancy: a case for caution. *Hum Reprod* 2000;15:1646–7