REVIEW

Sorting out the complexity of SR protein functions

BRENTON R. GRAVELEY

Department of Genetics and Developmental Biology, University of Connecticut Health Center, Farmington, Connecticut 06030, USA

INTRODUCTION

Members of the serine/arginine-rich (SR) protein family have multiple functions in the pre-mRNA splicing reaction. In addition to being required for the removal of constitutively spliced introns, SR proteins can function to regulate alternative splicing both *in vitro* and *in vivo* (Ge & Manley, 1990; Krainer et al., 1990a; Fu et al., 1992; Zahler et al., 1993a; Caceres et al., 1994; Wang & Manley, 1995). In the cell, SR proteins migrate from speckles—subnuclear domains that may function as storage sites for certain splicing factors—to sites of active transcription (Misteli et al., 1997; Misteli & Spector, 1999) and some SR proteins have been found to shuttle in and out of the nucleus (Caceres et al., 1998). The subcellular localization of SR proteins can be modulated by phosphorylation (Misteli & Spector, 1998; Misteli et al., 1998) and this undoubtedly underlies some regulated splicing events. However, once in the nucleus and localized to the nascent pre-mRNA, exactly how SR proteins engage the general splicing machinery to recognize specific splice sites is unclear and is an area of intense investigation.

All SR proteins have a modular organization and contain an N-terminal RNA-binding domain that interacts with the pre-mRNA and a C-terminal RS domain that functions as a protein interaction domain. SR proteins have been proposed to function by binding to the pre-mRNA and recruiting a number of different general splicing factors to the pre-mRNA during spliceosome assembly (Wu & Maniatis, 1993; Kohtz et al., 1994; Roscigno & Garcia-Blanco, 1995; Zuo & Maniatis, 1996). In addition, SR proteins are thought to mediate interactions between splicing factors bound to the 5' and 3' splice sites (Wu & Maniatis, 1993). However, it has not

been clear whether all of these activities occur during the removal of each intron. Recent studies now suggest that all of the proposed SR protein functions are carried out during each round of splicing, and at least some of these functions are performed by independent SR protein molecules. This review discusses recent advances in understanding the diverse functions of SR proteins in metazoan pre-mRNA splicing and presents a model that takes these new findings into account. Although the reader should keep in mind that the activity of SR proteins *in vivo* can be influenced by modulating their subcellular localization, this review focuses on biochemical experiments that specifically address the mechanisms by which SR proteins directly function in the splicing reaction. (The reader may also refer to related reviews recently published elsewhere (Tacke & Manley, 1999; Blencowe, 2000).)

SPLICEOSOME ASSEMBLY

Considerable progress has been made in the understanding of spliceosome assembly (Reed & Palandjian, 1997; Burge et al., 1999). The spliceosome consists of five small nuclear ribonucleoprotein particles (snRNPs), designated U1, U2, U4, U5, and U6, and a large number of non-snRNP proteins. Spliceosome assembly is directed, in part, by the RNA sequences at the splice sites. In mammals, the 5' splice site consensus sequence is AG/GURAGU (where / denotes the exon/intron boundary). Three distinct sequence elements are found at the 3' splice site—the branchpoint (YNYURAC), a polypyrimidine tract, and the actual 3' splice site (YAG/N). Spliceosome assembly proceeds in a step-wise fashion (Fig. 1) and is initiated upon the binding of U1 snRNP to the 5' splice site, SF1/mBBP to the branchpoint sequence and U2 auxiliary factor (U2AF) to the pyrimidine tract and 3' AG to form the E, or early, complex. Subsequently, U2 snRNP binds to the branchpoint to form A complex, followed by the

Reprint requests to: Brenton R. Graveley, Department of Genetics and Developmental Biology, University of Connecticut Health Center, 263 Farmington Avenue, Farmington, Connecticut 06030, USA; e-mail: graveley@neuron.uchc.edu.

FIGURE 1. Spliceosome assembly pathway. In the first step of spliceosome assembly, U1 snRNP binds to the 5' splice site (Michaud & Reed, 1991), SF1/mBBP binds to the branchpoint (Abovich & Rosbash, 1997), and U2 auxiliary factor (U2AF) binds to the pyrimidine tract (Ruskin et al., 1988; Zamore & Green, 1989; Bennett et al., 1992) and 3' YAG (Merendino et al., 1999; Wu et al., 1999; Zorio & Blumenthal, 1999). This complex commits the pre-mRNA to the splicing pathway and is called the E, or early, complex (Michaud & Reed, 1991). Next, E complex is converted to A complex when the U2 snRNP binds to the branchpoint (Bennett et al., 1992). Subsequently, B complex is formed when the U4, U5, and U6 snRNPs enter the spliceosome as a tri-snRNP particle (Reed & Palandjian, 1997). Finally, a massive rearrangement occurs in which U6 replaces U1 at the 5' splice site, U6 and U2 interact, U5 bridges the splice sites, and U1 and U4 become destabilized. This rearranged spliceosome is called the C complex and is catalytically active.

association of the U4/U6•U5 tri-snRNP to form B complex. Next, the spliceosome rearranges to form the catalytically active C complex. Some of the SR protein functions, such as the recruitment of U1 snRNP and U2AF to the 5' and 3' splice sites, respectively, are thought to act at the stage of E complex formation

(Reed, 1996). SR proteins also appear to function at later steps, such as the transition from A to B complex (Roscigno & Garcia-Blanco, 1995) and even perhaps after the first catalytic step of splicing (Chew et al., 1999).

THE SR PROTEIN FAMILY OF SPLICING FACTORS

SR proteins were independently discovered by a number of groups taking very different approaches. For instance, SF2/ASF (Splicing Factor 2/Alternative Splicing Factor; Ge et al., 1991; Krainer et al., 1991) was purified from HeLa cell nuclear extract as a factor required to reconstitute splicing in S100 splicing-deficient HeLa cell extract (Krainer et al., 1990b) and to induce splice site switching (Ge & Manley, 1990). In another approach, monoclonal antibodies directed against purified spliceosomes were used to identify SR proteins. In one case, this led to the identification of SC35 (Fu & Maniatis, 1990, 1992b). Another group, using the monoclonal antibody mAb104, identified an entire family of related proteins—including SC35 and SF2/ASF—that they termed the SR protein family based on their high serine and arginine content (Roth et al., 1991; Zahler et al., 1992). Subsequently, additional members of the SR protein family were identified (Ayane et al., 1991; Champlin et al., 1991; Diamond et al., 1993; Zahler et al., 1993b; Cavaloc et al., 1994; Sreaton et al., 1995; Zhang & Wu, 1996; Soret et al., 1998), and the human SR protein family currently contains 10 known members (Fig. 2A). Although SR proteins have been identified in all metazoan species examined and plants (Lazar et al., 1995; Lopato et al., 1996a, 1996b, 1999; Golovkin & Reddy, 1998, 1999), they are not found in all eukaryotes. For example, *Schizosaccharomyces pombe* contains at least two SR proteins (Gross et al., 1998; Lutzberger et al., 1999), while *Saccharomyces cerevisiae* contains none.

Until recently, metazoan SR proteins were thought to be encoded by essential genes. For instance, in *Drosophila melanogaster*, deletion of the gene encoding the B52 protein, which corresponds to human SRp55, results in lethality in the first or second instar larval stage (Ring & Lis, 1994). Likewise, conditional depletion of the ASF/SF2 protein in chicken DT40 B-cells results in cell death (Wang et al., 1996). Moreover, a late embryonic lethal phenotype was observed when the *Caenorhabditis elegans* SF2/ASF homolog, *rsp-3*, was targeted by RNAi (Longman et al., 2000). Surprisingly, no phenotype was observed when six other *C. elegans* SR protein genes were individually targeted by RNAi (Longman et al., 2000). However, when multiple *C. elegans* SR protein genes were simultaneously targeted by this method, developmental defects or lethality was observed (Longman et al., 2000). It will be important to determine the reasons why some SR pro-

A. Human SR Proteins

B. Human SR Related Proteins

U2 Auxiliary Factor

snRNP Components

Splicing Regulators

Splicing Coactivators

RNA Helicases

Protein Kinases

FIGURE 2. Schematic diagram of human SR proteins and SR related proteins. **A:** The domain structures of the known members of the human SR protein family are depicted. RRM: RNA recognition motif; RRMH: RRM homology; Z: zinc knuckle; RS: arginine/serine-rich domain. **B:** The domain structures for some of the human SRps that participate in pre-mRNA splicing are depicted. All proteins, with the exception of SRm300, are drawn to scale. RRM: RNA recognition motif; RS: arginine/serine-rich domain; Zn: zinc finger; DEXD/H Box: motif characteristic of RNA helicases.

teins are not essential. Do the nonessential SR proteins simply not participate in the splicing of essential genes, or can other SR proteins functionally substitute for the missing SR protein? The absence of an obvious phenotype in the worms treated with dsRNA, however, does not exclude the possibility that certain SR proteins are completely dispensable. For example, some SR proteins may be required for the proper function of specific neurons, and worms lacking these SR proteins may display subtle phenotypes that would be difficult to assess without rigorous behavioral or electrophysiological assays. Complete sets of knockouts in other metazoans must be created and analyzed to establish the generality of the results obtained in worms. The recent completion of the genomic DNA sequence of *D. melanogaster* should greatly facilitate these types of experiments.

Structural organization of SR proteins

SR proteins have a characteristic structural organization (Fig. 2A). They contain one or two RNA-binding domains at their N-termini and a variable-length arginine/

serine-rich domain at their C-termini (the RS domain). The RNA-binding and RS domains are modular. For example, the RNA-binding domains can be exchanged between SR proteins (Chandler et al., 1997; Mayeda et al., 1999) and can bind RNA in the absence of the RS domain (Caceres & Krainer, 1993; Zuo & Manley, 1993; Tacke & Manley, 1995). Likewise, the RS domains can be exchanged between SR proteins (Chandler et al., 1997; Wang et al., 1998b) and can even function when fused to a heterologous RNA-binding domain (Graveley & Maniatis, 1998).

A number of additional RS-domain-containing proteins, distinct from the SR proteins, are involved in various aspects of pre-mRNA metabolism and are collectively referred to as SR-related proteins or SRps (Fu, 1995; Blencowe et al., 1999). Examples include both subunits of U2AF, [U2AF³⁵ (Zhang et al., 1992) and U2AF⁶⁵ (Zamore et al., 1992)], snRNP components [(U1-70K (Query et al., 1989), U5-100K (Teigelkamp et al., 1997), U4/U6, U5-27K (Fetzer et al., 1997), and hLuc7p (Fortes et al., 1999; Nishii et al., 2000)], splicing regulators [Tra (Boggs et al., 1987), Tra2 (Amrein et al., 1988), hTra2α, hTra2β (Dauwalder

et al., 1996; Beil et al., 1997), and SWAP (Denhez & Lafyatis, 1994; Spikes et al., 1994)], splicing coactivators [SRm160/SRm300 (Blencowe et al., 1998, 2000)], RNA helicases [hPrp16 (Zhou & Reed, 1998), HRH1 (Ono et al., 1994), and Hel117 (Sukegawa & Blobel, 1995)], and protein kinases [Clk/Sty 1, 2, and 3 (Johnson & Smith, 1991; Hanes et al., 1994)]. Some of the human SRps that participate in pre-mRNA splicing are depicted in Figure 2B. Interestingly, additional SRps have been identified, and these proteins participate in a number of processes other than pre-mRNA splicing. For a detailed review on SRps see Blencowe et al. (1999).

The RNA-binding domain

To fully understand how SR proteins function, it is critical to know where they bind to the pre-mRNA. A number of studies over the last few years have therefore attempted to elucidate the RNA-binding specificities of each SR protein (Tacke & Manley, 1995; Shi et al., 1997; Tacke et al., 1997, 1998; Liu et al., 1998, 2000; Cavaloc et al., 1999; Schaal & Maniatis, 1999b). A surprising conclusion from these experiments is that SR proteins recognize a vast array of RNA sequences (see Table 1). Although SR proteins do display distinct RNA-binding specificities, the consensus sequences that they recognize are rather degenerate. In fact, in several cases, sequences identified as binding sites for one SR protein can also be recognized by other SR proteins (Tacke & Manley, 1995; Liu et al., 1998; Tacke

et al., 1998). The overlapping and promiscuous RNA-binding specificities of SR proteins may partially account for their apparent redundancy in their function. The possible significance of the degenerate SR protein recognition sequences will be discussed in more detail below.

The RS domain

It is now widely accepted that the RS domains of SR proteins function as protein interaction domains. SR proteins were initially shown to interact with one another, the splicing regulators Tra and Tra2, and with other components of the splicing machinery that contain RS domains such as U1-70K and U2AF³⁵ (Wu & Maniatis, 1993; Amrein et al., 1994; Kohtz et al., 1994). Importantly, these protein interactions all require the RS domains of each protein (Wu & Maniatis, 1993; Amrein et al., 1994; Kohtz et al., 1994). Only recently, however, have the RS domains been shown to be sufficient to mediate protein interactions. For instance, the RS domain of SF2/ASF has been shown to be sufficient to interact with RSF1, a *Drosophila* splicing repressor (Labourier et al., 1999). In addition, when artificially tethered to the pre-mRNA, the RS domains of several human SR proteins are sufficient to activate enhancer-dependent splicing (Graveley & Maniatis, 1998), an activity that presumably requires protein interactions. In contrast, the RS domain of SF2/ASF is unable to interact with U1-70K (Xiao & Manley, 1997). These studies indicate that the RS domains of SR pro-

TABLE 1. RNA sequences identified as SR protein-binding sites by in vitro selection methods.

Protein	Binding site ^a	Method ^b	Reference
SF2/ASF	RGAAGAAC AGGACRRAGC SRSASGA	SELEX	Tacke & Manley, 1995
SC35	AGSAGAGUA GUUCGAGUA UGUUCSAGWU GWUWCCUGCUA GGGUAUGCUG GAGCAGUAGKS AGGAGAU GRYYCSYR	Functional SELEX	Liu et al., 1998 Tacke & Manley, 1995
9G8	AGACKACGAY ACGAGAGAY	SELEX	Cavaloc et al., 1999
SRp40	UGGGAGCRGUYRGCUCGY ACDGS	SELEX Functional	Tacke et al., 1997 Liu et al., 1998
SRp55	USCGKM	SELEX	Liu et al., 1998
B52	GRUCAACCDNGGCGAACNG	SELEX	Shi et al., 1997
hTra2 β	(GAA) _n	SELEX	Tacke et al., 1998

^aN: any nucleotide; R: purine; Y: pyrimidine; S: G or C; K: U or G; W: A or U; D: A, G, or U; M: A or C.

^b"SELEX" indicates that the RNA sequence was determined to be a high affinity binding site for a purified SR protein; "functional" indicates that the RNA sequence was determined to function as an SR protein-specific splicing enhancer.

teins are indeed protein interaction domains, but suggest that different protein interactions may have distinct RS domain sequence requirements.

The modular domain organization of SR proteins has made it possible to assess the specificity of RS domain function in a variety of contexts by exchanging RS domains between different SR proteins. For example, it was found that the RS domains from other SR proteins, as well as the splicing regulator Tra2, can substitute for the SF2/ASF RS domain in chicken cells lacking endogenous SF2/ASF (Wang et al., 1998b). Likewise, it was found that several RS domains can replace the Tra2 C-terminal RS domain in *D. melanogaster* (Dauwalder & Mattox, 1998). Moreover, the RS domain of Tra can also substitute for the SWAP RS domain in *Drosophila* (Li & Bingham, 1991). Several in vitro studies have also been used to analyze RS domain specificity. Chandler et al. (1997) found that exchanging the RS domains of SF2/ASF and SC35 had no effect on the ability of the protein to complement splicing-deficient S100 extracts, suggesting that RS domains are fully interchangeable. However, when the RS domains of six different human SR proteins were tested for their ability to activate splicing when artificially tethered to the pre-mRNA, it was found that the activity of RS domains differed quantitatively (Graveley et al., 1998). In fact, the potency of an RS domain varies directly with the number of RS dipeptides present (Graveley et al., 1998). In summary, these results show that the RS domains of SR proteins are, for the most part, fully interchangeable and seem to differ from one another only quantitatively. However, subtle differences not detected in these rather crude assays could lead to profound differences in the animal.

RS domain phosphorylation

A number of studies have shown that the serine residues within the RS domains of SR proteins are extensively phosphorylated. This phosphorylation appears to influence the subcellular localization of SR proteins (Gui et al., 1994a; Colwill et al., 1996) as well as protein interactions involving SR proteins (Wang et al., 1998a; Xiao & Manley, 1998), both of which may change the ability of SR proteins to function in splicing. In fact, it has recently been shown that both hyper- and hypophosphorylated SR proteins are unable to support splicing (Kanopka et al., 1998; Prasad et al., 1999; Sanford & Buzik, 1999). The modulation of phosphorylation levels appears to be used to control the activity of SR proteins in adenovirus infection (Kanopka et al., 1998) and early development in the nematode *Ascaris lumbricoides* (Sanford & Buzik, 1999).

Three types of protein kinases have been identified that can phosphorylate RS domains in vitro. These include the SR protein kinase family [SRPK1 (Gui et al., 1994b) and SRPK2 (Wang et al., 1998a)], the Clk/Sty

family (see, for example, Colwill et al., 1996), and DNA topoisomerase I (Rossi et al., 1996). The most direct evidence for an in vivo role of SR protein kinases comes from an elegant series of experiments in *D. melanogaster*. *Drosophila* contains a homolog of Clk/Sty called *Doa*. DOA was found to phosphorylate the *Drosophila* SR protein RBP1, as well as Tra and Tra2, in vitro, and RBP1 in vivo (Du et al., 1998). Interestingly, *Doa* mutations disrupt the splicing of the *doublesex* pre-mRNA, but have no effect on the splicing of the *fruitless* pre-mRNA (Du et al., 1998)—both targets of regulation by Tra, Tra2, and RBP1. This result is of particular interest because Tra, Tra2, and RBP1 control alternative 3' splice site selection in the *doublesex* pre-mRNA, and alternative 5' splice site selection in the *fruitless* pre-mRNA. This observation suggests that the regulation of alternative 5' and 3' splice sites by SR proteins have different requirements for RS domain phosphorylation. In addition, these results demonstrate that an SR protein kinase can control the developmentally regulated alternative splicing of specific pre-mRNAs.

THE FUNCTIONS OF SR PROTEINS IN PRE-mRNA SPLICING

Despite the plethora of proposed SR protein functions, they can essentially be divided into two categories—exon dependent and exon independent.

The exon-dependent functions of SR proteins

Perhaps the best-characterized functions of SR proteins are those that involve binding to exon sequences. The exon-dependent functions can be subdivided into regulated 3' splice site selection, regulated 5' splice site selection, and constitutive functions. Although most of the work in this area has focused on how SR proteins function to regulate alternative splicing, it is now clear that the binding of SR proteins to constitutive exons also plays an important role in the splicing reaction.

Regulated 3' splice site selection

Soon after SR proteins were discovered, it was appreciated that they function not only as essential splicing factors, but also as splicing regulators. A flurry of papers demonstrated that SR proteins could bind to exon sequences, where they functioned to enhance the splicing of the adjacent intron (see, for example, Lavigne et al., 1993; Sun et al., 1993; Tian & Maniatis, 1993). These RNA elements have since been termed exonic splicing enhancers and have been identified in a number of metazoan exons. A well-characterized example of an exonic splicing enhancer is provided by the *D. melanogaster* sex-determination gene *doublesex*, which is alternatively spliced in males and females (Baker,

1989; Burtis & Baker, 1989; Hertel et al., 1997). In males, exon 4 is skipped, whereas in females, exon 4 is included. This splicing event requires a complex exonic splicing enhancer in exon 4 that consists of six 13-nt repeats, collectively referred to as the *dsx* repeat element (*dsxRE*). Each repeat of the *dsxRE* is recognized by the SR protein RBP1 and the splicing regulator Tra2 (Lynch & Maniatis, 1996). However, the binding of these factors to the splicing enhancer requires the presence of the splicing regulator Tra, which is expressed only in females. Tra, Tra2, and RBP1 interact with each other to cooperatively bind the *dsx* splicing enhancer (Lynch & Maniatis, 1996). This splicing enhancer complex, in turn, functions to activate the splicing of the upstream intron. The *dsx* splicing enhancer has served as a paradigm for regulated splicing enhancers.

The most controversial question in the SR protein field is how enhancer-bound SR proteins function to activate splicing, and several mechanisms have been proposed. The observations that many enhancer-dependent introns contain suboptimal pyrimidine tracts and that SR proteins can physically interact with U2AF³⁵ (Wu & Maniatis, 1993) led to the proposal that splicing enhancers function by recruiting U2AF⁶⁵ to the adjacent pyrimidine tract of regulated introns (Tian & Maniatis, 1993; Wu & Maniatis, 1993) (Fig. 3A). Subsequent *in vitro* experiments using purified recombinant proteins have shown that SR proteins can, in fact, recruit U2AF⁶⁵ to the upstream intron and that this recruitment requires U2AF³⁵ (Zuo & Maniatis, 1996). Similar results have also been obtained in experiments performed in nuclear extracts (Wang et al., 1995; Bouck et al., 1998; B.R. Graveley, K. Hertel, & T. Maniatis, unpubl. data). However, other studies in nuclear extracts have failed to detect a difference in U2AF binding in the presence or absence of a splicing enhancer, raising the possibility that enhancer-bound SR proteins may act through an alternative mechanism (Kan & Green, 1999; Li & Blencowe, 1999). For instance, it has been suggested that splicing enhancer complexes may function in conjunction with the splicing coactivator SRm160/300 (Eldridge et al., 1999) (Fig. 3B). In at least one case, splicing enhancers may function to counteract splicing inhibitory complexes (Kan & Green, 1999; Fig. 3C). Finally, although splicing enhancers appear to function primarily at the early stages of spliceosome assembly—prior to, or at the step of splice site selection—one set of recent experiments indicates that splicing enhancers may also function to enhance the second step of splicing (Chew et al., 1999).

Although SRm160/300 clearly participates in enhancer-dependent splicing and some enhancers may function in part by negating the effects of splicing inhibitors, at least one set of experiments is difficult to reconcile with models that do not invoke U2AF recruitment. It has been shown for several enhancer-

dependent introns that improving the weak pyrimidine tract can relieve the requirement for a splicing enhancer (Tian & Maniatis, 1994; Lorson & Androphy, 2000; B.R. Graveley, K. Hertel, & T. Maniatis, unpubl. results). For example, replacing the normal pyrimidine tracts of either the *Drosophila dsx* or the mouse IgM pre-mRNAs with sequences that have a 5–10-fold higher affinity for U2AF render those RNAs enhancer independent (B.R. Graveley, K. Hertel, & T. Maniatis, unpubl. data). Likewise, decreasing the strength of a pyrimidine tract of a constitutively spliced pre-mRNA, and therefore the affinity of U2AF, can make that intron enhancer dependent (Tian & Maniatis, 1992; Graveley & Maniatis, 1998). Thus, the requirement for a splicing enhancer is directly related to the intrinsic affinity of U2AF for the pyrimidine tract. The simplest explanation for these results is that splicing enhancers do indeed function to recruit U2AF. The failure to observe differences in U2AF binding in some cases may be a reflection of differences in experimental systems and the highly cooperative nature of spliceosome assembly. It is also possible that U2AF recruitment is not sufficient to activate splicing. Future experiments must more clearly address the detailed mechanism of enhancer function and definitively identify the target for the putative protein–protein interactions between enhancer complexes and the splicing machinery.

Regulated 5' splice site selection

In addition to promoting the recognition of alternative 3' splice sites, splicing enhancers also appear to promote the recognition of alternative 5' splice sites. Again, the best example comes from a *Drosophila* gene—*fruitless*. Just as in the case of *dsx* pre-mRNA, the *fruitless* pre-mRNA is alternatively spliced in males and females, except the latter case involves the selection of alternative 5' splice sites (Ryder et al., 1996). An analysis of the sequence of the *fruitless* gene revealed the presence of three nearly perfect copies of the *dsxRE* 13-nt repeats, immediately upstream of the female-specific 5' splice site (Ryder et al., 1996). This observation suggests that the same splicing enhancer complex that regulates *dsx* alternative splicing may regulate *fruitless* alternative splicing. In fact, Tra and Tra2 mutant females fail to utilize the female-specific 5' splice site (Ryder et al., 1996). This result directly implicates the Tra, Tra2, RBP1 heterotrimeric complex in 5' splice site selection in the *fruitless* pre-mRNA. Importantly, three other recent examples suggest that SR proteins bound to upstream splicing enhancers stimulate downstream 5' splice site utilization in other pre-mRNAs (Bourgeois et al., 1999; Côté et al., 1999; Selvakumar & Helfman, 1999). These results strongly suggest that upstream splicing enhancer complexes function to stimulate the binding of U1 snRNP to the downstream 5' splice site through an interaction with U1-70K (Fig. 3D).

FIGURE 3. The regulated exon-dependent functions of SR proteins. **A:** U2AF recruitment model. An enhancer-bound SR protein interacts with the RS domain of U2AF³⁵, thereby recruiting U2AF⁶⁵ to the pre-mRNA. **B:** Coactivator model. In this model, the splicing enhancer functions through interactions with the splicing coactivator SRm160/300, which also interacts with U1 snRNP and U2 snRNP. Some of these interactions may be indirect. **C:** Inhibitor model. In the absence of the enhancer-bound SR protein, a downstream splicing inhibitor functions to prevent the splicing of the upstream intron. The function of the splicing enhancer is to counteract the splicing inhibitor. **D:** Recruitment of U1 snRNP to a 5' splice site. An SR protein bound to the upstream exon interacts with U1-70K and recruits U1 snRNP to the 5' splice site.

However, it remains to be shown if this is the mechanism by which upstream SR proteins promote the recognition of the 5' splice site.

The constitutive functions

In addition to their exon-dependent role in regulating alternative splicing, SR proteins have recently been shown to have an exon-dependent function in constitutive splicing. The exon-definition model (Berget, 1995) proposes that interactions between components bound

to the 5' and 3' splice sites flanking an exon serve to distinguish exons, which are typically small, from introns, which are typically large. SR proteins have been proposed to participate in this process by binding to the exon where they simultaneously interact with U2AF³⁵ bound to the upstream 3' splice site and the 70K protein of U1 snRNP bound to the downstream 5' splice site. It is thought that the majority of constitutively spliced exons are defined by this mechanism (Reed, 1996). If correct, constitutively spliced exons should contain binding sites for one or more SR proteins. In support of this

model, a number of SR protein binding sites have recently been identified in constitutive exons (Mayeda et al., 1999; Schaal & Maniatis, 1999a). In fact, these sites have been shown to function as constitutive splicing enhancers (Mayeda et al., 1999; Schaal & Maniatis, 1999a).

Thus it appears that SR proteins have similar exon-dependent functions in regulated and constitutive splicing. In constitutive splicing, the SR proteins bind to the exon constitutively, whereas in alternative splicing the SR protein association is regulated. The functional interactions between SR proteins and constitutive exons place a significant constraint on the types of RNA sequences that can be present in coding exons, because codons and SR protein binding sites would overlap. Perhaps the degenerate RNA binding specificities of SR proteins have evolved to cope with this functional conflict. It is interesting to note that for *doublesex* and *fruitless*—two alternative splicing events that are essential to the survival of the species—the splicing enhancers, which are highly conserved, are present in untranslated regions. Thus, in instances where a high degree of RNA-binding specificity is necessary to achieve proper regulation, it may not be possible for splicing enhancers to coexist with coding sequence.

The exon-independent functions of SR proteins

In addition to their exon-dependent functions, SR proteins have some exon-independent functions. Most of these functions are considered exon-independent simply because it is not known if SR proteins must bind to exons to perform these functions. However, a recent study has shown that at least some essential SR protein functions do not require binding to exon sequences. Based in part on the observation that the second exon is not necessary for the first step of splicing (Smith et al., 1989; Anderson & Moore, 1997), Hertel and Maniatis (1999) performed an interesting set of experiments in which they determined the minimal number of nucleotides in the first exon that are required for the first step of splicing. They found that a pre-mRNA that contains only a single nucleotide in the first exon, and completely lacks the second exon, can undergo the first step of splicing. Interestingly, SR proteins are required for this splicing reaction with the minimal substrate. Moreover, all SR proteins tested in this reaction could promote splicing, indicating that there is functional redundancy with regards to the exon-independent function(s) of SR proteins. Thus, SR proteins perform some essential function(s) that does not require their binding to exons—perhaps mediating interactions between the 5' and 3' splice sites.

Cross-intron interactions

SR proteins have been proposed to play an important role in the pairing of 5' and 3' splice sites (Fig. 4A). This idea was originally based on the observation that the SR protein SC35 can promote an interaction between U1 and U2 snRNPs on the pre-mRNA (Fu & Maniatis, 1992a). Further evidence for this model was provided by the observation that SR proteins can simultaneously interact with U1-70K and U2AF³⁵, components bound to the 5' and 3' splice sites respectively (Wu & Maniatis, 1993). In addition, SR proteins are required for *trans*-splicing, a reaction in which the 5' and 3' splice sites are contained on separate RNA molecules (Bruzik & Maniatis, 1995; Chiara & Reed, 1995). Although SR proteins clearly have the potential to function in splice site pairing, it remains to be determined whether SR proteins actually participate in this process in the spliceosome.

Tri-snRNP incorporation

SR proteins have also been shown to play a role in the recruitment of the U4/U6•U5 tri-snRNP into the spliceosome (Roscigno & Garcia-Blanco, 1995; Tarn & Steitz, 1995) (Fig. 4A). Although the precise role of SR proteins in this activity has not been determined, it is most likely that this recruitment is mediated by an interaction between an SR protein present in the partially assembled spliceosome and a component of the tri-snRNP. There are at least two proteins within the human U4/U6•U5 tri-snRNP that contain RS domains—U4/U6•U5-27K (Fetzer et al., 1997) and U5-100K (Teigelkamp et al., 1997)—and are therefore potential interaction targets of SR proteins.

5' splice site recognition

As described earlier, enhancer-bound SR proteins can promote the utilization of a downstream 5' splice site, and this activity may involve the recruitment of U1 snRNP to the 5' splice site. However, SR proteins may also mediate U1 snRNP recruitment by a mechanism that does not involve binding to exon sequences. Experiments performed with purified components have shown that SF2/ASF can recruit U1 snRNP to an RNA containing a 5' splice site (Kohtz et al., 1994; Jamison et al., 1995). This activity requires an intact 5' splice site sequence, and at least one functional RNA binding domain and the RS domain of SF2/ASF (Kohtz et al., 1994; Jamison et al., 1995). Moreover, SF2/ASF has been shown to bind RNAs containing wild-type, but not mutant 5' splice sites (Zuo & Manley, 1994). Together, these results suggest that SF2/ASF binds directly to the 5' splice site, interacts with U1-70K, and recruits U1 snRNP (Fig. 4B). However, many details of this model still need to be resolved, such as whether SF2/

FIGURE 4. The exon-independent functions of SR proteins. **A:** Model for the splice site pairing and tri-snRNP recruitment activities of SR proteins. It is unknown whether these activities of SR proteins require RNA binding and, if so, where the SR protein binds. Therefore, the SR protein is not depicted bound to the pre-mRNA. The cross-intron bridging function is thought to be mediated by a simultaneous interaction of the SR protein with U1-70K and U2AF³⁵. SR proteins have been shown to participate in the recruitment of the U4/U6•U5 tri-snRNP, although the mechanism by which they do so has not been elucidated. Presumably this activity is mediated by an interaction between an SR protein present in the partially assembled spliceosome and a component of the tri-snRNP. Two candidate interaction targets of SR proteins are the U5-100K and U4/U6•U5-27K protein as depicted here. It should be noted, however, that these protein interactions are speculative and have not yet been demonstrated. **B:** An SR protein bound to the 5' splice site recruits U1 snRNP. This recruitment is mediated by an interaction between the SR protein and the 70K component of U1 snRNP. U1 snRNA then base pairs with the 5' splice site. It remains to be determined whether the SR protein can bind to the pre-mRNA:U1 snRNA duplex.

ASF can actually bind to the pre-mRNA:U1 snRNA duplex, and if not, where SF2/ASF binds to carry out this function.

SR proteins may also participate in the recognition of the 5' splice site after U1 snRNP binding. This insight came from experiments in which U1 snRNP was either debilitated by antisense oligonucleotides or depleted from nuclear extracts (Crispino et al., 1994; Tarn & Steitz, 1994). Extracts treated by either method are unable to support splicing on their own. However, the addition of high concentrations of SR proteins can re-

store splicing (Crispino et al., 1994; Tarn & Steitz, 1994). Thus, SR proteins appear to have the ability to bypass the requirement for U1 snRNP in 5' splice site selection. Subsequent experiments have suggested that the added SR proteins assist U6 snRNP in recognizing the 5' splice site (Crispino & Sharp, 1995; Tarn & Steitz, 1995)—a step that normally occurs late in spliceosome assembly. However, this function of SR proteins may be indirect. SR proteins have been shown to recruit the U4/U6•U5 tri-snRNP to the spliceosome (Rosignio & Garcia-Blanco, 1995; Tarn & Steitz, 1995). Thus, the

high concentrations of SR proteins added to U1 snRNP-depleted extracts may increase the number of spliceosomes that contain the U4/U6•U5 tri-snRNP. This, in turn, would increase the probability that U6 snRNA could functionally recognize the 5' splice site in a step that does not require SR proteins. Additional studies are required to determine whether SR proteins actually play a direct role in the U6 snRNP-5' splice site interaction in spliceosomes that contain U1 snRNP, and if so, how this is achieved.

MULTIPLE SIMULTANEOUS FUNCTIONS

Aside from the obvious question of how SR proteins function from a mechanistic point of view, there are a number of important questions that have not been given much consideration until recently. For instance, are all of the proposed functions of SR proteins performed during the removal of each intron or do different introns require SR proteins to perform different subsets of these functions? If all of the functions are carried out for each intron, does a single SR protein molecule perform all of these functions, or do different SR protein molecules carry out each function? At one extreme, it is possible that a single SR protein molecule bound to the downstream exon can simultaneously interact with U2AF, U1-70K, and a component of the U4/U6•U5 tri-snRNP. These interactions would serve to define and pair the 5' and 3' splice sites, and recruit the tri-snRNP, effectively performing all of the SR protein functions in one fell swoop. At the other extreme, each of the SR protein functions could be performed by individual SR protein molecules. Although these issues have not been fully answered, some recent experimental systems have allowed some of these questions to be addressed. Four papers contain results that together strongly indicate that multiple SR protein molecules participate in the removal of each intron (Graveley & Maniatis, 1998; Tacke et al., 1998; Xiao & Manley, 1998; Hertel & Maniatis, 1999). Specifically, these results suggest that each splicing event requires at least one SR protein molecule to perform a regulated exon-dependent function, and a second SR protein molecule to perform one or more of the other SR protein functions. Three of these papers have used pre-mRNAs containing SR protein-binding sites in the downstream exon. Thus, for the purposes of this discussion, the regulated exon-dependent functions of SR proteins will apply only to 3' splice site selection and not 5' splice site selection, which is likely to be constitutive in these cases.

Hybrid proteins consisting of the MS2 coat protein and an RS domain are sufficient to activate the splicing of enhancer-dependent pre-mRNAs containing an MS2 binding site in HeLa cell nuclear extracts (Graveley & Maniatis, 1998). In S100 extracts, however, both an SR protein and an MS2-RS protein, neither of which can function alone, are required for splicing (Graveley &

Maniatis, 1998). The MS2-RS proteins are probably only able to substitute for the regulated exon-dependent function of SR proteins for several reasons. First, because they can only bind to the MS2 site, their activity is spatially restricted. Second, the RS domain alone has been shown to be necessary but not sufficient to interact with the U1-70K protein (Xiao & Manley, 1997). Thus the MS2-RS proteins are probably unable to recruit U1 snRNP to the 5' splice site or mediate cross-intron interactions. The additional SR protein that is required in this situation is most likely performing one or both of these functions, and may also be involved in tri-snRNP incorporation. Thus, one interpretation of this experiment is that the MS2-RS proteins are performing the regulated exon-dependent function and the intact SR proteins are performing the constitutive exon-dependent and exon-independent functions (Fig. 5A). These results, as well as those described below, also indicate that the regulated exon-dependent functions of SR proteins can be uncoupled from the other SR protein functions.

The human homologs of the *Drosophila* Tra2 protein (hTra2 α and hTra2 β) function as activators of pre-mRNA splicing in vitro (Tacke et al., 1998). In S100 extracts, neither hTra2 α nor hTra2 β are sufficient to activate the splicing of a mouse IgM pre-mRNA containing a binding site for hTra2 as a splicing enhancer. However, the addition of either protein to a dilute nuclear extract, which alone does not support splicing, activates splicing of the IgM pre-mRNA containing the hTra2 splicing enhancer. These experiments are consistent with a model in which the activity of the hTra2 proteins is restricted to the regulated exon-dependent functions and the additional SR proteins present in the nuclear extract are required to perform the constitutive exon-dependent and exon-independent functions (Fig. 5B).

A study of the phosphorylation requirements for SR protein function has provided additional evidence that multiple SR proteins function in the splicing of each intron (Xiao & Manley, 1998). The HIV-1 *tat* pre-mRNA is known to contain SF2/ASF and to lack SC35-responsive elements (Fu, 1993; Mayeda et al., 1999). In S100 extracts, *tat* pre-mRNA splicing can be activated by the addition of SF2/ASF, but not SC35. Xiao and Manley (1998) have found that thiophosphorylated SF2/ASF is unable to activate *tat* pre-mRNA splicing in an S100 extract. However, they found that a combination of thiophosphorylated SF2/ASF and unmodified SC35 reconstituted *tat* pre-mRNA splicing in S100 extracts. Consistent with the model developed above, these results suggest that the thiophosphorylated SF2/ASF is performing the regulated exon-dependent functions and SC35 is performing the constitutive exon-dependent and exon-independent functions of SR proteins (Fig. 5C). Moreover, these results indicate that the regulated exon-dependent functions of SR

FIGURE 5. Summary of results indicating that multiple individual SR proteins participate in the removal of each intron. Throughout this figure the regulated exon-dependent functions of SR proteins in 3' splice site selection are illustrated as U2AF recruitment and the exon-independent functions are illustrated as cross-intron bridging and tri-snRNP recruitment. Note that the exon-dependent function of SR proteins in 5' splice site recognition may be occurring, but is not depicted in this illustration. **A:** MS2-RS proteins and SR proteins are required for enhancer-dependent splicing (Graveley & Maniatis, 1998). In this model, the MS2-RS protein is performing the regulated exon-dependent SR protein functions and the SR protein is performing the exon-independent and constitutive exon-dependent functions of SR proteins. **B:** hTra2 proteins and SR proteins are required for enhancer-dependent splicing (Tacke et al., 1998). In this model, hTra2 is performing the regulated exon-dependent functions and SR proteins are performing the exon-independent and constitutive exon-dependent functions. **C:** The phosphorylation requirements for the regulated exon-dependent functions are distinct from the other SR protein functions (Xiao & Manley, 1998). In this model, thiophosphorylated SF2/ASF is performing the regulated exon-dependent functions and SC35 is performing the exon-independent and constitutive exon-dependent functions. **D:** SR proteins have exon-independent functions (Hertel & Maniatis, 1999). In this model, a pre-mRNA substrate containing only 1 nt in the first exon and completely lacking the second exon can undergo the first step of splicing in a reaction that requires SR proteins. The SR protein in this case must be performing an exon-independent function.

proteins have phosphorylation requirements distinct from the other SR protein functions—the regulated exon-dependent functions do not require dephosphorylation whereas at least one of the other SR protein functions does.

The last experiments that support this model are those discussed earlier showing that a pre-mRNA containing only a single nucleotide in the first exon, and completely lacking the second exon, can undergo the first step of splicing (Hertel & Maniatis, 1999). The finding that SR proteins are required for the splicing of this minimal substrate demonstrates that SR proteins have an essential exon-independent function that, at least for some introns, is sufficient for splicing (Fig. 5D).

Taken together, this series of results provides very strong functional evidence that separate SR protein molecules perform the regulated exon-dependent functions and constitutive exon-dependent and exon-independent functions in the removal of each intron. This model can account for some of the apparent discrepancies in the literature. For instance, it was originally proposed that SR proteins were functionally redundant based on the observations that any individual SR protein could restore splicing in an S100 splicing-deficient extract (Fu et al., 1992; Mayeda et al., 1992; Zahler et al., 1992). Hertel and Maniatis (1999) have shown that the splicing of strong introns requires only the exon-independent functions of SR proteins and that all SR proteins tested for this activity were functionally redundant. Perhaps, then, only the exon-dependent functions of SR proteins will display SR protein specificity. This model could therefore account for earlier observations that SR proteins are functionally redundant in vitro.

This model can also explain some discrepancies in the interchangeability of RS domains. Although not discussed earlier, the RS domain of the large U2AF subunit can substitute for the C-terminal RS domain of Tra2 in *Drosophila*, and can function as a splicing activation domain in vitro when fused to the MS2 coat protein (B.R. Graveley, unpubl. data). In each case, the RNA binding domain of the fusion proteins restricts the activity of the RS domain to the regulated exon-dependent functions because Tra2 is a natural splicing regulator and MS2 is an artificial splicing regulator. In contrast, the U2AF⁶⁵ RS domain is unable to replace the SF2/ASF RS domain in vivo (DT40 chicken cells) or in vitro (Wang et al., 1998b). Although the in vivo result may be explained if the fusion protein failed to localize to the nucleus (which was not examined in this study), the in vitro results are not complicated by this possibility. Importantly, SF2/ASF can perform the exon-independent functions of SR proteins (Hertel & Maniatis, 1999). Therefore, the ability of the U2AF⁶⁵ RS domain to function when fused to Tra2 or MS2, but not when fused to SF2/ASF, may reflect differences in the RS domain sequence requirements for the regulated

exon-dependent and the exon-independent functions of SR proteins. This idea is consistent with the observation that thiophosphorylated SF2/ASF can perform the regulated exon-dependent functions of SR proteins, but is unable to perform the other SR protein functions (Xiao & Manley, 1998). Thus, the regulated exon-dependent and the other SR protein functions not only have different RS domain sequence requirements, but they also have different phosphorylation requirements.

CONCLUDING REMARKS

The last few years have seen several advances in our understanding of SR protein functions. The finding that SR proteins bind to both constitutive and alternative exons provides a nice biological explanation for their rather promiscuous binding specificities. The technological advances that have allowed for the uncoupling of SR protein activities have led to the discovery that different functional RS domain-mediated protein interactions have distinct protein sequence and phosphorylation requirements, and that multiple independent SR proteins participate in the removal of each intron. Hopefully, work over the next few years will be equally fruitful and will help to further sort through some of the confusing areas in the field.

ACKNOWLEDGMENTS

I would like to thank Javier Cáceres, Adrian Krainer, and Juan Valcárcel for communicating results prior to publication. In addition, I would like to thank Javier Cáceres, Klemens Hertel, Kristen Lynch, Tom Maniatis, and members of my laboratory for their useful comments on the manuscript.

Received May 11, 2000; returned for revision June 9, 2000; revised manuscript received June 23, 2000

REFERENCES

- Abovich N, Rosbash M. 1997. Cross-intron bridging interactions in the yeast commitment complex are conserved in mammals. *Cell* 89:403–412.
- Amrein H, Gorman M, Nothiger R. 1988. The sex-determining gene tra-2 of *Drosophila* encodes a putative RNA binding protein. *Cell* 55:1025–1035.
- Amrein H, Hedley ML, Maniatis T. 1994. The role of specific protein–RNA and protein–protein interactions in positive and negative control of pre-mRNA splicing by Transformer 2. *Cell* 76:735–746.
- Anderson K, Moore MJ. 1997. Bimolecular exon ligation by the human spliceosome. *Science* 276:1712–1716.
- Ayane M, Preuss U, Kohler G, Nielsen PJ. 1991. A differentially expressed murine RNA encoding a protein with similarities to two types of nucleic acid binding motifs. *Nucleic Acids Res* 19:1273–1278.
- Baker BS. 1989. Sex in flies: The splice of life. *Nature* 340:521–524.
- Beil B, Sreaton G, Stamm S. 1997. Molecular cloning of htra2-beta-1 and htra2-beta-2, two human homologs of tra-2 generated by alternative splicing. *DNA Cell Biol* 16:679–690.

- Bennett M, Michaud S, Kingston J, Reed R. 1992. Protein components specifically associated with prespliceosome and spliceosome complexes. *Genes & Dev* 6:1986–2000.
- Berget SM. 1995. Exon recognition in vertebrate splicing. *J Biol Chem* 270:2411–2414.
- Blencowe BJ. 2000. Exonic splicing enhancers: Mechanism of action, diversity and role in human genetic diseases. *Trends Biochem Sci* 25:106–110.
- Blencowe BJ, Bauren G, Eldridge AG, Issner R, Nickerson JA, Rosonina E, Sharp PA. 2000. The SRm160/300 splicing coactivator subunits. *RNA* 6:111–120.
- Blencowe BJ, Bowman JAL, McCracken S, Ronina E. 1999. SR-related proteins and the processing of messenger RNA precursors. *Biochem Cell Biol* 77:277–291.
- Blencowe BJ, Issner R, Nickerson JA, Sharp PA. 1998. A coactivator of pre-mRNA splicing. *Genes & Dev* 12:996–1009.
- Boggs RT, Gregor P, Idriss S, Belote JM, McKeown M. 1987. Regulation of sexual differentiation in *D. melanogaster* via alternative splicing of RNA from the transformer gene. *Cell* 50:739–747.
- Bouck J, Fu X-D, Skalka AM, Katz RA. 1998. Role of the constitutive splicing factors U2AF65 and SAP49 in suboptimal RNA splicing of novel retroviral mutants. *J Biol Chem* 273:15169–15176.
- Bourgeois CF, Popielarz M, Hildwein G, Stevenin J. 1999. Identification of a bidirectional splicing enhancer: Differential involvement of SR proteins in 5' and 3' splice site activation. *Mol Cell Biol* 19:7347–7356.
- Bruzik JP, Maniatis T. 1995. Enhancer-dependent interaction between 5' and 3' splice sites in trans. *Proc Natl Acad Sci USA* 92:7056–7059.
- Burge CB, Tuschl T, Sharp PA. 1999. Splicing of precursors to mRNAs by the spliceosomes. In: Gesteland RF, Cech TR, Atkins JF, eds., *The RNA world*, 2nd ed. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press. pp 525–560.
- Burtis KC, Baker BS. 1989. *Drosophila doublesex* gene controls somatic sexual differentiation by producing alternatively spliced mRNAs encoding related sex-specific polypeptides. *Cell* 56:997–1010.
- Caceres JF, Krainer AR. 1993. Functional analysis of pre-mRNA splicing factor SF2/ASF structural domains. *EMBO J* 12:4715–4726.
- Caceres JF, Screaton GR, Krainer AR. 1998. A specific subset of SR proteins shuttles continuously between the nucleus and the cytoplasm. *Genes & Dev* 12:55–66.
- Caceres JF, Stamm S, Helfman DM, Krainer AR. 1994. Regulation of alternative splicing in vivo by overexpression of antagonistic splicing factors. *Science* 265:1706–1709.
- Cavaloc Y, Bourgeois CF, Kister L, Stevenin J. 1999. The splicing factors 9G8 and SRp20 transactivate splicing through different and specific enhancers. *RNA* 5:468–483.
- Cavaloc Y, Popielarz M, Fuchs JP, Gattoni R, Stevenin J. 1994. Characterization and cloning of the human splicing factor 9G8: A novel 35 kDa factor of the serine/arginine protein family. *EMBO J* 13:2639–2649.
- Champlin DT, Frasch M, Saumweber H, Lis JT. 1991. Characterization of a *Drosophila* protein associated with boundaries of transcriptionally active chromatin. *Genes & Dev* 5:1611–1621.
- Chandler SD, Mayeda A, Yeakley JM, Krainer AR, Fu X-D. 1997. RNA splicing specificity determined by the coordinated action of RNA recognition motifs in SR proteins. *Proc Natl Acad Sci USA* 94:3596–3601.
- Chew SL, Liu HX, Mayeda A, Krainer AR. 1999. Evidence for the function of an exonic splicing enhancer after the first catalytic step of pre-mRNA splicing. *Proc Natl Acad Sci USA* 96:10655–10660.
- Chiara MD, Reed R. 1995. A two-step mechanism for 5' and 3' splice-site pairing. *Nature* 375:510–513.
- Colwill K, Pawson T, Andrews B, Prasad J, Manley JL, Bell JC, Duncan PI. 1996. The Clk/Sty protein kinase phosphorylates SR splicing factors and regulates their intranuclear distribution. *EMBO J* 15:265–275.
- Côté J, Simard MJ, Chabot B. 1999. An element in the 5' common exon of the NCAM alternative splicing unit interacts with SR proteins and modulates 5' splice site selection. *Nucleic Acids Res* 27:2529–2537.
- Crispino JD, Blencowe BJ, Sharp PA. 1994. Complementation by SR proteins of pre-mRNA splicing reactions depleted of U1 snRNP. *Science* 265:1866–1869.
- Crispino JD, Sharp PA. 1995. A U6 snRNA:pre-mRNA interaction can be rate-limiting for U1-independent splicing. *Genes & Dev* 9:2314–2323.
- Dauwalder B, Amaya-Manzanares F, Mattox W. 1996. A human homologue of the *Drosophila* sex determination factor transformer-2 has conserved splicing regulatory functions. *Proc Natl Acad Sci USA* 93:9004–9009.
- Dauwalder B, Mattox W. 1998. Analysis of the functional specificity of RS domains in vivo. *EMBO J* 17:6049–6060.
- Denhez F, Lafyatis R. 1994. Conservation of regulated alternative splicing and identification of functional domains in vertebrate homologs to the *Drosophila* splicing regulator, suppressor-of-white-apricot. *J Biol Chem* 269:16170–16179.
- Diamond RH, Du K, Lee VM, Mohn KL, Haber BA, Tewari DS, Taub R. 1993. Novel delayed-early and highly insulin-induced growth response genes. Identification of HRS, a potential regulator of alternative pre-mRNA splicing. *J Biol Chem* 268:15185–15192.
- Du C, McGuffin ME, Dauwalder B, Rabinow L, Mattox W. 1998. Protein phosphorylation plays an essential role in the regulation of alternative splicing and sex determination in *Drosophila*. *Mol Cell* 2:741–750.
- Eldridge AG, Li Y, Sharp PA, Blencowe BJ. 1999. The SRm160/300 splicing coactivator is required for exon-enhancer function. *Proc Natl Acad Sci USA* 96:6125–6130.
- Fetzer S, Lauber J, Will CL, Lüthmann R. 1997. The [U4/U6.U5] tri-snRNP-specific 27K protein is a novel SR protein that can be phosphorylated by the snRNP-associated protein kinase. *RNA* 3:344–355.
- Fortes P, Bilbao-Cortes D, Fornerod M, Rigaut G, Raymond W, Séraphin B, Mattaj JW. 1999. Luc7p, a novel yeast U1 snRNP protein with a role in 5' splice site selection. *Genes & Dev* 13:2425–2438.
- Fu X-D. 1993. Specific commitment of different pre-mRNAs to splicing by single SR proteins. *Nature* 365:82–85.
- Fu X-D. 1995. The superfamily of arginine/serine-rich splicing factors. *RNA* 1:663–680.
- Fu XD, Maniatis T. 1990. Factor required for mammalian spliceosome assembly is localized to discrete regions in the nucleus. *Nature* 343:437–441.
- Fu XD, Maniatis T. 1992a. The 35-kDa mammalian splicing factor SC35 mediates specific interactions between U1 and U2 small nuclear ribonucleoprotein particles at the 3' splice site. *Proc Natl Acad Sci USA* 89:1725–1729.
- Fu XD, Maniatis T. 1992b. Isolation of a complementary DNA that encodes the mammalian splicing factor SC35. *Science* 256:535–538.
- Fu XD, Mayeda A, Maniatis T, Krainer AR. 1992. General splicing factors SF2 and SC35 have equivalent activities in vitro, and both affect alternative 5' and 3' splice site selection. *Proc Natl Acad Sci USA* 89:11224–11228.
- Ge H, Manley JL. 1990. A protein factor, ASF, controls cell-specific alternative splicing of SV40 early pre-mRNA in vitro. *Cell* 62:25–34.
- Ge H, Zuo P, Manley JL. 1991. Primary structure of the human splicing factor ASF reveals similarities with *Drosophila* regulators. *Cell* 66:373–382.
- Golovkin M, Reddy AS. 1998. The plant U1 small nuclear ribonucleoprotein particle 70K protein interacts with two novel serine/arginine-rich proteins. *Plant Cell* 10:1637–1648.
- Golovkin M, Reddy AS. 1999. An SC35-like protein and a novel serine/arginine-rich protein interact with *Arabidopsis* U1-70K protein. *J Biol Chem* 274:36428–36438.
- Graveley BR, Hertel KJ, Maniatis T. 1998. A systematic analysis of the factors that determine the strength of pre-mRNA splicing enhancers. *EMBO J* 17:6747–6756.
- Graveley BR, Maniatis T. 1998. Arginine/serine-rich domains of SR proteins can function as activators of pre-mRNA splicing. *Mol Cell* 1:765–771.
- Gross T, Richert K, Mierke C, Lutzelberger M, Kaufer NF. 1998. Identification and characterization of srp1, a gene of fission yeast encoding a RNA binding domain and a RS domain typical of SR splicing factors. *Nucleic Acids Res* 26:505–511.
- Gui JF, Lane WS, Fu XD. 1994a. A serine kinase regulates intracellular localization of splicing factors in the cell cycle. *Nature* 369:678–682.

- Gui JF, Tronchere H, Chandler SD, Fu XD. 1994b. Purification and characterization of a kinase specific for the serine- and arginine-rich pre-mRNA splicing factors. *Proc Natl Acad Sci USA* 91:10824–10828.
- Hanes J, von der Kammer H, Klaudiny J, Scheit KH. 1994. Characterization by cDNA cloning of two new human protein kinases. Evidence by sequence comparison of a new family of mammalian protein kinases. *J Mol Biol* 244:665–672.
- Hertel KJ, Lynch KW, Maniatis T. 1997. Common themes in the function of transcription and splicing enhancers. *Curr Opin Cell Biol* 9:350–357.
- Hertel KJ, Maniatis T. 1999. Serine-arginine (SR)-rich splicing factors have an exon-independent function in pre-mRNA splicing. *Proc Natl Acad Sci USA* 96:2651–2655.
- Jamison SF, Pasman Z, Wang J, Will C, Lührmann R, Manley JL, Garcia-Blanco MA. 1995. U1 snRNP-ASF/SF2 interaction and 5' splice site recognition: Characterization of required elements. *Nucleic Acids Res* 23:3260–3267.
- Johnson KW, Smith KA. 1991. Molecular cloning of a novel human cdc2/CDC28-like protein kinase. *J Biol Chem* 266:3402–3407.
- Kan JLC, Green MR. 1999. Pre-mRNA Splicing of IgM exons M1 and M2 is directed by a juxtaposed splicing enhancer and inhibitor. *Genes & Dev* 13:462–471.
- Kanopka A, Muhlemann O, Petersen-Mahrt S, Estmer C, Ohrmalm C, Akusjarvi G. 1998. Regulation of adenovirus alternative RNA splicing by dephosphorylation of SR proteins. *Nature* 393:185–187.
- Kohtz JD, Jamison SF, Will CL, Zuo P, Lührmann R, Garcia-Blanco MA, Manley JL. 1994. Protein-protein interactions and 5' splice site recognition in mammalian mRNA precursors. *Nature* 368:119–124.
- Krainer AR, Conway GC, Kozak D. 1990a. The essential pre-mRNA splicing factor SF2 influences 5' splice site selection by activating proximal sites. *Cell* 62:35–42.
- Krainer AR, Conway GC, Kozak D. 1990b. Purification and characterization of pre-mRNA splicing factor SF2 from HeLa cells. *Genes & Dev* 4:1158–1171.
- Krainer AR, Mayeda A, Kozak D, Binns G. 1991. Functional expression of cloned human splicing factor SF2: Homology to RNA-binding proteins, U1 70K, and *Drosophila* splicing regulators. *Cell* 66:383–394.
- Labourier E, Bourbon H-M, Gallouzi I-E, Fostier M, Allemand E, Tazi J. 1999. Antagonism between RSF1 and SR proteins for both splice-site recognition in vitro and *Drosophila* development. *Genes & Dev* 13:740–753.
- Lavigne A, La Branche H, Dornblüth AR, Chabot B. 1993. A splicing enhancer in the human fibronectin alternate ED1 exon interacts with SR proteins and stimulates U2 snRNP binding. *Genes & Dev* 7:2405–2417.
- Lazar G, Schaal T, Maniatis T, Goodman HM. 1995. Identification of a plant serine-arginine-rich protein similar to the mammalian splicing factor SF2/ASF. *Proc Natl Acad Sci USA* 92:7672–7676.
- Li H, Bingham PM. 1991. Arginine/serine-rich domains of the su(wa) and tra RNA processing regulators target proteins to a subnuclear compartment implicated in splicing. *Cell* 67:335–342.
- Li Y, Blencowe BJ. 1999. Distinct factor requirements for exonic splicing enhancer function and binding of U2AF to the polypyrimidine tract. *J Biol Chem* 274:35074–35079.
- Liu H-X, Zhang M, Krainer AR. 1998. Identification of functional exonic splicing enhancer motifs recognized by individual SR proteins. *Genes & Dev* 12:1998–2012.
- Liu HX, Chew SL, Cartegni L, Zhang MQ, Krainer AR. 2000. Exonic splicing enhancer motif recognized by human SC35 under splicing conditions. *Mol Cell Biol* 20:1063–1071.
- Longman D, Johnstone IL, Cáceres JF. 2000. Functional characterization of SR and SR-related genes in *Caenorhabditis elegans*. *EMBO J* 19:1625–1637.
- Lopato S, Gattoni R, Fabini G, Stevenin J, Barta A. 1999. A novel family of plant splicing factors with a Zn knuckle motif: Examination of RNA binding and splicing activities. *Plant Mol Biol* 39:761–773.
- Lopato S, Mayeda A, Krainer AR, Barta A. 1996a. Pre-mRNA splicing in plants: Characterization of Ser/Arg splicing factors. *Proc Natl Acad Sci USA* 93:3074–3079.
- Lopato S, Waigmann E, Barta A. 1996b. Characterization of a novel arginine/serine-rich splicing factor in *Arabidopsis*. *Plant Cell* 8:2255–2264.
- Lorson CL, Androphy EJ. 2000. An exonic enhancer is required for inclusion of an essential exon in the SMA-determining gene SMN. *Hum Mol Genet* 9:259–265.
- Lutzelberger M, Gross T, Kaufer NF. 1999. Srp2, an SR protein family member of fission yeast: In vivo characterization of its modular domains. *Nucleic Acids Res* 27:2618–2626.
- Lynch KW, Maniatis T. 1996. Assembly of specific SR protein complexes on distinct regulatory elements of the *Drosophila* double-sex splicing enhancer. *Genes & Dev* 10:2089–2101.
- Mayeda A, Sreaton GR, Chandler SD, Fu XD, Krainer AR. 1999. Substrate specificities of SR proteins in constitutive splicing are determined by their RNA recognition motifs and composite pre-mRNA exonic elements. *Mol Cell Biol* 19:1853–1863.
- Mayeda A, Zahler AM, Krainer AR, Roth MB. 1992. Two members of a conserved family of nuclear phosphoproteins are involved in pre-mRNA splicing. *Proc Natl Acad Sci USA* 89:1301–1304.
- Merendino L, Guth S, Bilbao D, Martinez C, Valcarcel J. 1999. Inhibition of msl-2 splicing by Sex-lethal reveals interaction between U2AF35 and the 3' splice site AG. *Nature* 402:838–841.
- Michaud S, Reed R. 1991. An ATP-independent complex commits pre-mRNA to the mammalian spliceosome assembly pathway. *Genes & Dev* 5:2534–2546.
- Misteli T, Cáceres JF, Clement JQ, Krainer AR, Wilkinson MF, Spector DL. 1998. Serine phosphorylation of SR proteins is required for their recruitment to sites of transcription in vivo. *J Cell Biol* 143:297–307.
- Misteli T, Cáceres JF, Spector DL. 1997. The dynamics of a pre-mRNA splicing factor in living cells. *Nature* 387:523–527.
- Misteli T, Spector DL. 1998. The cellular organization of gene expression. *Curr Opin Cell Biol* 10:323–331.
- Misteli T, Spector DL. 1999. RNA polymerase II targets pre-mRNA splicing factors to transcription sites in vivo. *Mol Cell* 3:697–705.
- Nishii Y, Morishima M, Kakehi Y, Umehara K, Kioka N, Terano Y, Amachi T, Ueda K. 2000. CROP/Luc7A, a novel serine/arginine-rich protein, isolated from cisplatin-resistant cell line. *FEBS Lett* 465:153–165.
- Ono Y, Ohno M, Shimura Y. 1994. Identification of a putative RNA helicase (HRH1), a human homolog of yeast Prp22. *Mol Cell Biol* 14:7611–7620.
- Prasad J, Colwill K, Pawson T, Manley JL. 1999. The protein kinase Clk/Sty directly modulates SR protein activity: Both hyper- and hypophosphorylation inhibit splicing. *Mol Cell Biol* 19:6991–7000.
- Query CC, Bentley RC, Keene JD. 1989. A common RNA recognition motif identified within a defined U1 RNA binding domain of the 70K U1 snRNP protein. *Cell* 57:89–101.
- Reed R. 1996. Initial splice-site recognition and pairing during pre-mRNA splicing. *Curr Opin Genet Dev* 6:215–220.
- Reed R, Palandjian L. 1997. Spliceosome assembly. In: Krainer AR, ed., *Eukaryotic mRNA processing*. Oxford: IRL Press. pp 103–129.
- Ring HZ, Lis JT. 1994. The SR protein B52/SRp55 is essential for *Drosophila* development. *Mol Cell Biol* 14:7499–7506.
- Roscigno RF, Garcia-Blanco MA. 1995. SR proteins escort the U4/U6•U5 tri-snRNP to the spliceosome. *RNA* 1:692–706.
- Rossi F, Labourier E, Forne T, Divita G, Derancourt J, Riou JF, Antoine E, Cathala G, Brunel C, Tazi J. 1996. Specific phosphorylation of SR proteins by mammalian DNA topoisomerase I. *Nature* 381:80–82.
- Roth MB, Zahler AM, Stolk JA. 1991. A conserved family of nuclear phosphoproteins localized to sites of polymerase II transcription. *J Cell Biol* 115:587–596.
- Ruskin B, Zamore PD, Green MR. 1988. A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. *Cell* 52:207–219.
- Ryner LC, Goodwin SF, Castrillon DH, Anand A, Vilella A, Baker BS, Hall JC, Taylor BJ, Wasserman SA. 1996. Control of male sexual behavior and sexual orientation in *Drosophila* by the fruitless gene. *Cell* 87:1079–1089.
- Sanford JR, Bruzik JP. 1999. Developmental regulation of SR protein phosphorylation and activity. *Genes & Dev* 13:1513–1518.
- Schaal TD, Maniatis T. 1999a. Multiple distinct splicing enhancers in the protein-coding sequences of a constitutively spliced pre-mRNA. *Mol Cell Biol* 19:261–273.

- Schaal TD, Maniatis T. 1999b. Selection and characterization of pre-mRNA splicing enhancers: Identification of novel SR protein-specific enhancer sequences. *Mol Cell Biol* 19:1705–1719.
- Screaton GR, Cáceres JF, Mayeda A, Bell MV, Plebanski M, Jackson DG, Bell JI, Krainer AR. 1995. Identification and characterization of three members of the human SR family of pre-mRNA splicing factors. *EMBO J* 14:4336–4349.
- Selvakumar M, Helfman DM. 1999. Exonic splicing enhancers contribute to the use of both 3' and 5' splice site usage of rat beta-tropomyosin pre-mRNA. *RNA* 5:378–394.
- Shi H, Hoffman BE, Lis JT. 1997. A specific RNA hairpin loop structure binds the RNA recognition motifs of the *Drosophila* SR protein B52. *Mol Cell Biol* 17:2649–2657.
- Smith CW, Porro EB, Patton JG, Nadal-Ginard B. 1989. Scanning from an independently specified branch point defines the 3' splice site of mammalian introns. *Nature* 342:243–247.
- Soret J, Gattoni R, Guyon C, Sureau A, Popielarz M, Le Rouzic E, Dumon S, Apiou F, Dutrillaux B, Voss H, Ansorge W, Stevenin J, Perbal B. 1998. Characterization of SRp46, a novel human SR splicing factor encoded by a PR264/SC35 retropseudogene. *Mol Cell Biol* 18:4924–4934.
- Spikes DA, Kramer J, Bingham PM, Van Doren K. 1994. SWAP pre-mRNA splicing regulators are a novel, ancient protein family sharing a highly conserved sequence motif with the prp21 family of constitutive splicing proteins. *Nucleic Acids Res* 22:4510–4519.
- Sukegawa J, Blobel G. 1995. A putative mammalian RNA helicase with an arginine-serine-rich domain colocalizes with a splicing factor. *J Biol Chem* 270:15702–15706.
- Sun Q, Mayeda A, Hampson RK, Krainer AR, Rottman FM. 1993. General splicing factor SF2/ASF promotes alternative splicing by binding to an exonic splicing enhancer. *Genes & Dev* 7:2598–2608.
- Tacke R, Chen Y, Manley JL. 1997. Sequence-specific RNA binding by an SR protein requires RS domain phosphorylation: Creation of an SRp40-specific splicing enhancer. *Proc Natl Acad Sci USA* 94:1148–1153.
- Tacke R, Manley JL. 1995. The human splicing factors ASF/SF2 and SC35 possess distinct, functionally significant RNA binding specificities. *EMBO J* 14:3540–3551.
- Tacke R, Manley JL. 1999. Determinants of SR protein specificity. *Curr Opin Cell Biol* 11:358–362.
- Tacke R, Tohyama M, Ogawa S, Manley JL. 1998. Human Tra2 proteins are sequence-specific activators of pre-mRNA splicing. *Cell* 93:139–148.
- Tarn WY, Steitz JA. 1994. SR proteins can compensate for the loss of U1 snRNP functions in vitro. *Genes & Dev* 8:2704–2717.
- Tarn WY, Steitz JA. 1995. Modulation of 5' splice site choice in pre-messenger RNA by two distinct steps. *Proc Natl Acad Sci USA* 92:2504–2508.
- Teigelkamp S, Mundt C, Achsel T, Will CL, Lührmann R. 1997. The human U5 snRNP-specific 100-kD protein is an RS domain-containing, putative RNA helicase with significant homology to the yeast splicing factor Prp28p. *RNA* 3:1313–1326.
- Tian M, Maniatis T. 1992. Positive control of pre-mRNA splicing in vitro. *Science* 256:237–240.
- Tian M, Maniatis T. 1993. A splicing enhancer complex controls alternative splicing of doublesex pre-mRNA. *Cell* 74:105–114.
- Tian M, Maniatis T. 1994. A splicing enhancer exhibits both constitutive and regulated activities. *Genes & Dev* 8:1703–1712.
- Wang H-Y, Lin W, Dyck JA, Yeakley JM, Songyang Z, Cantley LC, Fu X-D. 1998a. SRPK2: A differentially expressed SR protein-specific kinase involved in mediating the interaction and localization of pre-mRNA splicing factors in mammalian cells. *J Cell Biol* 140:737–750.
- Wang J, Manley JL. 1995. Overexpression of the SR proteins ASF/SF2 and SC35 influences alternative splicing in vivo in diverse ways. *RNA* 1:335–346.
- Wang J, Takagaki Y, Manley JL. 1996. Targeted disruption of an essential vertebrate gene: ASF/SF2 is required for cell viability. *Genes & Dev* 10:2588–2599.
- Wang J, Xiao SH, Manley JL. 1998b. Genetic analysis of the SR protein ASF/SF2: Interchangeability of RS domains and negative control of splicing. *Genes & Dev* 12:2222–2233.
- Wang Z, Hoffmann HM, Grabowski PJ. 1995. Intrinsic U2AF binding is modulated by exon enhancer signals in parallel with changes in splicing activity. *RNA* 1:21–35.
- Wu JY, Maniatis T. 1993. Specific interactions between proteins implicated in splice site selection and regulated alternative splicing. *Cell* 75:1061–1070.
- Wu S, Romfo CM, Nilsen TW, Green MR. 1999. Functional recognition of the 3' splice site AG by the splicing factor U2AF35. *Nature* 402:832–835.
- Xiao S-H, Manley JL. 1997. Phosphorylation of the ASF/SF2 RS domain affects both protein-protein interactions and is necessary for splicing. *Genes & Dev* 11:334–344.
- Xiao S-H, Manley JL. 1998. Phosphorylation-dephosphorylation differentially affects activities of splicing factor ASF/SF2. *EMBO J* 17:6359–6367.
- Zahler AM, Lane WS, Stolk JA, Roth MB. 1992. SR proteins: A conserved family of pre-mRNA splicing factors. *Genes & Dev* 6:837–847.
- Zahler AM, Neugebauer KM, Lane WS, Roth MB. 1993a. Distinct functions of SR proteins in alternative pre-mRNA splicing. *Science* 260:219–222.
- Zahler AM, Neugebauer KM, Stolk JA, Roth MB. 1993b. Human SR proteins and isolation of a cDNA encoding SRp75. *Mol Cell Biol* 13:4023–4028.
- Zamore PD, Green MR. 1989. Identification, purification, and biochemical characterization of U2 small nuclear ribonucleoprotein auxiliary factor. *Proc Natl Acad Sci USA* 86:9243–9247.
- Zamore PD, Patton JG, Green MR. 1992. Cloning and domain structure of the mammalian splicing factor U2AF. *Nature* 355:609–614.
- Zhang M, Zamore PD, Carmo-Fonseca M, Lamond AI, Green MR. 1992. Cloning and intracellular localization of the U2 small nuclear ribonucleoprotein auxiliary factor small subunit. *Proc Natl Acad Sci USA* 89:8769–8773.
- Zhang W-J, Wu JY. 1996. Functional properties of p54, a novel SR protein active in constitutive and alternative splicing. *Mol Cell Biol* 16:5400–5408.
- Zhou Z, Reed R. 1998. Human homologs of yeast prp16 and prp17 reveal conservation of the mechanism for catalytic step II of pre-mRNA splicing. *EMBO J* 17:2095–2106.
- Zorio DA, Blumenthal T. 1999. Both subunits of U2AF recognize the 3' splice site in *Caenorhabditis elegans*. *Nature* 402:835–838.
- Zuo P, Maniatis T. 1996. The splicing factor U2AF³⁵ mediates critical protein-protein interactions in constitutive and enhancer-dependent splicing. *Genes & Dev* 10:1356–1368.
- Zuo P, Manley JL. 1993. Functional domains of the human splicing factor ASF/SF2. *EMBO J* 12:4727–4737.
- Zuo P, Manley JL. 1994. The human splicing factor ASF/SF2 can specifically recognize pre-mRNA 5' splice sites. *Proc Natl Acad Sci USA* 91:3363–3367.